



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### **Usage guidelines**

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



*8PT*  
*miss*

THE STANFORD  
COMPUTATION  
CENTER



Gift of  
Captain T. J. J. See

MATHEMATICAL  
SCIENCES  
LIBRARY



J. J. SEE

S. D. Lombard  
42 Ferkins Hall,  
Cambridge, Mass.

$$Z = r e^{i\theta} \text{ sec p. 20.}$$

$$\text{If } r = 1,$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^{i\frac{\pi}{2}} = \cos \frac{\pi}{2} + i \sin \frac{\pi}{2} = i.$$

Take the  $i$ th root:

$$e^{\frac{\pi}{2}} = \sqrt{-1} = \sqrt{i}, \text{ which is:}$$

$$e = 2.7182818284$$

$$\log e^{\frac{\pi}{2}} = \log e = 0.4342971 \dots 1.570796$$

$$\log \log e = 9.6377843$$

$$\log \frac{\pi}{2} = 0.1761199$$

$$\log (\log e^{\frac{\pi}{2}}) = 9.8339042$$

$$\log e^{\frac{\pi}{2}} = 0.6621883$$

$$e^{\frac{\pi}{2}} = 4.811479 \dots = \sqrt{i}$$

Q. S. D.

Also:

$$z = r e^{i\phi} = e^{i\pi}$$

$$\text{If } \phi = \pi,$$

$$r e^{i\phi} = e^{i\pi} = -1$$

$$\phi = \pi.$$

$$\phi = \pi,$$

$$r(-1), e^{i\pi} = -1.$$

$$1 = 0, \text{ End } \dots \text{ P. S. D.}$$





**T. J. J. SEE**  
**MARE ISLAND, CALIF.**

# **ELEMENTS**

**OF THE**

# **INTEGRAL CALCULUS,**

**WITH A**

***KEY TO THE SOLUTION OF DIFFERENTIAL  
EQUATIONS, AND A SHORT TABLE  
OF INTEGRALS.***

**BY**

**WILLIAM ELWOOD BYERLY, PH.D.,**  
**PROFESSOR OF MATHEMATICS IN HARVARD UNIVERSITY.**

---

***SECOND EDITION, REVISED AND ENLARGED.***

---

**BOSTON, U.S.A.:**  
**PUBLISHED BY GINN AND COMPANY.**  
**1895.**

517.3  
B993  
ed. 2  
c. 2

Entered according to Act of Congress, in the year 1888, by  
WILLIAM ELWOOD BYERLY,  
in the Office of the Librarian of Congress at Washington.

---

ALL RIGHTS RESERVED.

TYPOGRAPHY BY J. S. CUSHING & Co., BOSTON, U.S.A.

---

PRESSWORK BY GINN & Co., BOSTON, U.S.A.



## PREFACE.

---

THE following volume is a sequel to my treatise on the Differential Calculus, and, like that, is written as a text-book. The last chapter, however, a Key to the Solution of Differential Equations, may prove of service to working mathematicians.

I have used freely the works of Bertrand, Benjamin Peirce, Todhunter, and Boole; and I am much indebted to Professor J. M. Peirce for criticisms and suggestions.

I refer constantly to my work on the Differential Calculus as Volume I.; and for the sake of convenience I have added Chapter V. of that book, which treats of Integration, as an appendix to the present volume.

W. E. BYERLY.

CAMBRIDGE, 1881.



## PREFACE TO SECOND EDITION.

---

IN enlarging my Integral Calculus I have used freely Schlömilch's "Compendium der Höheren Analysis," Cayley's "Elliptic Functions," Meyer's "Bestimmte Integrale," Forsyth's "Differential Equations," and Williamson's "Integral Calculus."

The chapter on Theory of Functions was sketched out and in part written by Professor B. O. Peirce, to whom I am greatly indebted for numerous valuable suggestions touching other portions of the book, and who has kindly allowed me to have his Short Table of Integrals bound in with this volume.

W. E. BYERLY.

CAMBRIDGE, 1888.

## ANALYTICAL TABLE OF CONTENTS.

### CHAPTER I.

#### SYMBOLS OF OPERATION.

Article.	Page.
1. Functional symbols regarded as <i>symbols of operation</i> . . . . .	1
2. Compound function; compound operation . . . . .	1
3. Commutative or relatively free operations . . . . .	1
4. Distributive or linear operations . . . . .	2
5. The compounds of <i>distributive</i> operations are distributive . . . . .	2
6. Symbolic exponents . . . . .	2
7. The <i>law of indices</i> . . . . .	2
8. The interpretation of a zero exponent . . . . .	3
9. The interpretation of a negative exponent . . . . .	3
10. When operations are commutative and distributive, the symbols which represent them may be combined as if they were algebraic quantities . . . . .	3

### CHAPTER II.

#### IMAGINARIES.

11. Usual definition of an <i>imaginary</i> . Imaginaries first forced upon our attention in connection with quadratic equations . . . . .	5
12. Treatment of imaginaries purely arbitrary and conventional . . . . .	6
13. $\sqrt{-1}$ defined as a symbol of operation . . . . .	6
14. The rules in accordance with which the symbol $\sqrt{-1}$ is used. $\sqrt{-1}$ <i>distributive</i> and <i>commutative</i> with symbols of quantity . . . . .	7
15. Interpretation of powers of $\sqrt{-1}$ . . . . .	7
16. Imaginary roots of a quadratic . . . . .	8
17. Typical form of an imaginary. Equal imaginaries . . . . .	8
18. Geometrical representation of an imaginary. <i>Reals</i> and <i>pure imaginaries</i> . An interpretation of the operation $\sqrt{-1}$ . . . . .	8
19. The <i>sum</i> , the <i>product</i> , and the <i>quotient</i> of two imaginaries, $a + b\sqrt{-1}$ and $c + d\sqrt{-1}$ , are imaginaries of the typical form . . . . .	10

Article	Page
20. Second typical form $r(\cos \phi + \sqrt{-1} \sin \phi)$ . <i>Modulus and argument. Absolute value of an imaginary. Examples</i> . . . . .	10
21. The modulus of the sum of two imaginaries is never greater than the sum of their moduli . . . . .	11
22. Modulus and argument of the product of imaginaries . . . . .	12
23. Modulus and argument of the quotient of two imaginaries . . . . .	13
24. Modulus and argument of a power of an imaginary . . . . .	13
25. Modulus and argument of a root of an imaginary. Example . . . . .	14
26. Relation between the $n$ th roots of a real or an imaginary . . . . .	14
27. The imaginary roots of 1 and $-1$ . Examples . . . . .	15
28. <i>Conjugate</i> imaginaries. Examples . . . . .	17
29. <i>Transcendental</i> functions of an imaginary variable best defined by the aid of series . . . . .	17
30. Convergency of a series containing imaginary terms . . . . .	18
31. <i>Exponential</i> functions of an imaginary. Definition of $e^z$ where $z$ is imaginary . . . . .	19
32. The <i>law of indices</i> holds for imaginary exponentials. Example . . . . .	20
33. <i>Logarithmic</i> functions of an imaginary. Definition of $\log z$ . $\log z$ a <i>periodic</i> function. Example . . . . .	21
34. <i>Trigonometric</i> functions of an imaginary. Definition of $\sin z$ and $\cos z$ . Example . . . . .	22
35. $\sin z$ and $\cos z$ expressed in <i>exponential</i> form. The fundamental formulas of Trigonometry hold for imaginaries as well as for reals. Examples . . . . .	22
36. <i>Differentiation</i> of Functions of Imaginary Variables. The <i>derivative</i> of a function of an imaginary is in general indeterminate . . . . .	24
37. In differentiating, we may treat the $\sqrt{-1}$ like a constant factor. Example. Two forms of the differential of the independent variable . . . . .	24
38. Differentiation of a power of $z$ . . . . .	25
39. Differentiation of $e^z$ . Example . . . . .	26
40. Differentiation of $\log z$ . . . . .	26
41. Differentiation of $\sin z$ and $\cos z$ . . . . .	26
42. Formulas for <i>direct integration</i> (I., Art. 74) hold when $z$ is imaginary . . . . .	27
43. <i>Hyperbolic Functions</i> . . . . .	27
44. Examples. Properties of Hyperbolic Functions . . . . .	28
45. Differentiation of Hyperbolic Functions . . . . .	28
46. Anti-hyperbolic functions. Examples . . . . .	28
47. Anti-hyperbolic functions expressed as logarithms . . . . .	29
48. Formulas for the direct integration of some irrational forms . . . . .	30



TABLE OF CONTENTS.

vii

CHAPTER III.

GENERAL METHODS OF INTEGRATING.

Article.	Page.
49. Integral regarded as the inverse of a <i>differential</i> . . . . .	32
50. If $fx$ is any function whatever of $x$ , $fx.dx$ has an integral, and but one, except for the presence of an arbitrary constant . .	32
51. A <i>definite integral</i> contains no arbitrary constant, and is a function of the values between which the sum is taken. <i>Examples</i> . . . . .	33
52. Definite integral of a <i>discontinuous</i> function . . . . .	33
53. Formulas for direct integration . . . . .	34
54. Integration by <i>substitution</i> . <i>Examples</i> . . . . .	36
55. Integration by <i>parts</i> . <i>Examples</i> . <i>Miscellaneous examples in integration</i> . . . . .	37

CHAPTER IV.

RATIONAL FRACTIONS.

56. Integration of a <i>rational algebraic polynomial</i> . <i>Rational fractions, proper and improper</i> . . . . .	40
57. Every proper rational fraction can be reduced to a sum of simpler fractions with constant numerators . . . . .	40
58. Determination of the numerators of the partial fractions by indirect methods. <i>Examples</i> . . . . .	42
59. Direct determination of the numerators of the partial fractions	43
60. Illustrative examples . . . . .	45
61. Illustrative example . . . . .	46
62. Integration of the partial fractions . . . . .	48
63. Treatment of imaginary values which may occur in the partial fractions. <i>Examples</i> . . . . .	49

CHAPTER V.

REDUCTION FORMULAS.

64. Formulas for raising or lowering the exponents in the form $x^{m-1}(a+bx^n)^p dx$ . . . . .	52
65. Consideration of special cases. <i>Examples</i> . . . . .	54

## CHAPTER VI.

## IRRATIONAL FORMS.

Article.	Page
66. Integration of the form $f(x, \sqrt[n]{a+bx})dx$ . Examples . . . . .	56
67. Integration of the form $f(x, \sqrt[n]{c+\sqrt[m]{a+bx}})dx$ . Examples . .	57
68. Integration of the form $f(x, \sqrt{a+bx+cx^2})dx$ . . . . .	57
69. Illustrative example. Examples . . . . .	59
70. Integration of the form $f\left(x, \sqrt[n]{\frac{ax+b}{lx+m}}\right)dx$ . Example . . . . .	61
71. Application of the <i>Reduction Formulas</i> of Chapter V. to irrational forms. Examples . . . . .	61
72. A function rendered irrational through the presence under the radical sign of a polynomial of higher degree than the second cannot ordinarily be integrated. Elliptic Integrals .	62

## CHAPTER VII.

## TRANSCENDENTAL FUNCTIONS.

73. Use of the method of <i>Integration by Parts</i> . Examples . . . . .	63
74. Reduction Formulas for $\sin^n x$ and $\cos^n x$ . Examples . . . . .	64
75. Integration of $(\sin^{-1}x)^n dx$ . Examples . . . . .	65
76. Use of the method of <i>Integration by Substitution</i> . . . . .	66
77. Substitution of $z = \tan \frac{x}{2}$ in integrating trigonometric forms .	67
78. Integration of $\sin^n x \cos^n x dx$ . Examples . . . . .	68
79. Reduction formulas for $\int \tan^n x dx$ and $\int \frac{dx}{\tan^n x}$ . Examples . .	69

## CHAPTER VIII.

## DEFINITE INTEGRALS.

80. Definition of a <i>definite integral</i> as the limit of a sum of infinitesimals . . . . .	71
81. Computation of a definite integral as the limit of a sum. Illustrative examples. Examples . . . . .	72
82. Usual method of obtaining the value of a definite integral. Caution concerning multiple-valued functions. Examples .	76

TABLE OF CONTENTS.

ix

Article.	Page.
83. Consideration of the nature of the value of $\int_a^b fx \cdot dx$ when $fx$ becomes infinite for $x=a$ , or $x=b$ , or for some value of $x$ between $a$ and $b$ . Illustrative examples . . . . .	78
84. Test that must be satisfied in order that $\int_a^b fx \cdot dx$ may be finite and determinate if $fx$ is infinite for some value of $x$ between $a$ and $b$ . Illustrative examples. Examples . . . . .	80
85. Meaning of $\int_a^\infty fx \cdot dx$ . Condition that $\int_a^\infty fx \cdot dx$ shall be finite and determinate . . . . .	84
86. <i>Maximum-Minimum Theorem</i> . Proof that certain important definite integrals of the form $\int_a^\infty fx \cdot dx$ are finite and determinate. Examples . . . . .	85
87. Application of <i>reduction formulas</i> to definite integrals. Examples . . . . .	88
88. Application of the method of <i>integration by substitution</i> to definite integrals. Illustrative examples. Example . . . . .	90
89. Differentiation of a definite integral. Examples . . . . .	93
90. Many ingenious methods of finding the values of definite integrals are valid only in case the integral is finite and determinate . . . . .	96
91. <i>Integration by development in series</i> . Examples . . . . .	96
92. Values of $\int_0^{\frac{\pi}{2}} \log \sin x \cdot dx$ , $\int_0^\infty e^{-x^2} dx$ , $\int_0^\infty \frac{\sin mx}{x} \cdot dx$ obtained by ingenious devices. Examples . . . . .	98
93. Differentiation or integration with respect to a quantity which is independent of $x$ . Examples . . . . .	101
94. Additional illustrative examples. Examples . . . . .	102
95. Introduction of imaginary constants. . . . .	104
96. The <i>Gamma Function</i> . . . . .	105
97. Table giving $\log \Gamma(n)$ from $n=1$ to $n=2$ . Definite integrals expressed as Gamma Functions . . . . .	107
98. The <i>Beta Function</i> . Formula connecting the Beta Function with the Gamma Function. Value of $\Gamma(\frac{1}{2})$ . . . . .	109
99. More definite integrals expressed as Gamma Functions. Examples . . . . .	110

CHAPTER IX.

LENGTHS OF CURVES.

Article.	Page.
100. Formulas for $\sin \tau$ and $\cos \tau$ in terms of the length of the arc	113
101. The equation of the <i>Catenary</i> obtained. Example . . . . .	113
102. The equation of the <i>Tractrix</i> . Examples . . . . .	115
103. Length of an arc in rectangular coördinates . . . . .	117
104. Length of the arc of the <i>Cycloid</i> . Example . . . . .	118
105. Another method of rectifying the <i>Cycloid</i> . . . . .	119
106. Rectification of the <i>Epicycloid</i> . Examples . . . . .	119
107. Arc of the <i>Ellipse</i> . Auxillary angle. Example . . . . .	120
108. Length of an arc. Polar coördinates . . . . .	121
109. Equation of the <i>Logarithmic Spiral</i> . . . . .	121
110. Rectification of the Logarithmic Spiral. Examples . . . . .	122
111. Rectification of the <i>Cardioid</i> . . . . .	122
112. <i>Involutes</i> . Illustrative example. Example . . . . .	123
113. The <i>involute</i> of the <i>Cycloid</i> . Example . . . . .	125
114. <i>Intrinsic equation</i> of a curve. Example . . . . .	126
115. Intrinsic equation of the <i>Epicycloid</i> . Example . . . . .	127
116. Intrinsic equation of the <i>Logarithmic Spiral</i> . . . . .	128
117. Method of obtaining the <i>intrinsic equation</i> from the equation in rectangular coördinates. Examples . . . . .	128
118. Intrinsic equation of an <i>evolute</i> . . . . .	130
119. Illustrative examples. Examples . . . . .	130
120. The evolute of an <i>Epicycloid</i> . Example . . . . .	131
121. The intrinsic equation of an involute. Illustrative examples	132
122. Limiting form approached by an involute of an involute . .	133
123. Method of obtaining the equation in rectangular coördinates from the intrinsic equation. Illustrative example . . . . .	134
124. Rectification of <i>Curves in Space</i> . Examples . . . . .	135

CHAPTER X.

AREAS.

125. <i>Areas</i> expressed as definite integrals, rectangular coördinates. Examples . . . . .	137
126. <i>Areas</i> expressed as definite integrals, polar coördinates . .	139
127. Area between the catenary and the axis . . . . .	139
128. Area between the tractrix and the axis. Example . . . . .	139



TABLE OF CONTENTS.

Article.	Page.
129. Area between a curve and its asymptote. Examples . . .	140
130. Area of circle obtained by the aid of an auxiliary angle. Ex- amples . . . . .	141
131. Area between two curves (rect. coör.). Examples . . . .	142
132. Areas in Polar Coördinates. Examples . . . . .	143
133. Problems in areas can often be simplified by transformation of coördinates. Examples . . . . .	146
134. Area between a curve and its evolute. Examples . . . .	146
135. Holditch's Theorem. Examples . . . . .	147
136. Areas by a <i>double integration</i> (rect. coör.) . . . . .	149
137. Illustrative examples. Examples . . . . .	150
138. Areas by a <i>double integration</i> (polar coör.). Example . . .	151

CHAPTER XI.

AREAS OF SURFACES.

139. Area of a <i>surface of revolution</i> (rect. coör.). Example . . .	153
140. Illustrative examples. Examples . . . . .	154
141. Area of a surface of revolution by transformation of coördi- nates. Example . . . . .	155
142. Area of a <i>surface of revolution</i> (polar coör.). Examples . .	157
143. Area of a <i>cylindrical surface</i> . Examples . . . . .	157
144. Area of <i>any surface</i> by a double integration . . . . .	160
145. Illustrative example. Examples . . . . .	163
146. Illustrative example requiring transformation to polar coördi- nates. Examples . . . . .	165

CHAPTER XII.

VOLUMES.

147. Volume by a single integration. Example . . . . .	168
148. Volume of a <i>conoid</i> . Examples . . . . .	169
149. Volume of an <i>ellipsoid</i> . Examples . . . . .	170
150. Volume of a <i>solid of revolution</i> . Single integration. Exam- ples . . . . .	171
151. Volume of a <i>solid of revolution</i> . Double integration. Exam- ples . . . . .	172
152. Volume of a <i>solid of revolution</i> . Polar formula. Example .	174
153. Volume of <i>any solid</i> . Triple integration. Rectangular coör- dinates. Examples . . . . .	175
154. Volume of <i>any solid</i> . Triple integration. Polar coördinates. Examples . . . . .	179



## CHAPTER XIII.

## CENTRES OF GRAVITY.

Article.	Page.
155. Centre of Gravity defined . . . . .	180
156. General formulas for the coördinates of the Centre of Gravity of any mass. Example . . . . .	180
157. Centre of Gravity of a homogeneous body . . . . .	182
158. Centre of Gravity of a <i>plane area</i> . Examples . . . . .	182
159. Centre of Gravity of a <i>homogeneous solid</i> of revolution. Ex- amples . . . . .	185
160. Centre of Gravity of an <i>arc</i> ; of a <i>surface of revolution</i> . Ex- amples . . . . .	187
161. <i>Properties of Guldin</i> . Examples . . . . .	188

## CHAPTER XIV.

## LINE, SURFACE, AND SPACE INTEGRALS.

162. <i>Point function</i> . <i>Continuity</i> of a point function . . . . .	190
163. <i>Line-integral</i> , <i>surface-integral</i> , and <i>space-integral</i> of a point function . . . . .	190
164. Value of a line, surface, or space integral independent of the position in each element of the point at which the value of the function is taken . . . . .	191
165. Value of a line, surface, or space integral independent of the manner in which the line, surface, or space is broken up into infinitesimal elements . . . . .	191
166. Geometrical representation of a line-integral along a plane curve, and a surface-integral over a plane surface . . . . .	193
167. <i>Moments of inertia</i> . Examples . . . . .	193
168. Relation between a surface-integral over a plane surface, and a line-integral along the curve bounding the surface. Ex- ample . . . . .	196
169. Illustrative example. Examples . . . . .	198
170. Another form of the relation established in Art. 168. Example	199
171. Relation between a space-integral taken throughout a given space and a surface-integral over the surface bounding the space. Example . . . . .	199
172. Illustrative example. Example . . . . .	201



TABLE OF CONTENTS.

CHAPTER XV.

MEAN VALUE AND PROBABILITY.

Article.	Page.
173. References . . . . .	202
174. <i>Mean value of a continuously varying quantity.</i> The <i>mean distance</i> of all the points of the circumferences of a circle from a fixed point on the circumference. The <i>mean distance</i> of points on the surface of a circle from a fixed point on the circumference. The <i>mean distance</i> of points on the surface of a square from a corner of the square. The <i>mean distance</i> between two points within a given circle . . . . .	202
175. Problems in the application of the Integral Calculus to <i>probabilities</i> . Random straight lines. Examples . . . . .	204

CHAPTER XVI.

ELLIPTIC INTEGRALS.

176. Motion of a <i>simple pendulum</i> . <i>Vibration</i> . <i>Complete revolution</i>	211
177. The <i>length</i> of an arc of an <i>Ellipse</i> . . . . .	213
178. Algebraic forms of the <i>Elliptic Integrals</i> of the first, second, and third class. <i>Modulus</i> . <i>Parameter</i> . . . . .	213
179. Trigonometric forms of the <i>Elliptic Integrals</i> . <i>Amplitude</i> . <i>Delta</i> . <i>Complementary Modulus</i> . . . . .	214
180. <i>Landen's Transformation</i> . Reduction formula by which we can increase the modulus and diminish the amplitude of an <i>Elliptic Integral</i> of the first class. A method of computing $F(k, \phi)$ . . . . .	215
181. Reduction formula for diminishing the modulus and increasing the amplitude of an <i>Elliptic Integral</i> of the first class. A second method of computing $F(k, \phi)$ . . . . .	218
182. Actual computation of $F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$ and $F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right)$ . . . . .	219
183. <i>Landen's Transformation</i> . Reduction formula by which we can increase the modulus and diminish the amplitude of an <i>Elliptic Integral</i> of the second class. A method of computing $E(k, \phi)$ . . . . .	222
184. A reduction formula for diminishing the modulus and increasing the amplitude of an <i>Elliptic Integral</i> of the second class. A second method of computing $E(k, \phi)$ . . . . .	225

Article.	Page
185. Actual computation of $E\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$ and $E\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right)$ . . . . .	227
186. An Elliptic Integral of the first or second class, whose amplitude is greater than $\frac{\pi}{2}$ , can be made to depend upon one whose amplitude is less than $\frac{\pi}{2}$ , and upon the corresponding complete Elliptic Integral . . . . .	231
187. Three-place table of Elliptic Integrals of the first class and the second class . . . . .	233
188. <i>Addition Formulas</i> . Functions defined by the aid of definite integrals. $\log x$ , $\sin^{-1}x$ , $\tan^{-1}x$ , $F(k, x)$ , $E(k, x)$ . Addition formula for $\log x$ . . . . .	235
189. Addition formulas for $\sin^{-1}x$ and $\tan^{-1}x$ . . . . .	237
190. Addition formula for $F(k, x)$ . . . . .	239
191. Analogy between $\log^{-1}u$ , $\sin u$ , $\tan u$ , and $F^{-1}(k, u)$ . . . . .	241
192. The <i>Elliptic Functions</i> , $\operatorname{sn} u$ , $\operatorname{cn} u$ , and $\operatorname{dn} u$ . Their analogy with Trigonometric Functions. Formulas connecting the Elliptic Functions of a single quantity . . . . .	242
193. Formulas for Elliptic Functions of $(u+v)$ and $(u-v)$ . . . . .	244
194. Formulas for Elliptic Functions of $2u$ . . . . .	246
195. Formulas for Elliptic Functions of $\frac{u}{2}$ . . . . .	246
196. <i>Periodicity</i> of the Elliptic Functions. Real period, $4K$ . . . . .	247
197. Elliptic Functions of a pure imaginary. <i>Jacobi's Transformation</i> . Elliptic Functions have an imaginary period, $4K'\sqrt{-1}$ . Table of values of Elliptic Functions having the modulus $\frac{\sqrt{2}}{2}$ . . . . .	249
198. The Elliptic Integral of the second class expressed in terms of Elliptic Functions. <i>Addition formula</i> for Elliptic Integrals of the second class . . . . .	252
199. <i>Application to the rectification of the Lemniscate</i> . Examples. Bisection of the arc of a quadrant of the Lemniscate . . . . .	254
200. Rectification of the Ellipse. Examples . . . . .	257
201. Use of the Addition Formula in dealing with Elliptic arcs. <i>Fagnani's Point</i> . Examples . . . . .	258
202. Rectification of the <i>Hyperbola</i> . Examples . . . . .	260
203. The <i>simple pendulum</i> . Examples . . . . .	263

## CHAPTER XVII.

## INTRODUCTION TO THE THEORY OF FUNCTIONS.

Article.	Page.
204. <i>Single-valued</i> functions. <i>Multiple-valued</i> functions . . . . .	267
205. Importance of the graphical representation of imaginaries. <i>Complex quantity</i> . . . . .	267
206. When a complex variable is said to vary continuously . . . . .	268
207. A <i>continuous</i> function of a complex variable. <i>Critical values</i>	268
208. Criterion that a function shall have a determinate derivative. <i>Monogenic</i> functions . . . . .	269
209. Any function involving $z$ as a whole is a monogenic function	272
210. <i>Conjugate</i> functions. Their use as solutions of Laplace's Equation. Example . . . . .	272
211. <i>Preservation of angles</i> . . . . .	274
212. If two paths traced by the point representing the variable have a common beginning and a common end, and do not enclose a <i>critical point</i> , the corresponding paths traced by the point representing the function and having a common beginning will have a common end . . . . .	276
213. Examples where the paths traced by the point representing the variable enclose a critical point . . . . .	278
214. Critical points at which the derivative of the function is zero or infinite are to be avoided. <i>Branch points. Holomorphic</i> <i>functions</i> . . . . .	279
215. Definite integral of a function of a complex variable defined. Such an integral is generally indeterminate, and depends upon the path by which the point representing the variable passes from the lower limit to the upper limit of the integral	281
216. If the function is <i>holomorphic</i> , the definite integral is in gen- eral determinate . . . . .	283
217. The integral around a closed contour, embracing a point at which the function is infinite . . . . .	284
218. Illustrative examples . . . . .	285
219. Convergency of the series obtained by integrating the terms of a convergent series where the separate terms are holo- morphic functions . . . . .	287
220. Proof of Taylor's and Maclaurin's Theorems for functions of complex variables. <i>Circle of convergence</i> . . . . .	288
221. Investigation of the convergency of various series which are obtained by Taylor's and Maclaurin's Theorems. Examples	291

## CHAPTER XVIII.

## KEY TO THE SOLUTION OF DIFFERENTIAL EQUATIONS.

Article.	Page.
222. Description of Key . . . . .	296
223. Definition of the terms <i>differential equation, order, degree, linear, general solution or complete primitive, singular solution, exact differential equation</i> . . . . .	296
224. Examples illustrating the use of the Key . . . . .	298
225. Simplification of differential equations by change of variable.	308
KEY . . . . .	310
EXAMPLES UNDER KEY . . . . .	333



# INTEGRAL CALCULUS.

---

## CHAPTER I.

### SYMBOLS OF OPERATION.

1. It is often convenient to regard a functional symbol as indicating an operation to be performed upon the expression which is written after the symbol. From this point of view the symbol is called a *symbol of operation*, and the expression written after the symbol is called the *subject* of the operation.

Thus the symbol  $D_x$  in  $D_x(x^2y)$  indicates that the operation of differentiating with respect to  $x$  is to be performed upon the subject  $(x^2y)$ .

2. If the *result* of one operation is taken as the *subject* of a second, there is formed what is called a *compound function*.

Thus  $\log \sin x$  is a *compound function*, and we may speak of the taking of the  $\log \sin$  as a *compound operation*.

3. When two operations are so related that the compound operation, in which the result of performing the first on any subject is taken as the subject of the second, leads to the same result as the compound operation, in which the result of performing the second on the same subject is taken as the subject of the first, the two operations are *commutative* or *relatively free*.

Or to formulate ; if

$$fFu = Ffu,$$

the operations indicated by  $f$  and  $F$  are *commutative*.

4

For example; the operations of partial differentiation with respect to two independent variables  $x$  and  $y$  are commutative, for we know that

$$D_x D_y u = D_y D_x u. \quad (\text{I. Art. 197}).$$

The operations of taking the sine and of taking the logarithm are not commutative, for  $\log \sin u$  is not equal to  $\sin \log u$ .

4. If  $f(u \pm v) = fu \pm fv$

where  $u$  and  $v$  are any subjects, the operation  $f$  is *distributive* or *linear*.

The operation indicated by  $d$  and the operation indicated by  $D_x$  are distributive, for we know that

$$d(u \pm v) = du \pm dv,$$

and that

$$D_x(u \pm v) = D_x u \pm D_x v.$$

The operation  $\sin$  is not distributive, for  $\sin(u + v)$  is not equal to  $\sin u + \sin v$ .

5. *The compounds of distributive operations are distributive.*

Let  $f$  and  $F$  indicate distributive operations, then  $fF$  will be distributive; for

$$F(u \pm v) = Fu \pm Fv,$$

therefore  $fF(u \pm v) = f(Fu \pm Fv) = fFu \pm fFv$ .

6. The *repetition* of any operation is indicated by writing an *exponent*, equal to the number of times the operation is performed, after the symbol of the operation.

Thus  $\log^3 x$  means  $\log \log \log x$ ;  $d^3 u$  means  $dddu$ .

In the single case of the trigonometric functions a different use of the exponent is sanctioned by custom, and  $\sin^2 u$  means  $(\sin u)^2$  and not  $\sin \sin u$ .

7. If  $m$  and  $n$  are *whole numbers* it is easily proved that

$$f^m f^n u = f^{m+n} u,$$

*This formula is assumed for all values of m and n, and negative and fractional exponents are interpreted by its aid. It is called the law of indices.*

8. To find what interpretation must be given to a zero exponent, let

$$m = 0 \quad \text{in the formula of Art. 7.}$$

$$f^0 f^n u = f^{0+n} u = f^n u,$$

or, denoting  $f^n u$  by  $v$ ,  $f^0 v = v$ .

That is; a symbol of operation with the exponent zero has no effect on the subject, and may be regarded as multiplying it by unity.

9. To interpret a negative exponent, let

$$m = -n \quad \text{in the formula of Art. 7.}$$

$$f^{-n} f^n u = f^{-n+n} u = f^0 u = u.$$

If we call  $f^n u = v$ , then  $f^{-n} v = u$ .

If  $n = 1$

we get  $f^{-1} f u = u$ ,

and the exponent  $-1$  indicates what we have called the anti-function of  $f u$ . (I. Art. 72.)

The exponent  $-1$  is used in this sense even with trigonometric functions.

10. When two operations are commutative and distributive, the symbols which represent them may be combined precisely as if they were algebraic quantities.

For they obey the laws,

$$a(m + n) = am + an,$$

$$am = ma,$$

on which all the operations of arithmetic and algebra are founded.



For example ; if the operation  $(D_x + D_y)$  is to be performed  $n$  times in succession on a subject  $u$ , we can expand  $(D_x + D_y)^n$  precisely as if it were a binominal, and then perform on  $u$  the operations indicated by the expanded expression.

$$\begin{aligned}(D_x + D_y)^3 u &= (D_x^3 + 3 D_x^2 D_y + 3 D_x D_y^2 + D_y^3) u \\ &= D_x^3 u + 3 D_x^2 D_y u + 3 D_x D_y^2 u + D_y^3 u.\end{aligned}$$

## CHAPTER II.

## IMAGINARIES.

11. An *imaginary* is usually defined in algebra as *the indicated even root of a negative quantity*, and although it is clear that there can be no *quantity* that raised to an even power will be negative, the assumption is made that an imaginary can be treated like any algebraic quantity.

Imaginaries are first forced upon our notice in connection with the subject of quadratic equations. Considering the typical quadratic

$$x^2 + ax + b = 0,$$

we find that it has two roots, and that these roots possess certain important properties. For example; their sum is  $-a$  and their product is  $b$ . We are led to the conclusion that every quadratic has two roots whose sum and whose product are simply related to the coefficients of the equation.

On trial, however, we find that there are quadratics having but one root, and quadratics having no root.

For example; if we solve the equation

$$x^2 - 2x + 1 = 0,$$

we find that the only value of  $x$  which will satisfy it is *unity*; and if we attempt to solve

$$x^2 - 2x + 2 = 0,$$

we find that there is no value of  $x$  which will satisfy the equation.

As these results are apparently inconsistent with the conclusion to which we were led on solving the general equation, we naturally endeavor to reconcile them with it.

The difficulty in the case of the equation which has but one

root is easily overcome by regarding it as having two equal roots. Thus we can say that each of the two roots of the equation

$$x^2 - 2x + 1 = 0$$

is equal to 1; and there is a decided advantage in looking at the question from this point of view, for the roots of this equation will possess the same properties as those of a quadratic having unequal roots. The sum of the roots 1 and 1 is minus the coefficient of  $x$  in the equation, and their product is the constant term.

To overcome the difficulty presented by the equation which has no root we are driven to the conception of *imaginaries*.

12. *An imaginary is not a quantity, and the treatment of imaginaries is purely arbitrary and conventional.* We begin by laying down a few arbitrary rules for our imaginary expressions to obey, which must not involve any contradiction; and we must perform all our operations upon imaginaries, and must interpret all our results by the aid of these rules.

Since imaginaries occur as roots of equations, they bear a close analogy with ordinary algebraic quantities, and they have to be subjected to the same operations as ordinary quantities; therefore our rules ought to be so chosen that the results may be comparable with the results obtained when we are dealing with real quantities.

13. By adopting the convention that

$$\sqrt{-a^2} = a\sqrt{-1},$$

where  $a$  is supposed to be *real*, we can reduce all our imaginary algebraic expressions to forms where  $\sqrt{-1}$  is the only peculiar symbol. This symbol  $\sqrt{-1}$  we shall define and use as *the symbol of some operation, at present unknown, the repetition of which has the effect of changing the sign of the subject of the operation.* Thus in  $a\sqrt{-1}$  the symbol  $\sqrt{-1}$  indicates that an operation is performed upon  $a$  which, if repeated, will change the sign of  $a$ . That is,

$$a(\sqrt{-1})^2 = -a.$$

From this point of view it would be more natural to write the symbol before instead of after the subject on which it operates,  $(\sqrt{-1})a$  instead of  $a\sqrt{-1}$ , and this is sometimes done; but as the usage of mathematicians is overwhelmingly in favor of the second form, we shall employ it, merely as a matter of convenience, and remembering that  $a$  is the *subject* and the  $\sqrt{-1}$  the *symbol of operation*.

14. The rules in accordance with which we shall use our new symbol are, first,

$$a\sqrt{-1} + b\sqrt{-1} = (a + b)\sqrt{-1}. \quad [1]$$

In other words, the operation indicated by  $\sqrt{-1}$  is to be *distributive* (Art. 4); and second,

$$a\sqrt{-1} = (\sqrt{-1})a, \quad [2]$$

or our symbol is to be *commutative* with the symbols of quantity (Art. 3).

These two conventions will enable us to use our symbol in algebraic operations precisely as if it were a quantity (Art. 10).

When no coefficient is written before  $\sqrt{-1}$  the coefficient 1 will be understood, or unity will be regarded as the subject of the operation.

15. Let us see what interpretation we can get for powers of  $\sqrt{-1}$ ; that is, for repetitions of the operation indicated by the symbol.

$$(\sqrt{-1})^0 = 1 \quad (\text{Art. 8}),$$

$$(\sqrt{-1})^1 = \sqrt{-1},$$

$$(\sqrt{-1})^2 = -1, \quad \text{by definition (Art. 13),}$$

$$(\sqrt{-1})^3 = (\sqrt{-1})^2\sqrt{-1} = -\sqrt{-1}, \quad \text{by definition,}$$

$$(\sqrt{-1})^4 = -(\sqrt{-1})^2 = 1,$$

$$(\sqrt{-1})^5 = 1\sqrt{-1} = \sqrt{-1},$$

$$(\sqrt{-1})^6 = (\sqrt{-1})^2 = -1,$$

and so on, the values  $\sqrt{-1}$ ,  $-1$ ,  $-\sqrt{-1}$ ,  $1$ , occurring in cycles of four.

16. The definition we have given for the square root of a negative quantity, and the rules we have adopted concerning its use, enable us to remove entirely the difficulty felt in dealing with a quadratic which does not have real roots. Take the equation

$$x^2 - 2x + 5 = 0. \quad (1)$$

Solving by the usual method, we get

$$x = 1 \pm \sqrt{-4};$$

$$\sqrt{-4} = 2\sqrt{-1}, \text{ by Art. 13 [1];}$$

hence  $x = 1 + 2\sqrt{-1}$  or  $1 - 2\sqrt{-1}$ .

On substituting these results in turn in the equation (1), performing the operations by the aid of our conventions (Art. 14 [1] and [2]), and interpreting  $(\sqrt{-1})^2$  by Art. 15, we find that they both satisfy the equation, and that they can therefore be regarded as entirely analogous to real roots. We find, too, that their sum is 2 and that their product is 5, and consequently that they bear the same relations to the coefficients of the equation as real roots.

17. An imaginary root of a quadratic can always be reduced to the form  $a + b\sqrt{-1}$  where  $a$  and  $b$  are real, and this is taken as the general type of an imaginary; and part of our work will be to show that when we subject imaginaries to the ordinary functional operations, all our results are reducible to this typical form.

If two imaginaries  $a + b\sqrt{-1}$  and  $c + d\sqrt{-1}$  are equal,  $a$  must be equal to  $c$ , and  $b$  must be equal to  $d$ .

For we have  $a + b\sqrt{-1} = c + d\sqrt{-1}$ .

Therefore  $a - c = (d - b)\sqrt{-1}$ ,

or a real is equal to an imaginary, unless  $a - c = 0 = d - b$ .

Since obviously a real and an imaginary cannot be equal, it follows that  $a = c$  and  $b = d$ .


18. We have defined  $\sqrt{-1}$  as the symbol of an operation whose repetition changes the sign of the subject.

Several different interpretations of this operation have been suggested, and the following one, in which every imaginary is graphically represented by the position of a point in a plane, is commonly adopted, and is found exceedingly useful in suggesting and interpreting relations between different imaginaries and between imaginaries and reals.

In the *Calculus of Imaginaries*,  $a + b\sqrt{-1}$  is taken as the general symbol of quantity. If  $b$  is equal to zero,  $a + b\sqrt{-1}$  reduces to  $a$ , and is *real*; if  $a$  is equal to zero,  $a + b\sqrt{-1}$  reduces to  $b\sqrt{-1}$ , and is called a *pure imaginary*.

$a + b\sqrt{-1}$  is represented by the position of a point referred to a pair of rectangular axes, as in analytic geometry,  $a$  being taken as the abscissa of the point and  $b$  as its ordinate. Thus in the figure the position of the point  $P$  represents the imaginary  $a + b\sqrt{-1}$ .

If  $b = 0$ , and our quantity is real,  $P$  will lie on the axis of  $X$ , which on that account is called the *axis of reals*; if  $a = 0$ , and we have a *pure imaginary*,  $P$  will lie on the axis of  $Y$ , which is called the *axis of pure imaginaries*.



It follows from Art. 17 that if two imaginaries are equal, the points representing them will coincide.

Since  $a$  and  $a\sqrt{-1}$  are represented by points equally distant from the origin, and lying on the *axis of reals* and the *axis of pure imaginaries* respectively, we may regard the operation indicated by  $\sqrt{-1}$  as causing the point representing the subject of the operation to rotate about the origin through an angle of  $90^\circ$ . A repetition of the operation ought to cause the point to rotate  $90^\circ$  further, and it does; for

$$a(\sqrt{-1})^2 = -a,$$

and is represented by a point at the same distance from the

origin as  $a$ , and lying on the opposite side of the origin ; again repeat the operation,

$$a(\sqrt{-1})^3 = -a\sqrt{-1},$$

and the point has rotated  $90^\circ$  further ; repeat again,

$$a(\sqrt{-1})^4 = a,$$

and the point has rotated through  $360^\circ$ . We see, then, that if the subject is a *real* or a *pure imaginary* the effect of performing on it the operation indicated by  $\sqrt{-1}$  is to rotate it about the origin through the angle  $90^\circ$ . We shall see later that even when the subject is neither a real nor a pure imaginary, the effect of operating on it with  $\sqrt{-1}$  is still to produce the rotation just described.

19. The *sum*, the *product*, and the *quotient* of any two imaginaries,  $a + b\sqrt{-1}$  and  $c + d\sqrt{-1}$ , are imaginaries of the typical form.

$$a + b\sqrt{-1} + c + d\sqrt{-1} = a + c + (b + d)\sqrt{-1}. \quad [1]$$

$$(a + b\sqrt{-1})(c + d\sqrt{-1}) = ac - bd + (bc + ad)\sqrt{-1}. \quad [2]$$

$$\begin{aligned} \frac{a + b\sqrt{-1}}{c + d\sqrt{-1}} &= \frac{(a + b\sqrt{-1})(c - d\sqrt{-1})}{(c + d\sqrt{-1})(c - d\sqrt{-1})} = \frac{ac + bd + (bc - ad)\sqrt{-1}}{c^2 + d^2} \\ &= \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}\sqrt{-1}. \end{aligned} \quad [3]$$


All these results are of the form  $A + B\sqrt{-1}$ .

20. The graphical representation we have suggested for imaginaries suggests a second typical form for an imaginary. Given the imaginary  $x + y\sqrt{-1}$ , let the *polar coördinates* of the point  $P$  which represents  $x + y\sqrt{-1}$  be  $r$  and  $\phi$ .

$r$  is called the *modulus* and  $\phi$  the *argument* of the imaginary.



The figure enables us to establish very simple relations between  $x$ ,  $y$ ,  $r$ , and  $\phi$ .



$$\left. \begin{aligned} x &= r \cos \phi, \\ y &= r \sin \phi; \end{aligned} \right\} \quad [1]$$

$$\left. \begin{aligned} r &= \sqrt{x^2 + y^2}, \\ \phi &= \tan^{-1} \frac{y}{x}. \end{aligned} \right\} \quad [2]$$

$$\begin{aligned} x + y\sqrt{-1} &= r \cos \phi + (\sqrt{-1})r \sin \phi \\ &= r(\cos \phi + \sqrt{-1} \cdot \sin \phi), \end{aligned} \quad [3]$$

where the imaginary is expressed in terms of its modulus and argument.

The value of  $r$  given by our formulas [2] is ambiguous in sign; and  $\phi$  may have any one of an infinite number of values differing by multiples of  $\pi$ . In practice we always take the positive value of  $r$ , and a value of  $\phi$  which will bring the point in question into the right quadrant. In the case of any given imaginary then,  $r$  can have but one value, while  $\phi$  may have any one of an infinite number of values differing by multiples of  $2\pi$ .

The modulus  $r$  is sometimes called the *absolute value* of the imaginary.

#### EXAMPLES.

(1) Find the modulus and argument of 1; of  $\sqrt{-1}$ ; of  $-4$ ; of  $-2\sqrt{-1}$ ; of  $3+3\sqrt{-1}$ ; of  $2+4\sqrt{-1}$ ; and express each of these quantities in the form  $r(\cos \phi + \sqrt{-1} \cdot \sin \phi)$ .

(2) Show that every positive real has the argument zero; every negative real the argument  $\pi$ ; every positive pure imaginary the argument  $\frac{\pi}{2}$ ; and every negative pure imaginary the argument  $\frac{3\pi}{2}$ .

21. If we add two imaginaries, the *modulus of the sum* is never greater than the *sum of the moduli* of the given imaginaries.



The sum of  $a + b\sqrt{-1}$  and  $c + d\sqrt{-1}$  is  $a + c + (b + d)\sqrt{-1}$ . The modulus of this sum is  $\sqrt{(a + c)^2 + (b + d)^2}$ ; the sum of the moduli of  $a + b\sqrt{-1}$  and  $c + d\sqrt{-1}$  is  $\sqrt{a^2 + b^2} + \sqrt{c^2 + d^2}$ . We wish to show that

$$\sqrt{(a + c)^2 + (b + d)^2} < \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2};$$

the sign  $<$  meaning "equal to or less than."

$$\text{Now } \sqrt{(a + c)^2 + (b + d)^2} < \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2},$$

$$\text{if } (a + c)^2 + (b + d)^2 < a^2 + b^2 + 2\sqrt{(a^2 + b^2)(c^2 + d^2)} + c^2 + d^2,$$

$$\text{that is, if } ac + bd < \sqrt{a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2};$$

or, squaring, if

$$a^2c^2 + 2abcd + b^2d^2 < a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2;$$

or, if

$$0 < (ad - bc)^2.$$

This last result is necessarily true, as no real can have a square less than zero; hence our proposition is established.

22. *The modulus of the product of two imaginaries is the product of the moduli of the given imaginaries, and the argument of the product is the sum of the arguments of the imaginaries.*

Let us multiply

$$r_1(\cos \phi_1 + \sqrt{-1} \cdot \sin \phi_1) \quad \text{by} \quad r_2(\cos \phi_2 + \sqrt{-1} \cdot \sin \phi_2);$$

we get

$$r_1 r_2 [\cos \phi_1 \cos \phi_2 - \sin \phi_1 \sin \phi_2 + \sqrt{-1}(\sin \phi_1 \cos \phi_2 + \cos \phi_1 \sin \phi_2)],$$

$$\cos \phi_1 \cos \phi_2 - \sin \phi_1 \sin \phi_2 = \cos(\phi_1 + \phi_2),$$

$$\sin \phi_1 \cos \phi_2 + \cos \phi_1 \sin \phi_2 = \sin(\phi_1 + \phi_2)$$

by Trigonometry; hence

$$\begin{aligned} & r_1(\cos \phi_1 + \sqrt{-1} \cdot \sin \phi_1) r_2(\cos \phi_2 + \sqrt{-1} \cdot \sin \phi_2) \\ &= r_1 r_2 [\cos(\phi_1 + \phi_2) + \sqrt{-1} \cdot \sin(\phi_1 + \phi_2)], \end{aligned}$$

and our result is in the typical form,  $r_1 r_2$  being the modulus and  $\phi_1 + \phi_2$  the argument of the product.

If each factor has the modulus unity, this theorem enables us to construct very easily the product of the imaginaries; it also enables us to show that the interpretation of the operation  $\sqrt{-1}$ , suggested in Art. 18, is perfectly general.

Let us operate on any imaginary subject,

$$r(\cos \phi + \sqrt{-1} \cdot \sin \phi), \quad \text{with } \sqrt{-1},$$

that is, with  $1\left(\cos \frac{\pi}{2} + \sqrt{-1} \cdot \sin \frac{\pi}{2}\right).$

The modulus  $r$  will be unchanged, the argument  $\phi$  will be increased by  $\frac{\pi}{2}$ , and the effect will be to cause the point representing the given imaginary to rotate about the origin through an angle of  $90^\circ$ .

23. Since division is the inverse of multiplication,

$$r_1(\cos \phi_1 + \sqrt{-1} \cdot \sin \phi_1) + r_2(\cos \phi_2 + \sqrt{-1} \cdot \sin \phi_2)$$

will be equal to

$$\frac{r_1}{r_2}[\cos(\phi_1 - \phi_2) + \sqrt{-1} \cdot \sin(\phi_1 - \phi_2)],$$

since if we multiply this by  $r_2(\cos \phi_2 + \sqrt{-1} \cdot \sin \phi_2)$ , according to the method established in Art. 22, we must get

$$r_1(\cos \phi_1 + \sqrt{-1} \cdot \sin \phi_1).$$

*To divide one imaginary by another, we have then to take the quotient obtained by dividing the modulus of the first by the modulus of the second as our required modulus, and the argument of the first minus the argument of the second as our new argument.*

24. If we are dealing with the product of  $n$  equal factors, or, in other words, if we are raising  $r(\cos \phi + \sqrt{-1} \cdot \sin \phi)$  to the

$n$ th power,  $n$  being a positive whole number, we shall have, by Art. 22,

$$[r(\cos \phi + \sqrt{-1} \cdot \sin \phi)]^n = r^n(\cos n\phi + \sqrt{-1} \cdot \sin n\phi). \quad [1]$$

If  $r$  is unity, we have merely to multiply the argument by  $n$ , without changing the modulus; so that in this case increasing the exponent by unity amounts to rotating the point representing the imaginary through an angle equal to  $\phi$  without changing its distance from the origin.

25. Since extracting a root is the inverse of raising to a power,

$$\sqrt[n]{r(\cos \phi + \sqrt{-1} \cdot \sin \phi)} = \sqrt[n]{r} \left( \cos \frac{\phi}{n} + \sqrt{-1} \cdot \sin \frac{\phi}{n} \right); \quad [1]$$

for, by Art. 24,

$$\left[ \sqrt[n]{r} \left( \cos \frac{\phi}{n} + \sqrt{-1} \cdot \sin \frac{\phi}{n} \right) \right]^n = r(\cos \phi + \sqrt{-1} \cdot \sin \phi).$$

#### EXAMPLE.

Show that Art. 24 [1] holds even when  $n$  is negative or fractional.

26. As the *modulus of every quantity*, positive, negative, real, or imaginary, is *positive*, it is always possible to find the modulus of any required root; and as this modulus must be real and positive, it can never, in any given example, have more than one value. We know from algebra, however, that every equation of the  $n$ th degree containing one unknown has  $n$  roots, and that consequently every number must have  $n$   $n$ th roots. Our formula, Art. 25 [1], appears to give us but one  $n$ th root for any given quantity. It must then be incomplete.

We have seen (Art. 20) that while the modulus of a given imaginary has but one value, its argument is indeterminate and may have any one of an infinite number of values which differ by multiples of  $2\pi$ . If  $\phi_0$  is one of these values, the full form of

the imaginary is not  $r(\cos \phi_0 + \sqrt{-1} \cdot \sin \phi_0)$ , as we have hitherto written it, but is

$$r[\cos(\phi_0 + 2m\pi) + \sqrt{-1} \cdot \sin(\phi_0 + 2m\pi)],$$

where  $m$  is zero or any whole number positive or negative. Since angles differing by multiples of  $2\pi$  have the same trigonometric functions, it is easily seen that the introduction of the term  $2m\pi$  into the argument of an imaginary will not modify any of our results except that of Art. 25, which becomes

$$\begin{aligned} & \sqrt[n]{r[\cos(\phi_0 + 2m\pi) + \sqrt{-1} \cdot \sin(\phi_0 + 2m\pi)]} \\ &= \sqrt[n]{r} \left[ \cos\left(\frac{\phi_0}{n} + m\frac{2\pi}{n}\right) + \sqrt{-1} \cdot \sin\left(\frac{\phi_0}{n} + m\frac{2\pi}{n}\right) \right]. \quad [1] \end{aligned}$$

Giving  $m$  the values 0, 1, 2, 3 ....,  $n-1$ ,  $n$ ,  $n+1$ , successively, we get

$$\begin{aligned} & \frac{\phi_0}{n}, \quad \frac{\phi_0}{n} + \frac{2\pi}{n}, \quad \frac{\phi_0}{n} + 2\frac{2\pi}{n}, \quad \frac{\phi_0}{n} + 3\frac{2\pi}{n} \dots, \quad \frac{\phi_0}{n} + (n-1)\frac{2\pi}{n}, \\ & \frac{\phi_0}{n} + 2\pi, \quad \frac{\phi_0}{n} + \frac{2\pi}{n} + 2\pi, \end{aligned}$$

as arguments of our  $n$ th root.


Of these values the first  $n$ , that is, all except the last two, correspond to different points, and therefore to different roots; the next to the last gives the same point as the first, and the last the same point as the second, and it is easily seen that if we go on increasing  $m$  we shall get no new points. The same thing is true of negative values of  $m$ .

Hence we see that *every quantity, real or imaginary, has  $n$  distinct  $n$ th roots, all having the same modulus, but with arguments differing by multiples of  $\frac{2\pi}{n}$ .*

27. Any *positive real* differs from unity only in its modulus, and any *negative real* differs from  $-1$  only in its modulus. All the  $n$ th roots of any number or of its negative may be obtained

by multiplying the  $n$ th roots of 1 or of  $-1$  by the real positive  $n$ th root of the number.

Let us consider some of the roots of 1 and of  $-1$ ; for example, the cube roots of 1 and of  $-1$ . The modulus of 1 is 1, and its argument is 0. The modulus of each of the cube roots of 1 is 1, and their arguments are  $0$ ,  $\frac{2\pi}{3}$ , and  $\frac{4\pi}{3}$ ; that is,  $0^\circ$ ,  $120^\circ$ , and  $240^\circ$ . The roots in question, then, are represented by the points  $P_1$ ,  $P_2$ ,  $P_3$ , in the figure. Their values are




$$1(\cos 0 + \sqrt{-1} \cdot \sin 0),$$

$$1(\cos 120^\circ + \sqrt{-1} \cdot \sin 120^\circ),$$

$$\text{and } 1(\cos 240^\circ + \sqrt{-1} \cdot \sin 240^\circ),$$

$$\text{or } 1, -\frac{1}{2} + \frac{\sqrt{3}}{2} \sqrt{-1}, -\frac{1}{2} - \frac{\sqrt{3}}{2} \sqrt{-1}.$$

The modulus of  $-1$  is 1, and its argument is  $\pi$ . The modulus of the cube roots of  $-1$  is 1, and their arguments are  $\frac{\pi}{3}$ ,  $\frac{\pi}{3} + \frac{2\pi}{3}$ ,  $\frac{\pi}{3} + \frac{4\pi}{3}$ , that is,  $60^\circ$ ,  $180^\circ$ ,  $300^\circ$ . The roots in question, then, are represented by the points  $P_1$ ,  $P_2$ ,  $P_3$ , in the figure. Their values are



$\frac{1}{2} + \frac{\sqrt{3}}{2} \sqrt{-1}$ ,  $-1$ ,  $\frac{1}{2} - \frac{\sqrt{3}}{2} \sqrt{-1}$ .

#### EXAMPLES.

(1) What are the square roots of 1 and  $-1$ ? the 4th roots? the 5th roots? the 6th roots?

(2) Find the cube roots of  $-8$ ; the 5th roots of 32.

(3) Show that an imaginary can have no real  $n$ th root; that a positive real has two real  $n$ th roots if  $n$  is even, one if  $n$  is odd; that a negative real has one real  $n$ th root if  $n$  is odd, none if  $n$  is even.

28. Imaginaries having equal moduli, and arguments differing only in sign, are called *conjugate imaginaries*.

$r(\cos \phi + \sqrt{-1} \cdot \sin \phi)$ , and  $r[\cos(-\phi) + \sqrt{-1} \cdot \sin(-\phi)]$ , or  $r(\cos \phi - \sqrt{-1} \cdot \sin \phi)$  are *conjugate*.

They can be written  $x + y\sqrt{-1}$  and  $x - y\sqrt{-1}$ , and we see that the points corresponding to them have the same abscissa, and ordinates which are equal with opposite signs.

#### EXAMPLES.

(1) Prove that *conjugate imaginaries* have a real sum and a real product.

(2) Prove, by considering in detail the substitution of  $a + b\sqrt{-1}$  and  $a - b\sqrt{-1}$  in turn for  $x$  in any algebraic polynomial in  $x$  with real coefficients, that if any algebraic equation with real coefficients has an imaginary root the *conjugate* of that root is also a root of the equation.

(3) Prove that if in any fraction where the numerator and denominator are rational algebraic polynomials in  $x$ , we substitute  $a + b\sqrt{-1}$  and  $a - b\sqrt{-1}$  in turn for  $x$ , the results are conjugate.

#### *Transcendental Functions of Imaginaries.*

29. We have adopted a definition of an *imaginary* and laid down rules to govern its use, that enable us to deal with it, in all expressions involving only algebraic operations, precisely as if it were a quantity. If we are going further, and are to subject it to *transcendental* operations, we must carefully define each function that we are going to use, and establish the rules which the function must obey.

The principal *transcendental* functions are  $e^x$ ,  $\log x$ , and  $\sin x$ , and we wish to define and study these when  $x$  is replaced by an imaginary variable  $z$ .

As our conception and treatment of imaginaries have been entirely algebraic, we naturally wish to define our transcendental

functions by the aid of algebraic functions; and since we know that the transcendental functions of a *real* variable can be expressed in terms of algebraic functions only by the aid of infinite series, we are led to use such series in defining transcendental functions of an *imaginary* variable; but we must first establish a proposition concerning the convergency of a series containing imaginary terms.

30. *If the moduli of the terms of a series containing imaginary terms form a convergent series, the given series is convergent.*

Let  $u_0 + u_1 + u_2 + \dots + u_n + \dots$  be a series containing imaginary terms.

Let

$$u_0 = R_0(\cos \Phi_0 + \sqrt{-1} \cdot \sin \Phi_0), u_1 = R_1(\cos \Phi_1 + \sqrt{-1} \cdot \sin \Phi_1), \text{ \&c.,}$$

and suppose that the series  $R_0 + R_1 + R_2 + \dots + R_n + \dots$  is convergent; then will the series  $u_0 + u_1 + u_2 + \dots$  be convergent.

The series  $R_0 + R_1 + \dots$  is a convergent series composed of positive terms; if then we break up this series into parts in any way, each part will have a definite sum or will approach a definite limit as the number of terms considered is increased indefinitely.

The series  $u_0 + u_1 + u_2 + \dots + u_n + \dots$  can be broken up into the two series

$$R_0 \cos \Phi_0 + R_1 \cos \Phi_1 + R_2 \cos \Phi_2 + \dots + R_n \cos \Phi_n + \dots \quad (1)$$

and

$$\sqrt{-1}(R_0 \sin \Phi_0 + R_1 \sin \Phi_1 + R_2 \sin \Phi_2 + \dots + R_n \sin \Phi_n + \dots). \quad (2)$$

(1) can be separated into two parts, the first made up only of positive terms, the second only of negative terms, and can therefore be regarded as the difference between two series, each consisting of positive terms. Each term in either series will be a term of the modulus series  $R_0 + R_1 + R_2 + \dots$  multiplied by a quantity less than one, and the sum of  $n$  terms of each series will therefore approach a definite limit, as  $n$  increases indefinitely. The series (1), then, which is the abscissa of the point representing the given imaginary series, has a finite sum.

In the same way it may be shown that the coefficient of  $\sqrt{-1}$  in (2) has a finite sum, and this is the ordinate of the point representing the given series. The sum of  $n$  terms of the given series, then, approaches a definite limit as  $n$  is increased indefinitely, and the series is convergent.

31. We have seen (I. Art. 133 [2]) that

$$e^x = 1 + \frac{x}{1} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots \quad [1]$$

when  $x$  is real, and that this series is convergent for all values of  $x$ .

Let us define  $e^z$ , where  $z = x + y\sqrt{-1}$ , by the series

$$e^z = 1 + \frac{z}{1} + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} + \dots \quad [2]$$

This series is convergent, for if  $z = r(\cos \phi + \sqrt{-1} \cdot \sin \phi)$  the series

$$1 + \frac{r}{1} + \frac{r^2}{2!} + \frac{r^3}{3!} + \frac{r^4}{4!} + \dots$$

made up of the moduli of the terms of [2] is convergent by I. Art. 133, and therefore the value we have chosen for  $e^z$  is a determinate finite one.

Write  $x + y\sqrt{-1}$  for  $z$ , and we get

$$e^{x+y\sqrt{-1}} = 1 + \frac{x+y\sqrt{-1}}{1} + \frac{(x+y\sqrt{-1})^2}{2!} + \frac{(x+y\sqrt{-1})^3}{3!} + \dots \quad [3]$$

The terms of this series can be expanded by the Binomial Theorem. Consider all the resulting terms containing any given power of  $x$ , say  $x^p$ ; we have

$$\frac{x^p}{p!} \left( 1 + \frac{y\sqrt{-1}}{1} + \frac{(y\sqrt{-1})^2}{2!} + \frac{(y\sqrt{-1})^3}{3!} + \dots + \frac{(y\sqrt{-1})^n}{n!} + \dots \right);$$

or, separating the real terms and the imaginary terms,

$$\begin{aligned} & \frac{x^p}{p!} \left( 1 - \frac{y^2}{2!} + \frac{y^4}{4!} - \frac{y^6}{6!} + \dots \right) \\ & + \frac{x^p}{p!} \sqrt{-1} \left( y - \frac{y^3}{3!} + \frac{y^5}{5!} - \frac{y^7}{7!} + \dots \right), \end{aligned}$$



or  $\frac{x^p}{p!} (\cos y + \sqrt{-1} \cdot \sin y)$ , by I. Art. 134.

Giving  $p$  all values from 1 to  $\infty$  we get

$$e^{x + y\sqrt{-1}} = (\cos y + \sqrt{-1} \cdot \sin y) \left(1 + \frac{x}{1} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots\right) \\ = e^x (\cos y + \sqrt{-1} \cdot \sin y), \quad [4]$$

which, by the way, is in one of our typical imaginary forms.

If  $x = 0$ , in [4],

$$\text{we get } e^{y\sqrt{-1}} = \cos y + \sqrt{-1} \cdot \sin y, \quad [5]$$

which suggests a new way of writing our typical imaginary; namely,

$$r (\cos \phi + \sqrt{-1} \cdot \sin \phi) = r e^{\phi\sqrt{-1}}.$$

32. We have seen that

$$e^{x+y\sqrt{-1}} = e^x e^{y\sqrt{-1}};$$

let us see if all imaginary powers of  $e$  obey the *law of indices*; that is, if the equation

$$e^u e^v = e^{u+v} \quad [1]$$

is universally true.

$$\text{Let } u = x_1 + y_1\sqrt{-1} \text{ and } v = x_2 + y_2\sqrt{-1},$$

$$\text{then } e^u = e^{x_1 + y_1\sqrt{-1}} = e^{x_1} (\cos y_1 + \sqrt{-1} \cdot \sin y_1),$$

$$e^v = e^{x_2 + y_2\sqrt{-1}} = e^{x_2} (\cos y_2 + \sqrt{-1} \cdot \sin y_2),$$

$$e^u e^v = e^{x_1} e^{x_2} [\cos(y_1 + y_2) + \sqrt{-1} \cdot \sin(y_1 + y_2)]$$

$$= e^{x_1 + x_2} [\cos(y_1 + y_2) + \sqrt{-1} \cdot \sin(y_1 + y_2)]$$

$$= e^{x_1 + x_2 + (y_1 + y_2)\sqrt{-1}}$$

$$= e^{u+v},$$

and the *fundamental property of exponential functions holds for imaginaries as well as for reals.*

EXAMPLE.

Prove that  $a^u a^v = a^{u+v}$  when  $u$  and  $v$  are imaginary.

*Logarithmic Functions.*

33. As a logarithm is the inverse of an exponential, we ought to be able to obtain the logarithm of an imaginary from the formula for  $e^{x+r\sqrt{-1}}$ . We see readily that

$$z = r(\cos \phi + \sqrt{-1} \cdot \sin \phi) = e^{\log r + \phi \sqrt{-1}},$$

whence  $\log z = \log r + \phi \sqrt{-1}$ ;

or, more strictly, since

$$z = r[\cos(\phi_0 + 2n\pi) + \sqrt{-1} \cdot \sin(\phi_0 + 2n\pi)],$$

$$\log z = \log r + (\phi_0 + 2n\pi) \sqrt{-1} \quad [1]$$

where  $n$  is any integer.

If  $z = x + y\sqrt{-1}$ ,  $r = \sqrt{x^2 + y^2}$ , and  $\phi = \tan^{-1} \frac{y}{x}$ ;

whence  $\log z = \frac{1}{2} \log(x^2 + y^2) + \sqrt{-1} \cdot \tan^{-1} \frac{y}{x}$ . [2]

Each of the expressions for  $\log z$  is indeterminate, and represents an infinite number of values, differing by multiples of  $2\pi\sqrt{-1}$ .

This indeterminateness in the logarithm might have been expected *a priori*, for

$$e^{2\pi\sqrt{-1}} = \cos 2\pi + \sqrt{-1} \cdot \sin 2\pi = 1, \quad \text{by Art. 31.}$$

Hence, adding  $2\pi\sqrt{-1}$  to the logarithm of any quantity will have the effect of multiplying the quantity by 1, and therefore will not change its value.

**EXAMPLE.**

Show that if an expression is imaginary, all its logarithms are imaginary; if it is real and positive, one logarithm is real and the rest imaginary; if it is real and negative, all are imaginary.

*Trigonometric Functions.*

34. If  $z$  is real,

$$\sin z = z - \frac{z^3}{3!} + \frac{z^5}{5!} - \frac{z^7}{7!} + \dots \quad [1]$$

$$\cos z = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \frac{z^6}{6!} + \dots \quad [2]$$

by I. Art. 134.

If  $z = r(\cos \phi + \sqrt{-1} \cdot \sin \phi)$ ,

the series of the moduli,

$$r + \frac{r^3}{3!} + \frac{r^5}{5!} + \frac{r^7}{7!} + \dots,$$

$$1 + \frac{r^2}{2!} + \frac{r^4}{4!} + \frac{r^6}{6!} + \dots,$$

are easily seen to be convergent; therefore if  $z$  is imaginary, the series [1] and [2] are convergent. We shall take them as definitions of the sine and cosine of an imaginary.

## EXAMPLE.

From the formulas of Art. 31, and from Art. 34 [1] and [2], show that

$$e^{z\sqrt{-1}} = \cos z + \sqrt{-1} \cdot \sin z,$$

and  $e^{-z\sqrt{-1}} = \cos z - \sqrt{-1} \cdot \sin z$ , for all values of  $z$ .

35. From the relations

$$e^{s\sqrt{-1}} = \cos z + \sqrt{-1} \cdot \sin z,$$

$$e^{-s\sqrt{-1}} = \cos z - \sqrt{-1} \cdot \sin z,$$

we get

$$\cos z = \frac{e^{s\sqrt{-1}} + e^{-s\sqrt{-1}}}{2}, \quad [1]$$

$$\sin z = \frac{e^{s\sqrt{-1}} - e^{-s\sqrt{-1}}}{2\sqrt{-1}}, \quad [2]$$

for all values of  $s$ .

Let  $z = x + y\sqrt{-1}$ .

$$\begin{aligned}\cos(x+y\sqrt{-1}) &= \frac{e^{x\sqrt{-1}-y} + e^{-x\sqrt{-1}+y}}{2} \\ &= \frac{(\cos x + \sqrt{-1} \cdot \sin x)e^{-y} + (\cos x - \sqrt{-1} \cdot \sin x)e^y}{2}, \\ &\qquad\qquad\qquad \text{by Art. 34, Ex.,} \\ &= \cos x \frac{e^y + e^{-y}}{2} - \sqrt{-1} \cdot \sin x \frac{e^y - e^{-y}}{2}. \quad [3]\end{aligned}$$

In the same way it may be shown that

$$\begin{aligned}\sin(x+y\sqrt{-1}) &= \frac{(\cos x + \sqrt{-1} \cdot \sin x)e^{-y} - (\cos x - \sqrt{-1} \cdot \sin x)e^y}{2\sqrt{-1}} \\ &= \sin x \frac{e^y + e^{-y}}{2} + \sqrt{-1} \cdot \cos x \frac{e^y - e^{-y}}{2}. \quad [4]\end{aligned}$$

If  $z$  is real in [1] and [2], we have

$$\begin{aligned}\cos x &= \frac{e^{x\sqrt{-1}} + e^{-x\sqrt{-1}}}{2}, \\ \sin x &= -\frac{e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}}{2}\sqrt{-1}.\end{aligned}$$

If  $z = y\sqrt{-1}$ , and is a pure imaginary,

$$\cos y\sqrt{-1} = \frac{e^y + e^{-y}}{2}, \quad [5]$$

$$\sin y\sqrt{-1} = \frac{e^y - e^{-y}}{2}\sqrt{-1}; \quad [6]$$

whence we see that the cosine of a pure imaginary is real, while its sine is imaginary.

By the aid of [5] and [6], [3] and [4] can be written :

$$\cos(x+y\sqrt{-1}) = \cos x \cos y\sqrt{-1} - \sin x \sin y\sqrt{-1}, \quad [7]$$

$$\sin(x+y\sqrt{-1}) = \sin x \cos y\sqrt{-1} + \cos x \sin y\sqrt{-1}. \quad [8]$$

## EXAMPLES.

(1) From [1] and [2] show that  $\sin^2 z + \cos^2 z = 1$ .

(2) Prove that

$$\cos(u + v) = \cos u \cos v - \sin u \sin v,$$

$$\sin(u + v) = \sin u \cos v + \cos u \sin v,$$

where  $u$  and  $v$  are imaginary.

The relations to be proved in examples (1) and (2) are the fundamental formulas of Trigonometry, and they enable us to use trigonometric functions of imaginaries precisely as we use trigonometric functions of reals.

*Differentiation of Functions of Imaginaries.*

36. A function of an imaginary variable,

$$z = x + y\sqrt{-1},$$

is, strictly speaking, a function of two independent variables,  $x$  and  $y$ ; for we can change  $z$  by changing either  $x$  or  $y$ , or both  $x$  and  $y$ . Its differential will usually contain  $dx$  and  $dy$ , and not necessarily  $dz$ ; and if we divide its differential by  $dz$  to get its derivative with respect to  $z$ , the result will generally contain  $\frac{dy}{dx}$ , which will be wholly indeterminate, since  $x$  and  $y$  are entirely independent in the expression  $x + y\sqrt{-1}$ . It may happen, however, in the case of some simple functions, that  $dz$  will appear as a factor in the differential of the function, which in that case will have a single derivative.

37. In differentiating, the  $\sqrt{-1}$  may be treated like a constant; for the operation of finding the differential of a function is an algebraic operation, and in all algebraic operations  $\sqrt{-1}$  obeys the same laws as any constant.

EXAMPLE.

Prove that  $d(x^2\sqrt{-1}) = 2x\sqrt{-1}.dx$ ;  
and that  $d\sqrt{-1}. \sin x = \sqrt{-1}. \cos x.dx$ .

We have, by the aid of this principle,

if 
$$z = x + y\sqrt{-1},$$

$$dz = dx + \sqrt{-1}.dy; \tag{1}$$

if 
$$z = r(\cos \phi + \sqrt{-1}. \sin \phi),$$

$$dz = dr(\cos \phi + \sqrt{-1}. \sin \phi) + rd\phi(-\sin \phi + \sqrt{-1}. \cos \phi)$$

$$= (dr + r\sqrt{-1}.d\phi)(\cos \phi + \sqrt{-1}. \sin \phi). \tag{2}$$

38. Let us now consider the differentiation of  $z^m$ ,  $e^z$ ,  $\log z$ ,  $\sin z$ , and  $\cos z$ .

Let 
$$z = r(\cos \phi + \sqrt{-1}. \sin \phi),$$

then

$$z^m = r^m(\cos m\phi + \sqrt{-1}. \sin m\phi), \tag{by Art. 24 [1];}$$

$$dz^m = mr^{m-1}dr(\cos m\phi + \sqrt{-1}. \sin m\phi) + mr^m d\phi(-\sin m\phi + \sqrt{-1}. \cos m\phi),$$

$$dz^m = mr^{m-1}[\cos(m-1)\phi + \sqrt{-1}. \sin(m-1)\phi](\cos \phi + \sqrt{-1}. \sin \phi)dr$$

$$+ mr^m[\cos(m-1)\phi + \sqrt{-1}. \sin(m-1)\phi](\cos \phi + \sqrt{-1}. \sin \phi)\sqrt{-1}.d\phi,$$

$$dz^m = mr^{m-1}[\cos(m-1)\phi + \sqrt{-1}. \sin(m-1)\phi](dr + r\sqrt{-1}.d\phi)(\cos \phi + \sqrt{-1}. \sin \phi),$$

$$dz^m = mz^{m-1}dz, \tag{1} \text{ by Art. 37 [2],}$$

$$\frac{dz^m}{dz} = mz^{m-1}, \tag{2}$$

and a power of an imaginary variable has a single derivative.

39. If  $z = x + y\sqrt{-1}$ ,  
 $e^z = e^x(\cos y + \sqrt{-1} \cdot \sin y)$ , by Art. 31 [4],  
 $de^z = e^x dx(\cos y + \sqrt{-1} \cdot \sin y) + e^x(-\sin y$   
 $+ \sqrt{-1} \cdot \cos y) dy$ ,  
 $de^z = e^x(\cos y + \sqrt{-1} \cdot \sin y) (dx + \sqrt{-1} \cdot dy)$ ,  
 $de^z = e^z dz$ , [1]  
 $\frac{de^z}{dz} = e^z$ . [2]

## EXAMPLE.

Show that  $da^z = a^z \log a \cdot dz$ .

40. If  $z = r(\cos \phi + \sqrt{-1} \cdot \sin \phi)$ ,  
 $\log z = \log r + \phi \sqrt{-1}$ , by Art. 33,  
 $d \log z = \frac{dr}{r} + \sqrt{-1} \cdot d\phi = \frac{dr + r \sqrt{-1} \cdot d\phi}{r}$ ,  
 $d \log z = \frac{(dr + r \sqrt{-1} \cdot d\phi)(\cos \phi + \sqrt{-1} \cdot \sin \phi)}{r(\cos \phi + \sqrt{-1} \cdot \sin \phi)}$ ,  
 $d \log z = \frac{dz}{z}$ , [1]  
 $\frac{d \log z}{dz} = \frac{1}{z}$ . [2]

41.  $\sin z = \frac{e^{z\sqrt{-1}} - e^{-z\sqrt{-1}}}{2\sqrt{-1}}$ , by Art. 35 [2].  
 $d \sin z = \frac{e^{z\sqrt{-1}} + e^{-z\sqrt{-1}}}{2\sqrt{-1}} \sqrt{-1} \cdot dz$   
 $= \frac{e^{z\sqrt{-1}} + e^{-z\sqrt{-1}}}{2} dz$ , by Art. 35 [1],  
 $d \sin z = \cos z \cdot dz$ . [1]

$$\begin{aligned}\cos z &= \frac{e^{z\sqrt{-1}} + e^{-z\sqrt{-1}}}{2}, \\ d \cos z &= \frac{e^{z\sqrt{-1}} - e^{-z\sqrt{-1}}}{2} \sqrt{-1} . dz = -\frac{e^{z\sqrt{-1}} - e^{-z\sqrt{-1}}}{2\sqrt{-1}} dz, \\ d \cos z &= -\sin z . dz. \quad [2]\end{aligned}$$

42. We see, then, that we get the same formulas for the differentiation of simple functions of imaginaries as for the differentiation of the corresponding functions of reals. It follows that our formulas for direct integration (I. Art. 74) hold when  $x$  is imaginary.

#### *Hyperbolic Functions.*

43. We have (Art. 35 [5] and [6])

$$\cos x \sqrt{-1} = \frac{e^x + e^{-x}}{2},$$

and

$$\sin x \sqrt{-1} = \frac{e^x - e^{-x}}{2} \sqrt{-1},$$

where  $x$  is real.  $\frac{e^x + e^{-x}}{2}$  is called the hyperbolic cosine of  $x$ , and is written  $\cosh x$ ; and  $\frac{e^x - e^{-x}}{2}$  is called the hyperbolic sine of  $x$ , and is written  $\sinh x$ ;

$$\sinh x = \frac{e^x - e^{-x}}{2} = -\sqrt{-1} . \sin x \sqrt{-1}, \quad [1]$$

$$\cosh x = \frac{e^x + e^{-x}}{2} = \cos x \sqrt{-1}. \quad [2]$$

The hyperbolic tangent is defined as the ratio of  $\sinh$  to  $\cosh$ ; and the hyperbolic cotangent, secant, and cosecant are the reciprocals of the  $\tanh$ ,  $\cosh$ , and  $\sinh$  respectively.

These functions, which are real when  $x$  is real, resemble in their properties the ordinary trigonometric functions.



44. For example,

$$\cosh^2 x - \sinh^2 x = 1; \quad [1]$$

for 
$$\cosh^2 x = \frac{e^{2x} + 2 + e^{-2x}}{4},$$

and 
$$\sinh^2 x = \frac{e^{2x} - 2 + e^{-2x}}{4}.$$

EXAMPLES.

- (1) Prove that  $1 - \tanh^2 x = \operatorname{sech}^2 x$ .  
 (2) Prove that  $1 - \operatorname{ctnh}^2 x = -\operatorname{csch}^2 x$ .  
 (3) Prove that  $\sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y$ .  
 (4) Prove that  $\cosh(x + y) = \cosh x \cosh y + \sinh x \sinh y$ .

45. 
$$d \sinh x = d \frac{e^x - e^{-x}}{2} = \frac{e^x + e^{-x}}{2} dx,$$

$$d \sinh x = \cosh x \cdot dx.$$

EXAMPLES.

- (1) Prove  $d \cosh x = \sinh x \cdot dx$ .  
 $d \tanh x = \operatorname{sech}^2 x \cdot dx$ .  
 $d \operatorname{ctnh} x = -\operatorname{csch}^2 x \cdot dx$ .  
 $d \operatorname{sech} x = -\operatorname{sech} x \tanh x \cdot dx$ .  
 $d \operatorname{csch} x = -\operatorname{csch} x \operatorname{ctnh} x \cdot dx$ .

46. We can deal with anti-hyperbolic functions just as with anti-trigonometric functions.

To find  $d \sinh^{-1} x$ .

Let  $u = \sinh^{-1} x,$   
 then  $x = \sinh u,$   
 $dx = \cosh u \cdot du,$

$$du = \frac{dx}{\cosh u},$$

$$\cosh u = \sqrt{1 + \sinh^2 u}, \quad \text{by Art. 44 [1],}$$

$$\cosh u = \sqrt{1 + x^2},$$

$$d \sinh^{-1} x = \frac{dx}{\sqrt{1 + x^2}}. \quad [1]$$

## EXAMPLES.

Prove the formulas

$$d \cosh^{-1} x = \frac{dx}{\sqrt{x^2 - 1}}$$

$$d \tanh^{-1} x = \frac{dx}{1 - x^2}$$

$$d \operatorname{sech}^{-1} x = -\frac{dx}{x \sqrt{1 - x^2}}$$

$$d \operatorname{csch}^{-1} x = -\frac{dx}{x \sqrt{x^2 + 1}}$$

47. The anti-hyperbolic functions are easily expressed as logarithms.

Let

$$u = \sinh^{-1} x,$$

then

$$x = \sinh u = \frac{e^u - e^{-u}}{2},$$

$$2x = e^u - \frac{1}{e^u},$$

$$2xe^u = e^{2u} - 1,$$

$$e^{2u} - 2xe^u = 1,$$

$$e^{2u} - 2xe^u + x^2 = 1 + x^2,$$

$$e^u - x = \pm \sqrt{1 + x^2},$$

$$e^u = x \pm \sqrt{1 + x^2};$$

as  $e^u$  is necessarily positive, we may reject the negative value in the second member as impossible, and we have

$$e^u = x + \sqrt{1+x^2},$$

$$u = \log(x + \sqrt{1+x^2}),$$

$$\text{or} \quad \sinh^{-1} x = \log(x + \sqrt{1+x^2}). \quad [1]$$

EXAMPLES.

Prove the formulas

$$\cosh^{-1} x = \log(x + \sqrt{x^2 - 1}).$$

$$\tanh^{-1} x = \frac{1}{2} \log \frac{1+x}{1-x}.$$

$$\operatorname{sech}^{-1} x = \log\left(\frac{1}{x} + \sqrt{\frac{1}{x^2} - 1}\right).$$

$$\operatorname{cosech}^{-1} x = \log\left(\frac{1}{x} + \sqrt{\frac{1}{x^2} + 1}\right).$$

48. The principal advantage arising from the use of hyperbolic functions is that they bring to light some curious analogies between the integrals of certain irrational functions.

From I. Art. 71 we obtain the formulas for direct integration.

$$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x. \quad [1]$$

$$\int \frac{dx}{1+x^2} = \tan^{-1} x. \quad [2]$$

$$\int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x. \quad [3]$$

From Art. 46 we obtain the allied formulas :

$$\int \frac{dx}{\sqrt{1+x^2}} = \sinh^{-1} x = \log(x + \sqrt{1+x^2}). \quad [4]$$

$$\int \frac{dx}{\sqrt{x^2-1}} = \cosh^{-1} x = \log(x + \sqrt{x^2-1}). \quad [5]$$

$$\int \frac{dx}{1-x^2} = \tanh^{-1}x = \frac{1}{2} \log \frac{1+x}{1-x}. \quad [6]$$

$$-\int \frac{dx}{x\sqrt{1-x^2}} = \operatorname{sech}^{-1}x = \log\left(\frac{1}{x} + \sqrt{\frac{1}{x^2} - 1}\right). \quad [7]$$

$$-\int \frac{dx}{x\sqrt{x^2+1}} = \operatorname{csch}^{-1}x = \log\left(\frac{1}{x} + \sqrt{\frac{1}{x^2} + 1}\right). \quad [8]$$

## CHAPTER III.

## GENERAL METHODS OF INTEGRATING.

49. We have defined the *integral* of any function of a single variable as *the function which has the given function for its derivative* (I. Art. 53); we have defined a *definite integral* as *the limit of the sum of a set of differentials*; and we have shown that a definite integral is the *difference between two values of an ordinary integral* (I. Art. 183).

Now that we have adopted the differential notation in place of the derivative notation, it is better to regard an integral as the inverse of a *differential* instead of as the inverse of a *derivative*. Hence the integral of  $fx.dx$  will be the function whose differential is  $fx.dx$ ; and we shall indicate it by  $\int fx.dx$ . In our old notation we should have indicated precisely the same function by  $\int_x fx$ ; for if the derivative of a function is  $fx$  we know that its differential is  $fx.dx$ .

50. If  $fx$  is any function whatever of  $x$ ,  $fx.dx$  has an *integral*. For if we construct the curve whose equation is  $y = fx$ , we know that the area included by the curve, the axis of  $X$ , any fixed ordinate, and the ordinate corresponding to the variable  $x$ , has for its differential  $ydx$ , or, in other words,  $fx.dx$  (I. Art. 51). Such an area always exists, and it is a determinate function of  $x$ , except that, as the position of the initial ordinate is wholly arbitrary, the expression for the area will contain an arbitrary constant. Thus, if  $Fx$  is the area in question for some one position of the initial ordinate, we shall have

$$\int fx.dx = Fx + C,$$

where  $C$  is an arbitrary constant.

Moreover,  $Fx + C$  is a complete expression for  $\int fx.d x$ ; for if two functions of  $x$  have the same differential, they have the same derivative with respect to  $x$ , and therefore they change at the same rate when  $x$  changes (I. Art. 38); they can differ, then, at any instant only by the difference between their initial values, which is some constant.

Hence we see that every expression of the form  $fx.d x$  has an integral, and, except for the presence of an arbitrary constant, but one integral.

51. We have shown in I. Art. 183 that a definite integral is the difference between two values of an ordinary integral, and therefore contains no constant. Thus, if  $Fx + C$  is the integral of  $fx.d x$ ,

$$\int_a^b fx.d x = Fb - Fa.$$

In the same way we shall have

$$\int_a^b fx.d x = Fb - Fa;$$

and we see that a definite integral is a function of the values between which the sum is taken and not of the variable with respect to which we integrate.

Since

$$\int_a^b fx.d x = Fa - Fb,$$

$$\int_b^a fx.d x = - \int_a^b fx.d x.$$

EXAMPLE.

Show that  $\int_a^b fx.d x + \int_b^c fx.d x = \int_a^c fx.d x$ .

52. In what we have said concerning definite integrals we have tacitly assumed that the integral is a continuous function between the values between which the sum in question is taken. If it is not, we cannot regard the whole increment of  $Fx$  as equal


to the limit of the sum of the partial infinitesimal increments, and the reasoning of I. Art. 183 ceases to be valid.

Take, for example,  $\int_{-1}^1 \frac{dx}{x^2}$ .

$\int \frac{dx}{x^2} = \int x^{-2} dx = \frac{x^{-1}}{-1} = -\frac{1}{x}$ , by I. Art. 55 (7); and apparently

$$\int_{-1}^1 \frac{dx}{x^2} = \left(-\frac{1}{x}\right)_{x=1} - \left(-\frac{1}{x}\right)_{x=-1} = -2.$$

But  $\int_{-1}^1 \frac{dx}{x^2}$  ought to be the area between the curve  $y = \frac{1}{x^2}$ , the axis of  $x$ , and the ordinates corresponding to  $x=1$  and  $x=-1$ ,



which evidently is not  $-2$ ; and we see that the function  $\frac{1}{x^2}$  is discontinuous between the values  $x = -1$  and  $x = 1$ .

The area in question which the definite integral should represent is easily seen to be infinite, for

$$\int_{-1}^{-\epsilon} \frac{dx}{x^2} = \frac{1}{\epsilon} - 1, \text{ and } \int_{\epsilon}^1 \frac{dx}{x^2} = \frac{1}{\epsilon} - 1,$$

and each of these expressions increases without limit as  $\epsilon$  approaches zero.

53. Since a definite integral is the difference between two values of an indefinite integral, what we have to find first in any problem is the indefinite integral. This may be found by inspection if the function to be integrated comes under any of the forms we have already obtained by differentiation, and we are then said to integrate directly. Direct integration has been illustrated, and the most important of the forms which can be integrated directly have been given in I. Chapter V. For the sake of convenience we rewrite these forms, using the differential notation, and adding one or two new forms from our sections on hyperbolic functions.

$$\int x^n dx = \frac{x^{n+1}}{n+1}.$$

$$\int \frac{dx}{x} = \log x.$$

$$\int a^x dx = \frac{a^x}{\log a}.$$

$$\int e^x dx = e^x.$$

$$\int \sin x \cdot dx = -\cos x.$$

$$\int \cos x \cdot dx = \sin x.$$

$$\int \tan x \cdot dx = -\log \cos x.$$

$$\int \operatorname{ctn} x \cdot dx = \log \sin x.$$

$$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x.$$

$$\int \frac{dx}{\sqrt{1+x^2}} = \sinh^{-1} x = \log(x + \sqrt{1+x^2}).$$

$$\int \frac{dx}{\sqrt{x^2-1}} = \cosh^{-1} x = \log(x + \sqrt{x^2-1}).$$

$$\int \frac{dx}{1+x^2} = \tan^{-1} x.$$

$$\int \frac{dx}{1-x^2} = \tanh^{-1} x = \frac{1}{2} \log \frac{1+x}{1-x}.$$

$$\int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x.$$

$$\int \frac{dx}{x\sqrt{1-x^2}} = -\operatorname{sech}^{-1} x = -\log\left(\frac{1}{x} + \sqrt{\frac{1}{x^2}-1}\right).$$

$$\int \frac{dx}{x\sqrt{x^2+1}} = -\operatorname{csch}^{-1} x = -\log\left(\frac{1}{x} + \sqrt{\frac{1}{x^2}+1}\right).$$

$$\int \frac{dx}{\sqrt{2x-x^2}} = \operatorname{vers}^{-1} x.$$



54. We took up in I. Chap. V. the principal devices used in preparing a function for integration when it cannot be integrated directly.

The first of these methods, that of *integration by substitution*, is simplified by the use of the differential notation, because the formula for *change of variable* (I. Art. 75 [1]),

$$\int_x u = \int_y u D_y x \text{ becoming } \int u dx = \int u \frac{dx}{dy} dy,$$

reduces to an identity and is no longer needed, and all that is required is a simple substitution.

(a) For example, let us find  $\int \frac{dx}{x} \sqrt{1 + \log x}$ .

Let  $1 + \log x = z$ ;      then  $\frac{dx}{x} = dz$ ,

and  $\int \frac{dx}{x} \sqrt{1 + \log x} = \int z^{\frac{1}{2}} dz = \frac{2}{3} z^{\frac{3}{2}} = \frac{2}{3} (1 + \log x)^{\frac{3}{2}}$ .

When, as in this example, a factor of the quantity to be integrated is equal or proportional to the differential of some function occurring in the expression, the substitution of a new variable for the function in question will generally simplify the problem.

(b) Required  $\int \frac{dx}{e^x + e^{-x}}$ .

Let  $e^x = y$ ;      then  $e^x dx = dy$ ,

$$\frac{dx}{e^x + e^{-x}} = \frac{e^x dx}{e^{2x} + 1} = \frac{dy}{y^2 + 1},$$

and  $\int \frac{dx}{e^x + e^{-x}} = \int \frac{dy}{1 + y^2} = \tan^{-1} y = \tan^{-1} e^x$ .

(c) Required  $\int \sec x dx$ .

$$\sec x = \frac{1}{\cos x} = \frac{\cos x}{\cos^2 x}$$

Let  $z = \sin x$ ;      then  $dz = \cos x \cdot dx$ ,  
 $\cos^2 x = 1 - z^2$ ,

$$\int \frac{\cos x \cdot dx}{\cos^2 x} = \int \frac{dz}{1 - z^2} = \frac{1}{2} \log \frac{1+z}{1-z}, \quad \text{by Art. 53,}$$

$$\int \sec x \cdot dx = \frac{1}{2} \log \frac{1 + \sin x}{1 - \sin x} = \log \tan \left( \frac{\pi}{4} + \frac{x}{2} \right).$$

EXAMPLES.

Prove that (1)  $\int \csc x \cdot dx = \frac{1}{2} \log \frac{1 - \cos x}{1 + \cos x} = \log \tan \frac{x}{2}$ .

(2)  $\int \frac{x^2 dx}{\sqrt{1 - x^2}} = -\frac{1}{2} \cos^{-1} x - \frac{x\sqrt{1 - x^2}}{2}$ .

*Suggestion:* Let  $x = \cos z$ .

55. The formula for *integration by parts* (I. Art. 79 [1]) becomes

$$\int u dv = uv - \int v du, \quad [1]$$

when we use the differential notation. It is used as in I. Chap. V.

(a) For example, let us find  $\int x^n \log x \cdot dx$ .

Let  $u = \log x$ ,      and  $dv = x^n dx$ ;

then  $du = \frac{dx}{x}$ ,

and  $v = \frac{x^{n+1}}{n+1}$ ,

$$\int x^n \log x \cdot dx = \frac{x^{n+1}}{n+1} \log x - \int \frac{x^n}{n+1} dx = \frac{x^{n+1}}{n+1} \left( \log x - \frac{1}{n+1} \right).$$

(b) Required  $\int x \sin^{-1} x \cdot dx$ .

Let  $u = \sin^{-1} x$ ,      and  $dv = x dx$ ;

then  $du = \frac{dx}{\sqrt{1 - x^2}}$ ,

and

$$v = \frac{x^2}{2},$$

$$\int x \sin^{-1} x . dx = \frac{x^2}{2} \sin^{-1} x - \frac{1}{2} \int \frac{x^2 dx}{\sqrt{1-x^2}},$$

$$\int x \sin^{-1} x . dx = \frac{x^2}{2} \sin^{-1} x + \frac{1}{4} (\cos^{-1} x + x\sqrt{1-x^2}).$$

(c) Required  $\int \frac{xe^x dx}{(1+x)^2}$ .

Let  $u = xe^x$ ,

and

$$dv = \frac{dx}{(1+x)^2};$$

then

$$du = (xe^x + e^x) dx = e^x(1+x) dx,$$

and

$$v = -\frac{1}{1+x},$$

$$\int \frac{xe^x dx}{(1+x)^2} = -\frac{xe^x}{1+x} + \int e^x dx = -\frac{xe^x}{1+x} + e^x = \frac{e^x}{1+x}.$$

## EXAMPLES.

$$(1) \int \frac{dx}{\sqrt{1-3x-x^2}} = \sin^{-1} \frac{3+2x}{\sqrt{13}}.$$

$$(2) \int x \tan^{-1} x . dx = \frac{1+x^2}{2} \tan^{-1} x - \frac{1}{2} x.$$

$$(3) \int \frac{x dx}{(1-x)^3} = -\frac{1}{1-x} + \frac{1}{2(1-x)^2}.$$

$$(4) \int \frac{x dx}{\sqrt{2ax-x^2}} = -\sqrt{2ax-x^2} + a \operatorname{vers}^{-1} \frac{x}{a}.$$

$$(5) \int \sqrt{2ax-x^2} . dx = \frac{x-a}{2} \sqrt{2ax-x^2} + \frac{a^2}{2} \sin^{-1} \frac{x-a}{a}.$$

*Suggestion:* Throw  $2ax-x^2$  into the form  $a^2-(x-a)^2$ .

$$(6) \int \frac{1+\cos x}{x+\sin x} dx = \log(x+\sin x).$$

$$(7) \int \frac{x + \sin x}{1 + \cos x} dx = x \tan \frac{x}{2}.$$

*Suggestion:* Introduce  $\frac{x}{2}$  in place of  $x$ .

$$(8) \int \frac{dx}{x(\log x)^n} = -\frac{1}{(n-1)(\log x)^{n-1}}.$$

$$(9) \int \frac{\log(\log x)}{x} dx = \log x [\log(\log x) - 1].$$

$$(10) \int \frac{\sin^{-1} x \cdot dx}{(1-x^2)^{\frac{3}{2}}} = z \tan z + \log \cos z, \text{ where } z = \sin^{-1} x.$$

$$(11) \int \frac{dx}{\sin x + \cos x} = \frac{1}{\sqrt{2}} \log \tan \left( \frac{x}{2} + \frac{\pi}{8} \right).$$

$$(12) \int \frac{\sin x \, dx}{a + b \cos x} = -\frac{\log(a + b \cos x)}{b}.$$

$$(13) \int \frac{dx}{x^2 + 4x + 5} = \tan^{-1}(x + 2).$$

$$(14) \int \frac{x^2 dx}{1-x^2} = \frac{1}{6} \log \left( \frac{1+x^2}{1-x^2} \right).$$

$$(15) \int \frac{x^3 dx}{x^2 - x^4 - 6} = \frac{1}{20} \log \left( \frac{x^4 - 3}{x^4 + 2} \right).$$

$$(16) \int \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x} = \frac{1}{ab} \tan^{-1} \left( \frac{b}{a} \tan x \right).$$

## CHAPTER IV.

## RATIONAL FRACTIONS.

56. We shall now attempt to consider systematically the methods of integrating various functions; and to this end we shall begin with *rational algebraic expressions*. Any rational algebraic polynomial can be integrated immediately by the aid of the formula

$$\int x^n dx = \frac{x^{n+1}}{n+1}.$$

Take next a *rational fraction*, that is, a fraction whose numerator and denominator are rational algebraic polynomials. A rational fraction is *proper* if its numerator is of lower degree than its denominator; *improper* if the degree of the numerator is equal to or greater than the degree of the denominator. Since an improper fraction can always be reduced to a polynomial plus a proper fraction, by actually dividing the numerator by the denominator, we need only consider the treatment of proper fractions.

57. Every proper rational fraction can be reduced to the sum of a set of simpler fractions each of which has a constant for a numerator and some power of a binomial for its denominator; that is, a set of fractions any one of which is of the form  $\frac{A}{(x-a)^m}$ .

Let our given fraction be  $\frac{fx}{Fx}$ .

If  $a, b, c, \&c.$ , are the roots of the equation,

$$Fx = 0, \tag{1}$$

we have, from the Theory of Equations,

$$Fx = A(x-a)(x-b)(x-c) \dots \tag{2}$$

The equation (1) may have some equal roots, and then some of the factors in (2) will be repeated. Suppose  $a$  occurs  $p$  times as a root of (1),  $b$  occurs  $q$  times,  $c$  occurs  $r$  times, &c.,

then 
$$Fx = A(x-a)^p(x-b)^q(x-c)^r \dots \tag{3}$$

Call 
$$A(x-b)^q(x-c)^r \dots = \phi x;$$

then 
$$Fx = (x-a)^p \phi x,$$

and 
$$\begin{aligned} \frac{fx}{Fx} &= \frac{fx}{(x-a)^p \phi x} = \frac{fx - \frac{fa}{\phi a} \phi x}{(x-a)^p \phi x} + \frac{\frac{fa}{\phi a} \phi x}{(x-a)^p \phi x} \\ &= \frac{\frac{fa}{\phi a}}{(x-a)^p} + \frac{fx - \frac{fa}{\phi a} \phi x}{(x-a)^p \phi x}. \end{aligned}$$

$\frac{fx - \frac{fa}{\phi a} \phi x}{(x-a)^p \phi x}$  is a new proper fraction, but it can be reduced to a simpler form by dividing numerator and denominator by  $x-a$ , which is an exact divisor of the numerator because  $a$  is a root of the equation

$$fx - \frac{fa}{\phi a} \phi x = 0.$$

If we represent by  $f_1 x$  the quotient arising from the division of  $fx - \frac{fa}{\phi a} \phi x$  by  $x-a$ , we shall have

$$\frac{fx}{Fx} = \frac{\frac{fa}{\phi a}}{(x-a)^p} + \frac{f_1 x}{(x-a)^{p-1} \phi x},$$

where  $\frac{f_1 x}{(x-a)^{p-1} \phi x}$  is a proper fraction, and may be treated precisely as we have treated the original fraction.

Hence 
$$\frac{f_1 x}{(x-a)^{p-1} \phi x} = \frac{\frac{f_1 a}{\phi a}}{(x-a)^{p-1}} + \frac{f_2 x}{(x-a)^{p-2} \phi x}.$$

By continuing this process we shall get

$$\frac{fx}{Fx} = \frac{\frac{fa}{\phi a}}{(x-a)^p} + \frac{\frac{f_1 a}{\phi a}}{(x-a)^{p-1}} + \frac{\frac{f_2 a}{\phi a}}{(x-a)^{p-2}} + \dots + \frac{\frac{f_{p-1} a}{\phi a}}{x-a} + \frac{f_p x}{\phi x}.$$

## CHAPTER IV.

## RATIONAL FRACTIONS.

We shall now attempt to consider systematically the method of integrating rational functions: and to this end we shall first deal with the case of proper fractions. *Any rational function can be decomposed or evaluated by the aid of the following*

$$\int \frac{P}{Q} dx = \frac{P}{Q} + \dots$$

The proper fraction is defined as that is, a fraction whose numerator and denominator are rational algebraic polynomials, and whose denominator is of higher degree than its numerator, or whose denominator is of the same degree as the numerator, but whose degree of the numerator is less than the degree of the denominator. Since any such fraction can be reduced to a polynomial plus a proper fraction, the method of integration of a rational function will usually involve the treatment of proper fractions.

Any proper rational fraction can be reduced to the sum of a polynomial and a fraction which has a constant for a numerator and a linear factor for its denominator; and the latter fraction can be reduced to the form  $\frac{A}{x-a}$ .

$$\frac{P}{Q} = \frac{P}{(x-a)^2} + \dots$$

where  $a$  is one of the roots of the equation

$$Q = 0.$$

Let us suppose that  $Q$  is of the form

$$Q = (x-a)^2(x-b)(x-c) \dots$$

The equation (1) may have some equal roots and then some of the factors in (2) will be repeated. Suppose a certain  $\phi$  occurs as a root of (1); it occurs  $l$  times;  $\psi$  occurs  $m$  times; etc.,

then 
$$F(x) = \Delta(x-a)^l(x-b)^m(x-c)^n \dots \quad (3)$$

Call 
$$\Delta(x-a)^l(x-b)^m \dots = \phi(x);$$

then 
$$F(x) = (x-a)^l \phi(x),$$

and 
$$\begin{aligned} \frac{f(x)}{F(x)} &= \frac{f(x)}{(x-a)^l \phi(x)} = \frac{f(x) - \frac{f(a)}{\phi(a)} \phi(x)}{(x-a)^l \phi(x)} + \frac{\frac{f(a)}{\phi(a)} \phi(x)}{(x-a)^l \phi(x)} \\ &= \frac{\frac{f(x) - \frac{f(a)}{\phi(a)} \phi(x)}{\phi(x)}}{(x-a)^l} + \frac{\frac{f(a)}{\phi(a)}}{(x-a)^l} \end{aligned}$$

$\frac{f(x) - \frac{f(a)}{\phi(a)} \phi(x)}{(x-a)^l \phi(x)}$  is a new proper fraction, but it can be reduced to a simpler form by dividing numerator and denominator by  $x-a$ , which is an exact divisor of the numerator because  $a$  is a root of the equation

$$f(x) - \frac{f(a)}{\phi(a)} \phi(x) = 0.$$

If we represent by  $f_1(x)$  the numerator arising from the division of  $f(x) - \frac{f(a)}{\phi(a)} \phi(x)$  by  $x-a$ , we have

$$\frac{f_1(x)}{(x-a)^{l-1} \phi(x)},$$

which is a proper fraction, and may be treated as such.

$$\frac{f_1(x)}{(x-a)^{l-1} \phi(x)}$$

$$= \frac{f_{l-1}(a)}{l-1} + \frac{f_l(a)}{\phi(a)}.$$



In the same way  $\frac{f_p x}{\phi x}$  can be broken up into a set of fractions having  $(x-b)^r$ ,  $(x-b)^{r-1}$ , &c., for denominators, plus a fraction which can be broken up into fractions having  $(x-c)^r$ ,  $(x-c)^{r-1}$  ....., &c., for denominators; and we shall have, in the end,

$$\begin{aligned} \frac{f x}{F x} = & \frac{A_1}{(x-a)^p} + \frac{A_2}{(x-a)^{p-1}} + \dots + \frac{A_p}{x-a} + \frac{B_1}{(x-b)^r} \\ & + \frac{B_2}{(x-b)^{r-1}} + \dots + \frac{B_r}{x-b} + \dots + K, \end{aligned} \quad [1]$$

where  $K$  is the quotient obtained when we divide out the last factor of the denominator, and is consequently a constant. More than this,  $K$  must be zero, for as (1) is identically true, it must be true when  $x = \infty$ ; but when  $x = \infty$ ,  $\frac{f x}{F x}$  becomes zero, because its denominator is of higher degree than its numerator, and each of the fractions in the second member also becomes zero; whence  $K = 0$ .

58. Since we now know the form into which any given rational fraction can be thrown, we can determine the numerators by the aid of known properties of an identical equation.

Let it be required to break up  $\frac{3x-1}{(x-1)^2(x+1)}$  into simpler fractions.

By Art. 57,

$$\frac{3x-1}{(x-1)^2(x+1)} = \frac{A}{(x-1)^2} + \frac{B}{x-1} + \frac{C}{x+1},$$

and we wish to determine  $A$ ,  $B$ , and  $C$ . Clearing of fractions, we have

$$3x-1 = A(x+1) + B(x-1)(x+1) + C(x-1)^2. \quad (1)$$

As this equation is identically true, the coefficients of like powers of  $x$  in the two members must be equal; and we have

$$\begin{aligned} B + C &= 0, \\ A - 2C &= 3, \\ A - B + C &= -1; \end{aligned}$$

whence we find

$$\begin{aligned} A &= 1, \\ B &= 1, \\ C &= -1; \end{aligned}$$

and

$$\frac{3x-1}{(x-1)^2(x+1)} = \frac{1}{(x-1)^2} + \frac{1}{x-1} - \frac{1}{x+1}. \quad (2)$$

The labor of determining the required constants can often be lessened by simple algebraic devices.

For example; since the identical equation we start with is true for all values of  $x$ , we have a right to substitute for  $x$  values that will make terms of the equation disappear. Take equation [1]:

$$3x-1 = A(x+1) + B(x+1)(x-1) + C(x-1)^2. \quad [1]$$

Let  $x = 1$ ,

$$2 = 2A,$$

$$A = 1,$$

then

$$2x-2 = B(x+1)(x-1) + C(x-1)^2;$$

divide by  $x-1$ ,

$$2 = B(x+1) + C(x-1).$$

Let  $x = 1$ ,

$$2 = 2B,$$

$$B = 1,$$

then

$$-x+1 = C(x-1),$$

$$C = -1.$$

#### EXAMPLES.

(1) Show that when we equate the coefficients of the same powers of  $x$  on the two sides of our identical equation, we shall always have equations enough to determine all our required numerators.

(2) Break up  $\frac{9x^2+9x-128}{(x-3)^2(x+1)}$  into simpler fractions.

59. The partial fractions corresponding to any given factor of the denominator can be determined directly.

Let us suppose that the factor in question is of the first degree and occurs but once; represent it by  $x - a$ .

$$\frac{fx}{Fx} = \frac{A}{x-a} + \frac{f_1x}{\phi x}, \quad (1)$$

by Art. 57, where

$$\phi x = \frac{Fx}{x-a},$$

so that

$$Fx = (x-a)\phi x.$$

Clear (1) of fractions.

$$fx = A\phi x + (x-a)f_1x. \quad (2)$$

As (1) is an identical equation, (2) will be true for any value of  $x$ . Let  $x = a$ ,

$$fa = A\phi a,$$

$$A = \frac{fa}{\phi a}, \quad (3)$$

a result agreeing with Art. 57.

Hence, to find the numerator of the fraction corresponding to a factor  $(x-a)$  of the first degree, we have merely to strike out from the denominator of our original fraction the factor in question, and then substitute  $a$  for  $x$  in the result.

If the factor of the denominator is of the  $n$ th degree, there are  $n$  partial fractions corresponding to it. Let  $(x-a)^n$  be the factor in question.

$$\frac{fx}{Fx} = \frac{A_1}{(x-a)^n} + \frac{A_2}{(x-a)^{n-1}} + \frac{A_3}{(x-a)^{n-2}} + \dots + \frac{A_n}{x-a} + \frac{\psi x}{\phi x}, \quad (4)$$

where

$$Fx = (x-a)^n \phi x.$$

Multiply (4) by  $(x-a)^n$ , and represent  $(x-a)^n \frac{fx}{Fx}$  by  $\Phi x$ .

$$\begin{aligned} \Phi x &= A_1 + A_2(x-a) + A_3(x-a)^2 + \dots + A_n(x-a)^{n-1} \\ &\quad + \frac{\psi x}{\phi x}(x-a)^n. \end{aligned}$$

Differentiate successively both members of this identity, and put  $x = a$  after differentiation, and we get

$$\begin{aligned} A_1 &= \Phi a, \\ A_2 &= \Phi' a, \\ A_3 &= \frac{1}{2!} \Phi'' a, \\ A_4 &= \frac{1}{3!} \Phi''' a, \\ &\dots, \\ A_n &= \frac{1}{(n-1)!} \Phi^{(n-1)} a. \end{aligned}$$

Although these results form a complete solution of the problem, and one exceedingly neat in theory, the labor of getting the successive derivatives of  $\Phi x$  is so great that it is usually easier in practice to use the methods of Art. 58 when we have to deal with factors of higher degree than the first. So far as the fractions corresponding to factors of the first degree and to the highest powers of factors not of the first degree are concerned, the method of this article can be profitably combined with that of Art. 58.

60. As an example where the method of the last article applies well, consider

$$\begin{aligned} \frac{3x-1}{x(x-2)(x+1)} &= \frac{A}{x} + \frac{B}{x-2} + \frac{C}{x+1}, \\ A &= \left[ \frac{3x-1}{(x-2)(x+1)} \right]_{x=0} = \frac{1}{2}, \\ B &= \left[ \frac{3x-1}{x(x+1)} \right]_{x=2} = \frac{5}{6}, \\ C &= \left[ \frac{3x-1}{x(x-2)} \right]_{x=-1} = -\frac{4}{3}, \\ \frac{3x-1}{x(x-2)(x+1)} &= \frac{1}{2} \cdot \frac{1}{x} + \frac{5}{6} \cdot \frac{1}{x-2} - \frac{4}{3} \cdot \frac{1}{x+1}. \end{aligned} \quad [1]$$

Again, consider

$$\frac{1}{1+x^2} = \frac{1}{(x+\sqrt{-1})(x-\sqrt{-1})} = \frac{A}{x+\sqrt{-1}} + \frac{B}{x-\sqrt{-1}},$$

$$A = \left[ \frac{1}{x-\sqrt{-1}} \right]_{x=-\sqrt{-1}} = -\frac{1}{2\sqrt{-1}} = \frac{\sqrt{-1}}{2},$$

$$B = \left[ \frac{1}{x+\sqrt{-1}} \right]_{x=-\sqrt{-1}} = \frac{1}{2\sqrt{-1}} = -\frac{\sqrt{-1}}{2},$$

$$\frac{1}{1+x^2} = \frac{\sqrt{-1}}{2} \cdot \frac{1}{x+\sqrt{-1}} - \frac{\sqrt{-1}}{2} \cdot \frac{1}{x-\sqrt{-1}}. \quad [2]$$

61. Let us now consider a more difficult example, where it is worth while to combine our methods.

To break up  $\frac{x^2+1}{(x-1)^4(x^3+1)}$ .

$$\begin{aligned} x^3+1 &= (x+1)(x^2-x+1) \\ &= (x+1)\left(x-\frac{1}{2}-\frac{1}{2}\sqrt{-3}\right)\left(x-\frac{1}{2}+\frac{1}{2}\sqrt{-3}\right), \\ \frac{x^2+1}{(x-1)^4(x^3+1)} &= \frac{x^2+1}{(x-1)^4(x+1)(x^2-x+1)} = \frac{A_1}{(x-1)^4} \\ &+ \frac{A_2}{(x-1)^3} + \frac{A_3}{(x-1)^2} + \frac{A_4}{x-1} + \frac{B}{x+1} + \frac{C}{x-\frac{1}{2}-\frac{1}{2}\sqrt{-3}} \\ &+ \frac{D}{x-\frac{1}{2}+\frac{1}{2}\sqrt{-3}}. \end{aligned} \quad (1)$$

$$B = \left[ \frac{x^2+1}{(x-1)^4(x^2-x+1)} \right]_{x=-1} = \frac{1}{24},$$

$$A_1 = \left[ \frac{x^2+1}{(x+1)(x^2-x+1)} \right]_{x=1} = 1,$$

$$C = \left[ \frac{x^2+1}{(x-1)^4(x+1)\left(x-\frac{1}{2}+\frac{1}{2}\sqrt{-3}\right)} \right]_{x=\frac{1}{2}+\frac{1}{2}\sqrt{-3}} = -\frac{1}{3},$$

$$D = \left[ \frac{x^2+1}{(x-1)^4(x+1)\left(x-\frac{1}{2}-\frac{1}{2}\sqrt{-3}\right)} \right]_{x=\frac{1}{2}-\frac{1}{2}\sqrt{-3}} = -\frac{1}{3}.$$

$$\frac{-\frac{1}{3}}{x-\frac{1}{2}-\frac{1}{2}\sqrt{-3}} + \frac{-\frac{1}{3}}{x-\frac{1}{2}+\frac{1}{2}\sqrt{-3}} = -\frac{1}{3} \cdot \frac{(2x-1)}{x^2-x+1}.$$

Substitute these values and clear (1) of fractions.

$$24(x^2+1) = 24(x+1)(x^2-x+1) + 24A_2(x-1)(x+1)(x^2-x+1) \\ + 24A_3(x-1)^2(x+1)(x^2-x+1) + 24A_4(x-1)^3(x+1) \\ (x^2-x+1) + (x-1)^4(x^2-x+1) - 8(2x-1)(x-1)^4(x+1);$$

$$15x^5 - 51x^4 + 45x^3 + 6x^2 - 51x^2 + 45x - 9 = 24A_1(x-1)(x+1) \\ (x^2-x+1) + 24A_3(x-1)^2(x+1)(x^2-x+1) + 24A_4 \\ (x-1)^3(x+1)(x^2+x-1).$$

The second member of this equation is divisible by

$$(x-1)(x+1)(x^2-x+1),$$

therefore the first member must be divisible by the same quantity.

Dividing, we have

$$15x^2 - 36x + 9 = 24A_2 + 24A_3(x-1) + 24A_4(x-1)^2.$$

$$\text{Let } x = 1, \quad -12 = 24A_3,$$

$$A_3 = -\frac{1}{2},$$

and we get

$$15x^2 - 36x + 21 = 24A_2(x-1) + 24A_4(x-1)^2.$$

Divide by  $x-1$ ;

$$15x - 21 = 24A_2 + 24A_4(x-1).$$

$$\text{Let } x = 1, \quad -6 = 24A_2,$$

$$A_2 = -\frac{1}{4},$$

$$15x - 15 = 24A_4(x-1).$$

$$\text{Divide by } x-1; \quad 15 = 24A_4,$$

$$A_4 = \frac{5}{8}.$$

Hence

$$\frac{x^2+1}{(x-1)^4(x^2+1)} = \frac{1}{(x-1)^4} - \frac{1}{2} \cdot \frac{1}{(x-1)^3} - \frac{1}{4} \cdot \frac{1}{(x-1)^2} + \frac{5}{8} \cdot \frac{1}{x-1} \\ + \frac{1}{24} \cdot \frac{1}{x+1} - \frac{1}{3} \cdot \frac{1}{x-\frac{1}{2}-\frac{1}{2}\sqrt{-3}} - \frac{1}{3} \cdot \frac{1}{x-\frac{1}{2}+\frac{1}{2}\sqrt{-3}}. \quad (2)$$

62. Having shown that any rational fraction can be reduced to a sum of fractions which always come under one of the two forms  $\frac{A}{(x-a)^n}$  and  $\frac{A}{x-a}$ , it remains to show that these forms can be integrated.

To find  $\int \frac{A dx}{(x-a)^n}$ ,

let  $z = x - a$ ,

then  $dz = dx$ ,

and

$$\int \frac{A dx}{(x-a)^n} = A \int \frac{dz}{z^n} = -\frac{1}{(n-1)} \cdot \frac{A}{z^{n-1}} = -\frac{1}{(n-1)} \cdot \frac{A}{(x-a)^{n-1}}. \quad [1]$$

To find  $\int \frac{A dx}{x-a}$ ,

let  $z = x - a$ ,

then  $dz = dx$ ,

and  $\int \frac{A dx}{x-a} = A \int \frac{dz}{z} = A \log z = A \log(x-a). \quad [2]$

Turning back to Art. 58 (2), we find

$$\int \frac{(3x-1) dx}{(x-1)^2(x+1)} = \int \frac{dx}{(x-1)^2} + \int \frac{dx}{x-1} - \int \frac{dx}{x+1} = -\frac{1}{x-1} \\ + \log(x-1) - \log(x+1) = -\frac{1}{x-1} + \log \frac{x-1}{x+1}.$$

Turning to Art. 60 (1), we have

$$\int \frac{(3x-1) dx}{x(x-2)(x-1)} = \frac{1}{2} \int \frac{dx}{x} + \frac{3}{8} \int \frac{dx}{x-2} - \frac{4}{3} \int \frac{dx}{x+1} \\ = \frac{1}{2} \log x + \frac{3}{8} \log(x-2) - \frac{4}{3} \log(x+1).$$

63. If imaginary values come in when we break up our given fraction, they will disappear if we combine our results properly after integrating.

We know (Art. 28, Ex. 2) that if the denominator of our given fraction contains an imaginary factor,  $(x - a - b\sqrt{-1})^n$ , it will also contain the conjugate of that factor, namely,  $(x - a + b\sqrt{-1})^n$ . Moreover, since by Art. 59 the numerator of the partial fraction corresponding to  $(x - a - b\sqrt{-1})^n$  will be the same rational algebraic function of  $a + b\sqrt{-1}$  that the numerator of the partial fraction corresponding to  $(x - a + b\sqrt{-1})^n$  is of  $a - b\sqrt{-1}$ , these two numerators must be conjugate imaginaries by Art. 28, Ex. 3. Hence, for every fraction of the form  $\frac{A + B\sqrt{-1}}{(x - a - b\sqrt{-1})^n}$  we shall have a second of the form

$$\frac{A - B\sqrt{-1}}{(x - a + b\sqrt{-1})^n}$$

$$\int \frac{A + B\sqrt{-1}}{(x - a - b\sqrt{-1})^n} dx = -\frac{1}{(n-1)} \cdot \frac{(A + B\sqrt{-1})}{(x - a - b\sqrt{-1})^{n-1}},$$

by Art. 62 [1].

$$\int \frac{A - B\sqrt{-1}}{(x - a + b\sqrt{-1})^n} dx = -\frac{1}{(n-1)} \cdot \frac{(A - B\sqrt{-1})}{(x - a + b\sqrt{-1})^{n-1}}.$$

Let  $(x - a + b\sqrt{-1})^{n-1} = X + Y\sqrt{-1}$ ,

$X$  and  $Y$  being real functions of  $x$ ;

then  $(x - a - b\sqrt{-1})^{n-1} = X - Y\sqrt{-1}$ .

$$\begin{aligned} & \int \frac{A + B\sqrt{-1}}{(x - a - b\sqrt{-1})^n} dx + \int \frac{A - B\sqrt{-1}}{(x - a + b\sqrt{-1})^n} dx \\ &= -\frac{1}{(n-1)} \cdot \frac{(A + B\sqrt{-1})}{X - Y\sqrt{-1}} - \frac{1}{(n-1)} \cdot \frac{(A - B\sqrt{-1})}{X + Y\sqrt{-1}} \\ &= -\frac{1}{(n-1)} \cdot \frac{(2AX - 2BY)}{(x^2 - 2ax + a^2 + b^2)^{n-1}}, \end{aligned} \tag{1}$$

a result which is free from imaginaries.



If  $n = 1$ ,  
we have the pair of fractions,  $\frac{A + B\sqrt{-1}}{x - a - b\sqrt{-1}}$  and  $\frac{A - B\sqrt{-1}}{x - a + b\sqrt{-1}}$ .

$$\int \frac{A + B\sqrt{-1}}{x - a - b\sqrt{-1}} dx = (A + B\sqrt{-1}) \log(x - a - b\sqrt{-1}),$$

by Art. 62 [2],

$$\int \frac{A - B\sqrt{-1}}{x - a + b\sqrt{-1}} dx = (A - B\sqrt{-1}) \log(x - a + b\sqrt{-1});$$

$$\log(x - a - b\sqrt{-1}) = \frac{1}{2} \log[(x - a)^2 + b^2] - \sqrt{-1} \cdot \tan^{-1} \frac{b}{x - a},$$

$$\log(x - a + b\sqrt{-1}) = \frac{1}{2} \log[(x - a)^2 + b^2] + \sqrt{-1} \cdot \tan^{-1} \frac{b}{x - a}.$$

Hence 
$$\int \frac{A + B\sqrt{-1}}{x - a - b\sqrt{-1}} dx + \int \frac{A - B\sqrt{-1}}{x - a + b\sqrt{-1}} dx$$

$$= A \log[(x - a)^2 + b^2] + 2B \tan^{-1} \frac{b}{x - a}, \quad [2]$$

which is real.

The form of [2] can be modified by adding a constant.

$$\frac{\pi}{2} + \tan^{-1} \frac{b}{x - a} = \frac{\pi}{2} + \operatorname{ctn}^{-1} \frac{x - a}{b} = \frac{\pi}{2} - \operatorname{ctn}^{-1} \frac{a - x}{b} = \tan^{-1} \frac{a - x}{b}.$$

Hence 
$$A \log[(x - a)^2 + b^2] + 2B \tan^{-1} \frac{a - x}{b} \quad [3]$$

differs from [2] by the constant  $B\pi$ , and therefore is a true

value of 
$$\int \frac{A + B\sqrt{-1}}{x - a - b\sqrt{-1}} dx + \int \frac{A - B\sqrt{-1}}{x - a + b\sqrt{-1}} dx.$$

Turning back to Art. 61 (2) we find

$$\int \frac{x^2 + 1}{(x - 1)^4 (x^2 + 1)} dx = \int \frac{dx}{(x - 1)^4} - \frac{1}{2} \int \frac{dx}{(x - 1)^3} - \frac{1}{4} \int \frac{dx}{(x - 1)^2}$$

$$+ \frac{5}{8} \int \frac{dx}{x - 1} + \frac{1}{4} \int \frac{dx}{x + 1} - \frac{1}{8} \int \frac{dx}{x - \frac{1}{2} - \frac{1}{2}\sqrt{-3}} - \frac{1}{8} \int \frac{dx}{x - \frac{1}{2} + \frac{1}{2}\sqrt{-3}}$$

$$= -\frac{1}{3} \cdot \frac{1}{(x - 1)^3} + \frac{1}{4} \cdot \frac{1}{(x - 1)^2} + \frac{1}{4} \cdot \frac{1}{x - 1} + \frac{5}{8} \log(x - 1)$$

$$+ \frac{1}{4} \log(x + 1) - \frac{1}{8} \log(x^2 - x + 1).$$

## EXAMPLES.

- (1)  $\int \frac{x^2 - 3x + 3}{(x-1)(x-2)} dx = x + \log \frac{x-2}{x-1}.$
- (2)  $\int \frac{x^2 - 1}{x^2 - 4} dx = x + \frac{1}{2} \log \frac{x-2}{x+2}.$
- (3)  $\int \frac{dx}{x^2 + 1} = \tan^{-1} x.$
- (4)  $\int \frac{dx}{x^2 - 1} = \frac{1}{2} \log \frac{(x-1)^2}{x^2 + x + 1} - \frac{1}{\sqrt{3}} \tan^{-1} \frac{2x+1}{\sqrt{3}}.$
- (5)  $\int \frac{dx}{a^4 - x^4} = \frac{1}{2a^3} \tan^{-1} \frac{x}{a} + \frac{1}{4a^3} \log \frac{a+x}{a-x}.$
- (6)  $\int \frac{dx}{(x^2+1)(x^2+x+1)} = \frac{1}{2} \log \frac{x^2+x+1}{x^2+1} + \frac{1}{\sqrt{3}} \tan^{-1} \frac{2x+1}{\sqrt{3}}.$
- (7)  $\int \frac{x^2 dx}{x^4 + x^2 - 2} = \frac{1}{2} \log \frac{x-1}{x+1} + \frac{\sqrt{2}}{3} \tan^{-1} \frac{x}{\sqrt{2}}.$
- (8)  $\int \frac{x^2 - 1}{x^4 + x^2 + 1} dx = \frac{1}{2} \log \frac{x^2 - x + 1}{x^2 + x + 1}.$
- (9)  $\int \frac{dx}{(x-1)^2(x^2+1)^2} = -\frac{1}{4(x-1)} - \frac{1}{2} \log(x-1)$   
 $+ \frac{1}{2} \tan^{-1} x - \frac{1}{4(x^2+1)} + \frac{1}{2} \log(x^2+1).$
- (10)  $\int \frac{x^2 dx}{x^4 + 1} = \frac{1}{4\sqrt{2}} \log \frac{x^2 - x\sqrt{2} + 1}{x^2 + x\sqrt{2} + 1}$   
 $+ \frac{1}{2\sqrt{2}} [\tan^{-1}(x\sqrt{2} + 1) + \tan^{-1}(x\sqrt{2} - 1)].$
- (11)  $\int \frac{dx}{x^4 + 1} = \frac{1}{4\sqrt{2}} \log \frac{x^2 + x\sqrt{2} + 1}{x^2 - x\sqrt{2} + 1} + \frac{1}{2\sqrt{2}} \tan^{-1} \left( \frac{x\sqrt{2}}{1-x^2} \right).$

## CHAPTER V.

## REDUCTION FORMULAS.

64. The method given in the last chapter for the integration of rational fractions is open to the practical objection that it is often exceedingly laborious. In many cases much of the labor can be saved by making the required integration depend upon the integration of a simpler form. This is usually done by the aid of what is called a *reduction formula*.

Let the function to be integrated be of the form  $x^{m-1}(a+bx)^p$ , where  $m$ ,  $n$ , and  $p$  may be positive or negative. If they are integers, the function in question is either an *algebraic polynomial* or a *rational fraction*; if they are fractions, the expression is irrational. The formulas we shall obtain will apply to either case.

Denote  $a+bx^n$  by  $z$ ; then we want  $\int x^{m-1}z^p dx$ .

Let  $z^p = u$

and  $x^{m-1}dx = dv$ , and integrate by parts.

$$du = pz^{p-1}dz = bnpz^{n-1}z^{p-1}dx,$$

$$v = \frac{x^m}{m},$$

$$\int x^{m-1}z^p dx = \frac{x^m z^p}{m} - \frac{bnp}{m} \int x^{m+n-1}z^{p-1} dx. \quad [1]$$

This formula makes our integral depend upon the integral of an expression like the given one, except that the exponent of  $x$  has been increased while that of  $z$  has been decreased.

We get from [1], by transposition,

$$\int x^{m+n-1}z^{p-1} dx = \frac{x^m z^p}{bnp} - \frac{m}{bnp} \int x^{m-1}z^p dx.$$

Change  $m + n$  into  $m$  and  $p - 1$  into  $p$ , whence  $m$  is changed into  $m - n$  and  $p$  into  $p + 1$ , and we get

$$\int x^{m-1} z^p dx = \frac{x^{m-n} z^{p+1}}{bn(p+1)} - \frac{m-n}{bn(p+1)} \int x^{m-n-1} z^{p+1} dx, \quad [2]$$

a formula that lowers the exponent of  $x$  while it raises that of  $z$ .

Since  $z = a + bx^n$ ,

$$z^p = z^{p-1}(a + bx^n),$$

hence

$$\begin{aligned} \int x^{m-1} z^p dx &= \int x^{m-1} z^{p-1}(a + bx^n) dx = a \int x^{m-1} z^{p-1} dx \\ &\quad + b \int x^{m+n-1} z^{p-1} dx; \end{aligned}$$

therefore, by [1],

$$\frac{x^m z^p}{m} - \frac{bnp}{m} \int x^{m+n-1} z^{p-1} dx = a \int x^{m-1} z^{p-1} dx + b \int x^{m+n-1} z^{p-1} dx,$$

$$\int x^{m-1} z^{p-1} dx = \frac{x^m z^p}{am} - \frac{b(m+np)}{am} \int x^{m+n-1} z^{p-1} dx.$$

Change  $p$  into  $p + 1$ .

$$\int x^{m-1} z^p dx = \frac{x^m z^{p+1}}{am} - \frac{b(m+np+n)}{am} \int x^{m+n-1} z^p dx. \quad [3]$$

Change  $m$  into  $m - n$ , and transpose.

$$\int x^{m-1} z^p dx = \frac{x^{m-n} z^{p+1}}{b(m+np)} - \frac{a(m-n)}{b(m+np)} \int x^{m-n-1} z^p dx. \quad [4]$$

We have seen that

$$\int x^{m-1} z^p dx = a \int x^{m-1} z^{p-1} dx + b \int x^{m+n-1} z^{p-1} dx,$$

and, from [1],

$$b \int x^{m+n-1} z^{p-1} dx = \frac{x^m z^p}{np} - \frac{m}{np} \int x^{m-1} z^p dx;$$

hence

$$\int x^{m-1} z^p dx = a \int x^{m-1} z^{p-1} dx + \frac{x^m z^p}{np} - \frac{m}{np} \int x^{m-1} z^p dx,$$

$$\int x^{m-1} z^p dx = \frac{x^m z^p}{m + np} + \frac{anp}{m + np} \int x^{m-1} z^{p-1} dx. \quad [5]$$

Change  $p$  into  $p + 1$ , and transpose.

$$\int x^{m-1} z^p dx = -\frac{x^m z^{p+1}}{an(p+1)} + \frac{m + np + n}{an(p+1)} \int x^{m-1} z^{p+1} dx. \quad [6]$$

Formula [3] enables us to raise, and formula [4] to lower, the exponent of  $x$  by  $n$  without affecting the exponent of  $z$ ; while formula [5] enables us to lower, and formula [6] to raise, the exponent of  $z$  by unity without affecting the exponent of  $x$ .

Formulas [1] and [3] cannot be used when  $m = 0$ ;

formulas [2] and [6] cannot be used when  $p = -1$ ;

formulas [4] and [5] cannot be used when  $m = -np$ ;

for in all these cases infinite values will be brought into the second member of the formula.

65. If  $n = 1$ ,  $z = a + bx$ ,

and our last four reduction formulas become

$$\int x^{m-1} z^p dx = \frac{x^m z^{p+1}}{am} - \frac{b(m+p+1)}{am} \int x^m z^p dx. \quad [3]$$

$$\int x^{m-1} z^p dx = \frac{x^{m-1} z^{p+1}}{b(m+p)} - \frac{a(m-1)}{b(m+p)} \int x^{m-2} z^p dx. \quad [4]$$

$$\int x^{m-1} z^p dx = \frac{x^m z^p}{m+p} + \frac{ap}{m+p} \int x^{m-1} z^{p-1} dx. \quad [5]$$

$$\int x^{m-1} z^p dx = -\frac{x^m z^{p+1}}{a(p+1)} + \frac{m+p+1}{a(p+1)} \int x^{m-1} z^{p+1} dx. \quad [6]$$

If  $m$  and  $p$  are integers, and  $m > 0$  and  $p > 0$ , a repeated use of [5] will reduce  $p$  to zero, and we shall have to find merely the  $\int x^{m-1} dx$ .

If  $m < 0$  and  $p > 0$ , [3] will enable us to raise  $m$  to 0, and then [5] will enable us to lower  $p$  to 0, and we shall need only  $\int \frac{dx}{x}$ .

If  $m > 0$  and  $p < 0$ , [6] will raise  $p$  to  $-1$ , and [4] will then lower  $m$  to 1, and we shall need  $\int \frac{dx}{z}$ .

If  $m < 0$  and  $p < 0$ , [6] will raise  $p$  to  $-1$ , and [3] will raise  $m$  to 0, and we shall need  $\int \frac{dx}{xz}$ .

$$\int x^{m-1} dx = \frac{x^m}{m},$$

$$\int \frac{dx}{x} = \log x,$$

$$\int \frac{dx}{z} = \int \frac{dx}{a+bx} = \frac{1}{b} \log(a+bx),$$

$$\int \frac{dx}{xz} = \int \frac{dx}{x(a+bx)} = -\frac{1}{a} \log \frac{a+bx}{x}.$$

Hence, when  $n = 1$ , and  $m$  and  $p$  are integers, our reduction formulas always lead to the desired result.

#### EXAMPLES.

$$(1) \int \frac{dx}{x^2(a+bx)} = -\frac{b^4}{a^5} \log \frac{a+bx}{x} + \frac{b^3}{a^4 x} - \frac{b^2}{2a^3 x^2} + \frac{b}{3a^2 x^3} - \frac{1}{4ax^4}.$$

(2) Consider the case where  $n = 2$ , rewriting the reduction formulas to suit the case, and giving an exhaustive investigation.

$$(3) \int \frac{x^2 dx}{(a+bx^2)^3} = -\frac{x}{4b(a+bx^2)^2} + \frac{x}{8ab(a+bx^2)} + \frac{1}{8(ab)^{\frac{1}{2}}} \tan^{-1} x \sqrt{\frac{b}{a}}.$$

## CHAPTER VI.

## IRRATIONAL FORMS.

66. We have seen that algebraic polynomials and rational fractions can always be integrated. When we come to irrational expressions, however, very few forms are integrable, and most of these have to be rationalized by ingenious substitutions.

If an algebraic function is irrational because of the presence of an expression of the first degree under the radical sign, it can be easily made rational.

Let  $f(x, \sqrt[n]{a+bx})$  be the function in question.

Let  $z = \sqrt[n]{a+bx}$ ;

then  $z^n = a + bx$ .

$$nz^{n-1}dz = bdx,$$

$$dx = \frac{nz^{n-1}dz}{b};$$

$$x = \frac{z^n - a}{b}.$$

Hence  $\int f(x, \sqrt[n]{a+bx}) dx = \frac{n}{b} \int f\left(\frac{z^n - a}{b}, z\right) z^{n-1} dz,$

which is rational and can be treated by the methods of Chapter IV.

## EXAMPLES.

$$(1) \int \frac{\sqrt{x+1}}{\sqrt{x-1}} dx = x + 4\sqrt{x+4} \log(\sqrt{x-1}).$$

$$(2) \int \sqrt[n]{(ax+b)^m} dx = \frac{n\sqrt[n]{(ax+b)^{m+n}}}{a(m+n)}.$$

$$(3) \int [x\sqrt[n]{x+a} + \sqrt{x+a}] dx \\ = \frac{n\sqrt[n]{(x+a)^{2n+1}}}{2n+1} - \frac{na\sqrt[n]{(x+a)^{n+1}}}{n+1} + \frac{2}{3}\sqrt{x+a}^3.$$

67. A case not unlike the last is  $\int f(x, \sqrt[n]{c + \sqrt[n]{a + bx}}) dx$ .

Let

$$z = \sqrt[n]{c + \sqrt[n]{a + bx}};$$

$$z^n = c + \sqrt[n]{a + bx},$$

$$(z^n - c)^n = a + bx,$$

$$x = \frac{(z^n - c)^n - a}{b},$$

$$dx = \frac{mn(z^n - c)^{n-1} z^{n-1} dz}{b}.$$

Hence

$$\int f(x, \sqrt[n]{c + \sqrt[n]{a + bx}}) dx$$

$$= \frac{mn}{b} \int f\left[\frac{(z^n - c)^n - a}{b}, z\right] (z^n - c)^{n-1} z^{n-1} dz.$$

EXAMPLES.

- (1) Find  $\int \frac{xdx}{\sqrt{c + \sqrt{a + bx}}}$ .
- (2) Find  $\int \frac{dx}{\sqrt[3]{1 + \sqrt{1 - x}}}$ .

68. If the expression under the radical is of a higher degree than the first the function cannot in general be rationalized. The only important exceptional case is where the function to be integrated is irrational by reason of containing the square root of a quantity of the second degree.

Required  $\int f(x, \sqrt{a + bx + cx^2}) dx$ .

*First Method.* Let  $c$  be positive; take out  $\sqrt{c}$  as a factor, and the radical may be written  $\sqrt{A + Bx + x^2}$ .

Let

$$\sqrt{A + Bx + x^2} = x + z,$$

$$A + Bx + x^2 = x^2 + 2xz + z^2,$$

$$x = \frac{z^2 - A}{B - 2z},$$



$$dx = -\frac{2(z^2 - Bz + A)dz}{(B - 2z)^2},$$

$$\sqrt{A + Bx + x^2} = x + z = -\frac{z^2 - Bz + A}{B - 2z},$$

and the substitution of these values will render the given function rational.

*Second Method.* Let  $c$  be positive; take out  $\sqrt{c}$  as a factor, and, as before, the radical may be written  $\sqrt{A + Bx + x^2}$ .

$$\text{Let } \sqrt{A + Bx + x^2} = \sqrt{A + xz};$$

$$A + Bx + x^2 = A + 2\sqrt{A} \cdot xz + x^2z^2,$$

$$x = \frac{2\sqrt{A} \cdot z - B}{1 - z^2},$$

$$dx = \frac{2(\sqrt{A} \cdot z^2 - Bz + \sqrt{A})dz}{(1 - z^2)^2},$$

$$\sqrt{A + Bx + x^2} = \sqrt{A + xz} = \frac{\sqrt{A} \cdot z^2 - Bz + \sqrt{A}}{1 - z^2},$$

and the substitution of these values will render the given function rational.

If  $c$  is negative the radical can be reduced to the form  $\sqrt{A + Bx - x^2}$ , and the method just given will present no difficulty.

*Third Method.* Let  $c$  be positive; the radical will reduce to  $\sqrt{A + Bx + x^2}$ . Resolve the quantity under the radical into the product of two binomial factors  $(x - a)(x - \beta)$ ,  $a$  and  $\beta$  being the roots of the equation  $A + Bx + x^2 = 0$ .

$$\text{Let } \sqrt{(x - a)(x - \beta)} = (x - a)z;$$

$$(x - a)(x - \beta) = (x - a)^2z^2,$$

$$x = \frac{\beta - az^2}{1 - z^2},$$

$$dx = \frac{2z(\beta - a)dz}{(1 - z^2)^2},$$

$$\sqrt{(x - a)(x - \beta)} = (x - a)z = \frac{(\beta - a)z}{1 - z^2},$$

and the substitution of these values will make the given function rational.

If  $c$  is negative the radical will reduce to  $\sqrt{A+Bx-x^2}$ , and may be written  $\sqrt{(a-x)(x-\beta)}$  where  $a$  and  $\beta$  are the roots of  $x^2-Bx-A=0$ , and the method just explained will apply.

In general, that one of the three methods is preferable which will avoid introducing imaginary constants; the first, if  $c > 0$ ; the second, if  $c < 0$  and  $\frac{a}{-c} > 0$ ; the third, if  $c < 0$  and  $\frac{a}{-c} < 0$ . If the roots  $a$  and  $\beta$  are imaginary, and  $A = \frac{a}{-c}$  is negative, it will be impossible to avoid imaginaries, for in that case  $A+Bx-x^2$  will be negative for all real values of  $x$ .

69. Let us compare the working of the three methods just given by applying them in turn to the example  $\int \frac{dx}{\sqrt{2+3x+x^2}}$ .

1st. Let  $\sqrt{2+3x+x^2} = x+z$ ;

$$\int \frac{dx}{\sqrt{2+3x+x^2}} = \int \frac{2(x^2-3z+2)dz}{(3-2z)^2} \cdot \frac{3-2z}{x^2-3z+2} = \int \frac{2dz}{3-2z}$$

$$= -\log(3-2z),$$

$$\int \frac{dx}{\sqrt{2+3x+x^2}} = -\log(3+2x-2\sqrt{2+3x+x^2})$$

$$= \log \frac{1}{3+2x-2\sqrt{2+3x+x^2}}$$

$$= \log \frac{3+2x+2\sqrt{2+3x+x^2}}{9+12x+4x^2-8-12x-4x^2}$$

$$= \log [3+2x+2\sqrt{2+3x+x^2}]. \quad (1)$$

2d. Let  $\sqrt{2+3x+x^2} = \sqrt{2}+xz$ ;

$$\int \frac{dx}{\sqrt{2+3x+x^2}} = 2 \int \frac{(\sqrt{2} \cdot z^2 - 3z + \sqrt{2})dz}{(1-z^2)^2} \cdot \frac{1-z^2}{\sqrt{2} \cdot z^2 - 3z + \sqrt{2}}$$

$$= 2 \int \frac{dz}{1-z^2} = \log \frac{1+z}{1-z}. \quad (\text{Art. 53})$$

$$\begin{aligned} \int \frac{dx}{\sqrt{2+3x+x^2}} &= \log \frac{x - \sqrt{2} + \sqrt{2+3x+x^2}}{x + \sqrt{2} - \sqrt{2+3x+x^2}} \\ &= \log \frac{x^2 + 2x\sqrt{2+3x+x^2} + 2 + 3x + x^2 - 2}{x^2 + 2\sqrt{2} \cdot x + 2 - 2 - 3x - x^2} \\ &= \log \frac{3 + 2x + 2\sqrt{2+3x+x^2}}{2\sqrt{2}-3} \\ &= \log(3 + 2x + 2\sqrt{2+3x+x^2}) - \log(2\sqrt{2}-3), \end{aligned}$$

or, dropping the constant  $\log(2\sqrt{2}-3)$ ,

$$\int \frac{dx}{\sqrt{2+3x+x^2}} = \log(3 + 2x + 2\sqrt{2+3x+x^2}). \quad (2)$$

3d. Let  $\sqrt{2+3x+x^2} = \sqrt{(x+1)(x+2)} = (x+1)z$ ;

$$\int \frac{dx}{\sqrt{2+3x+x^2}} = 2 \int \frac{-zdz}{(1-z^2)^2} \frac{1-z^2}{-z} = 2 \int \frac{dz}{1-z^2} = \log \frac{1+z}{1-z}$$

$$\begin{aligned} \int \frac{dx}{\sqrt{2+3x+x^2}} &= \log \frac{1 + \sqrt{\frac{x+2}{x+1}}}{1 - \sqrt{\frac{x+2}{x+1}}} = \log \frac{\sqrt{x+1} + \sqrt{x+2}}{\sqrt{x+1} - \sqrt{x+2}} \\ &= \log \frac{x+1 + 2\sqrt{2+3x+x^2} + x+2}{x+1 - x-2} \\ &= \log(3 + 2x + 2\sqrt{2+3x+x^2}) + \log(-1), \end{aligned}$$

or, dropping the imaginary constant  $\log(-1)$ ,

$$\int \frac{dx}{\sqrt{2+3x+x^2}} = \log(3 + 2x + 2\sqrt{2+3x+x^2}). \quad (3)$$

#### EXAMPLES.

$$(1) \int \frac{dx}{(2+3x)\sqrt{4-x^2}} = \frac{1}{4\sqrt{2}} \log \frac{\sqrt{4+2x} - \sqrt{2-x}}{\sqrt{4+2x} + \sqrt{2-x}}$$

$$(2) \int \frac{dx}{\sqrt{x^2+x}} = \log\left(\frac{1}{2} + x + \sqrt{x^2+x}\right).$$

$$(3) \int \frac{dx}{\sqrt{a+bx+cx^2}} = \frac{1}{\sqrt{c}} \log\left(\frac{b}{2\sqrt{c}} + x\sqrt{c} + \sqrt{a+bx+cx^2}\right).$$

70. If the function is irrational through the presence, under the radical sign, of a fraction whose numerator and denominator are of the first degree, it can always be rationalized.

Required  $\int f(x, \sqrt[n]{\frac{ax+b}{lx+m}}) dx.$

Let 
$$z = \sqrt[n]{\frac{ax+b}{lx+m}},$$

$$z^n = \frac{ax+b}{lx+m},$$

$$x = \frac{b-mz^n}{lz^n-a},$$

$$dx = \frac{n(am-bl)z^{n-1} dz}{(lz^n-a)^2},$$

and the substitution of these values will make the given function rational.

EXAMPLE.

$$\int \frac{dx}{(1+x)^2} \sqrt[3]{\frac{1-x}{1+x}} = -\frac{8}{3} \sqrt[3]{\frac{(1-x)^4}{(1+x)}}.$$

71. If the function to be integrated is of the form  $x^{m-1}(a+bx^n)^p$ ,  $m$ ,  $n$ , and  $p$  being any numbers positive or negative, and one at least of them being fractional, the reduction formulas of Art. 64 will often lead to the desired integral.

EXAMPLES.

- (1)  $\int \frac{x^4 dx}{(1-x^2)^{\frac{1}{2}}} = \frac{8}{3} \sin^{-1} x - \frac{x \sqrt{1-x^2}}{8} (3+2x^2).$
- (2)  $\int \frac{dx}{x^3 \sqrt{1-x^2}} = \frac{1}{2} \log \frac{1-\sqrt{1-x^2}}{x} - \frac{\sqrt{1-x^2}}{2x^2}.$
- (3)  $\int \frac{x^3 dx}{(2ax-x^2)^{\frac{1}{2}}} = -(2ax-x^2)^{\frac{1}{2}} \left( \frac{x}{2} + \frac{3a}{2} \right) + 3a^2 \sin^{-1} \sqrt{\frac{x}{2a}}.$
- (4)  $\int \frac{x^3 dx}{(a^2+x^2)^{\frac{3}{2}}} = -\frac{(2a^2+3x^2)}{3(a^2+x^2)^{\frac{3}{2}}}.$

72. We have said that when an irrational function contains a quantity of a higher degree than the second, under the square-root sign, it cannot *ordinarily* be integrated. It would be more correct to say that its integral cannot ordinarily be finitely expressed in terms of the functions with which we are familiar.

The integrals of a large class of such irrational expressions have been specially studied under the name of Elliptic Integrals. They have peculiar properties, and can be expressed in terms of ordinary functions only by the aid of infinite series.

CHAPTER VII.

TRANSCENDENTAL FUNCTIONS.

73. In dealing with the integration of transcendental functions the method of *integration by parts* is generally the most effective.

For example. Required  $\int x(\log x)^2 dx$ .

Let

$$u = (\log x)^2,$$

$$dv = x dx;$$

$$du = \frac{2 \log x dx}{x},$$

$$v = \frac{x^2}{2},$$

$$\int x(\log x)^2 = \frac{x^2(\log x)^2}{2} - \int x \log x dx = \frac{x^2}{2} [(\log x)^2 - \log x + \frac{1}{2}].$$

Again. Required  $\int e^x \sin x dx$ .

$$u = \sin x,$$

$$dv = e^x dx;$$

$$du = \cos x dx,$$

$$v = e^x,$$

$$\int e^x \sin x dx = e^x \sin x - \int e^x \cos x dx,$$

$$\int e^x \cos x dx = e^x \cos x + \int e^x \sin x dx;$$

whence 
$$\int e^x \sin x dx = \frac{e^x(\sin x - \cos x)}{2},$$

and 
$$\int e^x \cos x dx = \frac{e^x(\sin x + \cos x)}{2}.$$

## EXAMPLES.

$$(1) \int x^m (\log x)^3 dx = \frac{x^{m+1}}{m+1} \left[ (\log x)^3 - \frac{3(\log x)^2}{m+1} + \frac{6 \log x}{(m+1)^2} - \frac{6}{(m+1)^3} \right].$$

$$(2) \int \frac{\log x \cdot dx}{(1-x)^2} = \frac{x \log x}{1-x} + \log(1-x).$$

$$(3) \int e^{ax} \sqrt{1-e^{2ax}} \cdot dx = \frac{1}{2a} \left[ e^{ax} \sqrt{1-e^{2ax}} + \sin^{-1} e^{ax} \right].$$

74. The method of integration by parts gives us important reduction formulas for transcendental functions. Let us consider  $\int \sin^n x \cdot dx$ .

$$u = \sin^{n-1} x,$$

$$dv = \sin x \cdot dx;$$

$$du = (n-1) \sin^{n-2} x \cos x \cdot dx,$$

$$v = -\cos x;$$

$$\begin{aligned} \int \sin^n x \cdot dx &= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \cos^2 x \cdot dx \\ &= -\sin^{n-1} x \cos x + (n-1) \int (\sin^{n-2} x - \sin^n x) dx; \end{aligned}$$

$$\int \sin^n x \cdot dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x \cdot dx. \quad [1]$$

Transposing, and changing  $n$  into  $n+2$ , we get

$$\int \sin^n x \cdot dx = \frac{1}{n+1} \sin^{n+1} x \cos x + \frac{n+2}{n+1} \int \sin^{n+2} x \cdot dx. \quad [2]$$

In like manner we get

$$\int \cos^n x \cdot dx = \frac{1}{n} \sin x \cos^{n-1} x + \frac{n-1}{n} \int \cos^{n-2} x \cdot dx, \quad [3]$$

$$\int \cos^n x \cdot dx = -\frac{1}{n+1} \sin x \cos^{n+1} x + \frac{n+2}{n+1} \int \cos^{n+2} x \cdot dx. \quad [4]$$

If  $n$  is a positive integer, formulas [1] and [3] will enable us to reduce the exponent of the sine or cosine to one or to zero,

and then we can integrate by inspection. If  $n$  is a negative integer, formulas [2] and [4] will enable us to raise the exponent to zero or to minus one. In the latter case we shall need  $\int \frac{dx}{\cos x}$ , or  $\int \frac{dx}{\sin x}$ , which have been found in Art. 54 (c).

EXAMPLES.

$$(1) \int \sin^4 x \, dx = -\frac{\sin x \cos x}{4} \left( \sin^2 x + \frac{3}{2} \right) + \frac{3}{8} x.$$

$$(2) \int \cos^4 x \, dx = \frac{\sin x \cos^3 x}{6} \left( \cos^2 x + \frac{5}{4} \right) + \frac{5}{16} (\sin x \cos x + x).$$

$$(3) \int \frac{dx}{\sin^3 x} = -\frac{\cos x}{2 \sin^2 x} + \frac{1}{2} \log \tan \frac{x}{2}.$$

(4) Obtain the formulas

$$\int \sinh^n x \, dx = \frac{1}{n} \sinh^{n-1} x \cosh x - \frac{n-1}{n} \int \sinh^{n-2} x \, dx.$$

$$\int \sinh^n x \, dx = \frac{1}{n+1} \sinh^{n+1} x \cosh x - \frac{n+2}{n+1} \int \sinh^{n+2} x \, dx.$$

$$\int \cosh^n x \, dx = \frac{1}{n} \sinh x \cosh^{n-1} x + \frac{n-1}{n} \int \cosh^{n-2} x \, dx.$$

$$\int \cosh^n x \, dx = -\frac{1}{n+1} \sinh x \cosh^{n+1} x + \frac{n+2}{n+1} \int \cosh^{n+2} x \, dx.$$

$$(5) \int \frac{dx}{\sinh^3 x} = -\frac{1}{2} \frac{\cosh x}{\sinh^2 x} - \frac{1}{4} \log \frac{\cosh x - 1}{\cosh x + 1}.$$

75. The  $(\sin^{-1} x)^n dx$  can be integrated by the aid of a reduction formula.

Let  $z = \sin^{-1} x$ ;

then  $x = \sin z$ ,

$$dx = \cos z \, dz,$$

and  $\int (\sin^{-1} x)^n dx = \int z^n \cos z \, dz.$



Let

$$\begin{aligned} u &= z^n, \\ dv &= \cos z \cdot dz; \\ du &= nz^{n-1} dz, \\ v &= \sin z; \end{aligned}$$

$$\int z^n \cos z \cdot dz = z^n \sin z - n \int z^{n-1} \sin z \cdot dz.$$

$\int z^{n-1} \sin z \cdot dz$  can be reduced in the same way, and is equal to  $-z^{n-1} \cos z + (n-1) \int z^{n-2} \cos z \cdot dz$ ;

hence

$$\int z^n \cos z \cdot dz = z^n \sin z + nz^{n-1} \cos z - n(n-1) \int z^{n-2} \cos z \cdot dz, \quad [1]$$

or

$$\begin{aligned} \int (\sin^{-1} x)^n dx &= x(\sin^{-1} x)^n + n\sqrt{1-x^2}(\sin^{-1} x)^{n-1} \\ &\quad - n(n-1) \int (\sin^{-1} x)^{n-2} dx. \end{aligned} \quad [2]$$

If  $n$  is a positive integer, this will enable us to make our required integral depend upon  $\int dx$  or  $\int \sin^{-1} x \cdot dx$ , the latter of which forms has been found in (I. Art. 81).

#### EXAMPLES.

(1) Obtain a formula for  $\int (\text{vers}^{-1} x)^n dx$ .

(2)  $\int (\sin^{-1} x)^4 dx = x[(\sin^{-1} x)^4 - 4 \cdot 3 \cdot (\sin^{-1} x)^2 + 4 \cdot 3 \cdot 2 \cdot 1]$   
 $+ 4 \sqrt{1-x^2} \sin^{-1} x [(\sin^{-1} x)^2 - 3 \cdot 2].$

76. Integration by substitution is sometimes a valuable method in dealing with transcendental forms, and in the case of the trigonometric functions often enables us to reduce the given form to an algebraic one. Let it be required to find  $\int (f \sin x) \cos x \cdot dx$ .

Let

$$\begin{aligned} z &= \sin x, \\ dz &= \cos x \cdot dx; \\ \int (f \sin x) \cos x \cdot dx &= \int fz \cdot dz. \end{aligned}$$

In the same way we see that

$$\int (f \cos x) \sin x \, dx = - \int f z \, dz \quad \text{if } z = \cos x,$$

and

$$\int [f(\sin x, \cos x)] \cos x \, dx = \int [f(z, \sqrt{1-z^2})] dz \quad \text{if } z = \sin x,$$

$$\int [f(\cos x, \sin x)] \sin x \, dx = - \int [f(z, \sqrt{1-z^2})] dz \quad \text{if } z = \cos x,$$

or, more generally,

$$\int f(\sin x, \cos x) \, dx = \int f(z, \sqrt{1-z^2}) \frac{dz}{\sqrt{1-z^2}} \quad \text{if } z = \sin x,$$

$$\int f(\cos x, \sin x) \, dx = - \int f(z, \sqrt{1-z^2}) \frac{dz}{\sqrt{1-z^2}} \quad \text{if } z = \cos x.$$

Since any trigonometric function of  $x$  may be expressed in terms of  $\sin x$  and  $\cos x$ , the formulas just given enable us to make the integration of any trigonometric function depend on the integration of an algebraic function, which, however, is frequently complicated by the presence of the radical  $\sqrt{1-z^2}$ .

77. A better substitution than that of the last article, when the form to be treated does not contain  $\sin x$  or  $\cos x$  as a factor, is  $z = \tan \frac{x}{2}$ .

This gives us 
$$dx = \frac{2 \, dz}{1 + z^2},$$

$$\sin x = \frac{2z}{1 + z^2},$$

$$\cos x = \frac{1 - z^2}{1 + z^2};$$

whence 
$$\int f(\sin x, \cos x) \, dx = 2 \int f\left(\frac{2z}{1+z^2}, \frac{1-z^2}{1+z^2}\right) \frac{dz}{1+z^2}. \quad [1]$$

As an example, let us find 
$$\int \frac{dx}{a + b \cos x}.$$

Here we have

$$\begin{aligned}\int \frac{dx}{a+b \cos x} &= 2 \int \frac{dz}{(1+z^2) \left[ a+b \frac{1-z^2}{1+z^2} \right]} = 2 \int \frac{dz}{a+b+(a-b)z^2} \\ &= \frac{2}{a-b} \int \frac{dz}{\frac{a+b}{a-b} + z^2} = \frac{2}{\sqrt{a^2-b^2}} \tan^{-1} \left( \sqrt{\frac{a-b}{a+b}} \cdot z \right) \\ &\qquad\qquad\qquad \text{by I. Art. 77, Ex. 1.}\end{aligned}$$

Hence  $\int \frac{dx}{a+b \cos x} = \frac{2}{\sqrt{a^2-b^2}} \tan^{-1} \left( \sqrt{\frac{a-b}{a+b}} \cdot \tan \frac{x}{2} \right)$ , if  $a > b$ .

78.  $\int \sin^m x \cos^n x \cdot dx$  can be readily found by the method of Art. 76 if  $m$  and  $n$  are positive integers, and if either of them is odd. Let  $n$  be odd, then

$$\cos^n x = \cos^{n-1} x \cos x = (1 - \sin^2 x)^{\frac{n-1}{2}} \cos x,$$

$$\int \sin^m x \cos^n x \cdot dx = \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} \cos x \cdot dx.$$

Let

$$z = \sin x,$$

$$dz = \cos x \cdot dx,$$

$$\int \sin^m x \cos^n x \cdot dx = \int z^m (1 - z^2)^{\frac{n-1}{2}} dz,$$

which can be expanded into an algebraic polynomial and integrated directly.

If  $m$  and  $n$  are positive integers, and are both even,

$$\int \sin^m x \cos^n x \cdot dx = \int \sin^m x (1 - \sin^2 x)^{\frac{n}{2}} dx.$$

$\sin^m x (1 - \sin^2 x)^{\frac{n}{2}}$  can be expanded and thus integrated by Art. 74 [1].

If  $m$  or  $n$  is negative, and odd, we can write

$$\cos^n x = \cos^{n-1} x \cos x, \quad \text{or} \quad \sin^m x = \sin^{m-1} x \sin x,$$

and reduce the function to be integrated to a rational fraction by the substitution of

$$z = \cos x, \quad \text{or} \quad z = \sin x.$$

$\int \sin^m x \cos^n x \cdot dx$  can also be treated by the aid of reduction formulas easily obtained.

79.  $\int \tan^n x dx$  and  $\int \frac{dx}{\tan^n x}$  can be handled by the methods of Art. 78, but they can be simplified greatly by a reduction formula.

We have

$$\begin{aligned} \int \tan^n x dx &= \int \tan^{n-2} x \tan^2 x dx = \int \tan^{n-2} x (\sec^2 x - 1) dx \\ &= \int \tan^{n-2} x d(\tan x) - \int \tan^{n-2} x dx, \end{aligned}$$

whence  $\int \tan^n x dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x dx;$  [1]

and  $\int \frac{dx}{\tan^n x} = \int \frac{\sec^2 x - \tan^2 x}{\tan^n x} dx = \int \frac{d(\tan x)}{\tan^n x} - \int \frac{dx}{\tan^{n-2} x},$

whence  $\int \frac{dx}{\tan^n x} = -\frac{1}{(n-1)\tan^{n-1} x} - \int \frac{dx}{\tan^{n-2} x}.$  [2]

EXAMPLES.

(1)  $\int \sin^3 x \cos^7 x dx = \frac{\cos^{10} x}{10} - \frac{\cos^8 x}{8}.$

(2)  $\int \cos^3 x \sqrt{\sin x} dx = \frac{2 \sin^{\frac{3}{2}} x}{3} - \frac{2 \sin^{\frac{1}{2}} x}{7}.$

(3)  $\int \frac{\sin^3 x dx}{\sqrt{\cos x}} = \frac{2 \cos^{\frac{3}{2}} x}{5} - 2 \cos^{\frac{1}{2}} x.$

(4)  $\int \cos^2 x \sin^4 x dx = \frac{\sin x \cos x}{2} \left( \frac{\sin^4 x}{3} - \frac{\sin^2 x}{12} - \frac{1}{8} \right) + \frac{x}{16}.$

(5)  $\int \frac{dx}{\sin x \cos^2 x} = \sec x + \log \tan \frac{x}{2}.$

(6)  $\int \frac{dx}{\sin^3 x \cos^2 x} = \sec x - \frac{\cos x}{2 \sin^2 x} + \frac{3}{2} \log \tan \frac{x}{2}.$

(7)  $\int \frac{dx}{\tan^3 x} = -\frac{1}{4 \tan^4 x} + \frac{1}{2 \tan^2 x} + \log \sin x.$

$$(8) \int \frac{dx}{a + b \cos x} = \frac{1}{\sqrt{b^2 - a^2}} \log \frac{\sqrt{b+a} + \sqrt{b-a} \cdot \tan \frac{x}{2}}{\sqrt{b+a} - \sqrt{b-a} \cdot \tan \frac{x}{2}}.$$

$$(9) \int \frac{dx}{5 + 4 \sin x} = \frac{2}{3} \tan^{-1} \left( \frac{4 + 5 \tan \frac{x}{2}}{3} \right).$$

$$(10) \int \frac{dx}{3 \sin x + \sin 2x} = \frac{1}{5} \log \sin \frac{x}{2} - \log \cos \frac{x}{2} + \frac{2}{5} \log(3 + 2 \cos x).$$

$$(11) \int \frac{\cos x dx}{(5 + 4 \cos x)^2} = \frac{5}{9} \frac{\sin x}{5 + 4 \cos x} - \frac{8}{27} \tan^{-1} \left( \frac{1}{3} \tan \frac{x}{2} \right).$$

$$(12) \int \frac{dx}{a + b \sin x + c \cos x} = \frac{2}{\sqrt{a^2 - b^2 - c^2}} \tan^{-1} \left[ \frac{(a-c) \tan \frac{x}{2} + b}{\sqrt{a^2 - b^2 - c^2}} \right].$$

CHAPTER VIII.

DEFINITE INTEGRALS.

80. In I. Art. 183, a definite integral has been defined as the limit of a sum of infinitesimal terms, and has been proved equal to the difference between two values of an ordinary integral.


We are now ready to put our definition into more precise, and at the same time more general, form.

If  $f x$  is finite, continuous, and single-valued between the values  $x = a$  and  $x = b$ , and we form the sum

$$(x_1 - a)fa + (x_2 - x_1)fx_1 + (x_3 - x_2)fx_2 + \dots + (x_{n-1} - x_{n-2})fx_{n-2} + (b - x_{n-1})fx_{n-1},$$

where  $x_1, x_2, x_3 \dots x_{n-1}$  are  $n - 1$  successive values of  $x$  lying between  $a$  and  $b$ , the limit approached by this sum as  $n$  is indefinitely increased, while at the same time each of the increments  $(x_1 - a), (x_2 - x_1)$ , etc., is made to approach zero, is the definite integral of  $f x$  from  $a$  to  $b$ , and will be denoted by  $\int_a^b f x . dx$ .

If we construct the curve  $y = f x$  in rectangular co-ordinates, this definition clearly requires us to break up the projection on the axis of  $X$  of the portion of the curve between the points  $A$  and  $B$  into  $n$  intervals, to multiply each interval by the ordinate at its beginning, and to take the limit of the sum of these products as each interval is indefinitely decreased; that is, the limit of the sum of the small rectangles in the figure, and this is easily proved to be the area  $ABA_1B_1$ .



Now the area  $ABA_1B_1$ , found by the method of I. Chap. V., is

$$\left[ \int f x . dx \right]_{z=b} - \left[ \int f x . dx \right]_{z=a} .$$

Therefore  $\int_a^b fx \cdot dx = \left[ \int fx \cdot dx \right]_{x=b} - \left[ \int fx \cdot dx \right]_{x=a}$ . [1]

That is,  $\int_a^b fx \cdot dx$  is the increment produced in  $\int fx \cdot dx$  by changing  $x$  from  $a$  to  $b$ .

It is to be noted that the successive increments  $(x_1 - a)$ ,  $(x_2 - x_1)$ ,  $(x_3 - x_2)$ , etc., that is, the successive values of  $dx$ , are not necessarily equal; and also, that if we multiply each interval, not by the ordinate at its beginning, but by an ordinate erected at any point of its length, the limit of our sum will be unaltered. (v. I. Arts. 161, 149.)

81. It is instructive to find a few definite integrals by actually performing the summation suggested in the definition (Art. 80), and then finding the limit of the sum.

$$(a) \quad \int_a^b x \cdot dx.$$

Let us divide the interval from  $a$  to  $b$  into  $n$  equal parts, and call each of them  $dx$ .

$$\text{Then} \quad ndx = b - a.$$

Our sum is

$$\begin{aligned} S &= adx + (a+dx) dx + (a+2dx) dx + \dots + (a+(n-1) dx) dx \\ &= nadx + (1+2+3+\dots+(n-1)) dx^2 \\ &= a(b-a) + \frac{n(n-1)}{2} dx^2, \end{aligned}$$

since  $ndx = b - a$ , and the sum of the arithmetical progression

$$1+2+3+\dots+(n-1) = \frac{n(n-1)}{2}.$$

$$\frac{n(n-1)}{2} dx^2 = \frac{1}{2} (n^2 dx^2 - n dx^2) = \frac{(b-a)^2}{2} - \frac{(b-a)dx}{2}.$$

$$\text{Hence} \quad S = \frac{b^2 - a^2}{2} - \frac{(b-a)}{2} dx.$$

As we increase  $n$  indefinitely,  $dx$  approaches zero, and

$$\int_a^b x \cdot dx = \lim_{dx \rightarrow 0} \left[ \frac{b^2 - a^2}{2} - \frac{(b-a) dx}{2} \right] = \frac{b^2}{2} - \frac{a^2}{2}.$$

(b) 
$$\int_a^b e^x dx.$$

Let 
$$dx = \frac{b-a}{n}.$$

$$S = e^a dx + e^{a+dx} dx + e^{a+2dx} dx + \dots + e^{a+(n-1)dx} dx \\ = e^a dx [1 + e^{dx} + e^{2dx} + e^{3dx} + \dots + e^{(n-1)dx}];$$

but  $1 + e^{dx} + e^{2dx} + \dots + e^{(n-1)dx}$  is a geometrical progression, and its sum is

$$\frac{e^{n dx} - 1}{e^{dx} - 1} = \frac{e^{b-a} - 1}{e^{dx} - 1}.$$

Hence 
$$S = \frac{e^{b-a} - 1}{e^{dx} - 1} \cdot e^a dx = (e^b - e^a) \frac{dx}{e^{dx} - 1},$$

and 
$$\int_a^b e^x dx = (e^b - e^a) \lim_{dx \rightarrow 0} \left[ \frac{dx}{e^{dx} - 1} \right];$$

but as  $dx$  approaches zero,  $\frac{dx}{e^{dx} - 1}$  approaches the indeterminate form  $\frac{0}{0}$ ; but since the true value of

$$\left[ \frac{a}{e^a - 1} \right]_{a=0} = \left[ \frac{1}{e^a} \right]_{a=0} = 1,$$

$$\int_a^b e^x dx = e^b - e^a.$$

(c) 
$$\int_0^\pi \cos^3 x \cdot dx.$$

Let  $dx = \frac{\pi}{n}$ , and let  $n$  be an odd number.

Then

$$S = dx + \cos^3 dx \cdot dx + \cos^3 2 dx \cdot dx + \dots + \cos^3 (n-2) dx \cdot dx \\ + \cos^3 (n-1) dx \cdot dx \\ = dx + \cos^3 dx \cdot dx + \cos^3 2 dx \cdot dx + \dots + \cos^3 (\pi - 2 dx) \cdot dx \\ + \cos^3 (\pi - dx) \cdot dx \\ = dx + \cos^3 dx \cdot dx + \cos^3 2 dx \cdot dx + \dots - \cos^3 2 dx \cdot dx \\ - \cos^3 dx \cdot dx,$$



since  $\cos(\pi - \phi) = -\cos \phi$ .

Hence the terms cancel in pairs, and we have left

$$S = dx$$

and  $\int_0^{\pi} \cos^3 x \cdot dx = \lim_{dx \rightarrow 0} \left[ dx \right] = 0$ .

$$(d) \quad \int_0^{\frac{\pi}{2}} \sin^2 x \cdot dx.$$

Let  $dx = \frac{\pi}{2n}$ , and let  $n$  be an odd number.

$$\begin{aligned} S &= \sin^2 0 \cdot dx + \sin^2 dx \cdot dx + \sin^2 2 dx \cdot dx + \dots + \sin^2(n-2) dx \cdot dx \\ &\quad + \sin^2(n-1) dx \cdot dx \\ &= \sin^2 dx \cdot dx + \sin^2 2 dx \cdot dx + \dots + \sin^2\left(\frac{\pi}{2} - 2 dx\right) dx + \sin^2\left(\frac{\pi}{2} - dx\right) dx \\ &= \sin^2 dx \cdot dx + \sin^2 2 dx \cdot dx + \dots + \cos^2 2 dx \cdot dx + \cos^2 dx \cdot dx, \end{aligned}$$

since  $\sin\left(\frac{\pi}{2} - \phi\right) = \cos \phi$ .

Then  $S = dx + dx + dx \dots = \frac{n-1}{2} dx$ ,

since  $\sin^2 \phi + \cos^2 \phi = 1$ .

Therefore  $S = \frac{\pi}{4} - \frac{dx}{2}$ ,

and  $\int_0^{\frac{\pi}{2}} \sin^2 x \cdot dx = \frac{\pi}{4}$ .

$$(e) \quad \int_a^b \frac{dx}{x}$$

Here it is best to divide the interval between  $a$  and  $b$  into unequal parts.

Let the values  $x_1, x_2, x_3 \dots x_{n-1}$  be such as to form with  $a$  and  $b$  a geometrical progression.

For this purpose take  $q = \sqrt[n]{\frac{b}{a}}$ , so that  $aq^n = b$ .

Then the values in question are  $aq, aq^2, aq^3 \dots aq^{n-1}$ , and the intervals are  $a(q-1), aq(q-1), aq^2(q-1) \dots aq^{n-1}(q-1)$ , and the sum

$$S = \frac{a(q-1)}{a} + \frac{aq(q-1)}{aq} + \frac{aq^2(q-1)}{aq^2} + \dots + \frac{aq^{n-1}(q-1)}{aq^{n-1}} \\ = n(q-1).$$

To prove our division legitimate we have only to show that each of our intervals,  $a(q-1), aq(q-1) \dots aq^{n-1}(q-1)$ , approaches the limit zero as  $n$  increases indefinitely. Since

$$q^n = \frac{b}{a}$$

the limiting value of  $q$  as  $n$  increases must be 1, as otherwise  $\lim_{n \rightarrow \infty} q^n$  would not be finite.

$$\text{Therefore } \lim_{n \rightarrow \infty} [aq^n(q-1)] = \lim_{q \rightarrow 1} [aq^n(q-1)] = 0.$$

We have then

$$\int_a^b \frac{dx}{x} = \lim_{n \rightarrow \infty} [S] = \lim_{n \rightarrow \infty} [n(q-1)] = \lim_{q \rightarrow 1} [n(q-1)] \\ = \lim_{q \rightarrow 1} \left[ \frac{\log \frac{b}{a}}{\log q} (q-1) \right],$$

$$\text{since } n \log q = \log \frac{b}{a}.$$

$$\text{But } \lim_{q \rightarrow 1} \left[ \frac{\log \frac{b}{a}}{\log q} (q-1) \right] = \log \frac{b}{a} \lim_{q \rightarrow 1} \left[ \frac{q-1}{\log q} \right] = \log \frac{b}{a}.$$

$$\text{For } \left[ \frac{q-1}{\log q} \right]_{q \rightarrow 1} = \left[ \frac{1}{\frac{1}{q}} \right]_{q \rightarrow 1} = 1.$$

$$\text{Therefore } \int_a^b \frac{dx}{x} = \log b - \log a.$$

## EXAMPLES.

(1) Prove by the methods of this article that

$$\int_a^c a^x dx = \frac{a^c - a^a}{\log a}.$$

(2) By the aid of the trigonometric formulas

$$\begin{aligned} & \cos \theta + \cos 2\theta + \cos 3\theta + \dots + \cos (n-1)\theta \\ &= \frac{1}{2} \left[ \sin n\theta \operatorname{ctn} \frac{\theta}{2} - 1 - \cos n\theta \right], \end{aligned}$$

$$\begin{aligned} & \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin (n-1)\theta \\ &= \frac{1}{2} \left[ (1 - \cos n\theta) \operatorname{ctn} \frac{\theta}{2} - \sin n\theta \right], \end{aligned}$$

prove that  $\int_a^b \cos x \cdot dx = \sin b - \sin a,$

and  $\int_a^b \sin x \cdot dx = \cos a - \cos b.$

(3) Show that  $\int_0^{2\pi} \sin^5 x \cdot dx = 0,$

and that  $\int_0^{\pi} \cos^2 x \cdot dx = \frac{\pi}{2}.$

(4) Show that  $\int_a^b x^m dx = \frac{b^{m+1} - a^{m+1}}{m+1},$  using the method of Art. 81 (e).

82. When the indefinite integral can be found, the definite integral  $\int_a^b fx \cdot dx$  can usually be most easily obtained by employing the formula [1] Art. 80, and this can always be done with safety when  $fx$  is finite, continuous, and single-valued between  $x = a$  and  $x = b$ .

Of course, if the indefinite integral is a multiple-valued function, we must choose the values of the indefinite integral corresponding to  $x = a$  and  $x = b$ , so that they may be ordinates of the same branch of the curve  $y = \int fx \cdot dx$ .

Consider, for example,  $\int_{-1}^1 \frac{dx}{1+x^2}$ . The indefinite integral  $\int \frac{dx}{1+x^2} = \tan^{-1}x$  and  $\tan^{-1}x$  is a multiple-valued function. Indeed,  $y = \tan^{-1}x$  is a curve consisting of an infinite number of separate branches so related that ordinates corresponding to the same value of  $x$  differ by multiples of  $\pi$ . On the branch which passes through the origin, when  $x = -1$ ,  $y = \tan^{-1}x = -\frac{\pi}{4}$ ; on the same branch, when  $x = 1$ ,  $y = \tan^{-1}x = \frac{\pi}{4}$ . On the next branch above, when  $x = -1$ ,  $y = \tan^{-1}x = \frac{3\pi}{4}$ ; and when  $x = 1$ ,  $y = \frac{5\pi}{4}$ . On any branch, when  $x = -1$ ,  $y = \tan^{-1}x = -\frac{\pi}{4} + n\pi$ ; and on the same branch, when  $x = 1$ ,  $y = \frac{\pi}{4} + n\pi$ .

$$\text{Hence } \int_{-1}^1 \frac{dx}{1+x^2} = \tan^{-1}(1) - \tan^{-1}(-1) = \frac{\pi}{4} + \frac{\pi}{4} = \frac{\pi}{2};$$

$$\text{or } \int_{-1}^1 \frac{dx}{1+x^2} = \frac{5\pi}{4} - \frac{3\pi}{4} = \frac{\pi}{2},$$

$$\text{or } \int_{-1}^1 \frac{dx}{1+x^2} = \frac{\pi}{4} + n\pi - \left(-\frac{\pi}{4} + n\pi\right) = \frac{\pi}{2}.$$

By  $\int_a^{\infty} f(x).dx$  we mean the limit approached by  $\int_a^b f(x).dx$  as  $b$  is indefinitely increased.

## EXAMPLES.

(1) Work the examples of Art. 81 by the method of Art. 82

$$(2) \int_0^{\frac{\pi}{2}} \frac{\sin x . dx}{\cos^2 x} = \sqrt{2} - 1.$$

$$(3) \int_0^a \frac{dx}{\sqrt{x+a} + \sqrt{x}} = \frac{4}{3} \sqrt{a} (\sqrt{2} - 1).$$

$$(4) \int_0^{\infty} \frac{dx}{a^2 + x^2} = \frac{\pi}{2a}.$$

$$(5) \int_0^{\infty} \frac{ax}{a^2 + x^2} dx = \frac{\pi}{2} \text{ if } a > 0, \text{ and } -\frac{\pi}{2} \text{ if } a < 0, \text{ and } 0 \text{ if } a = 0.$$

$$(6) \int_0^{\infty} e^{-ax} dx = \frac{1}{a} \text{ if } a > 0.$$


$$(7) \int_0^{\infty} e^{-ax} \sin mx dx = \frac{m}{a^2 + m^2} \text{ if } a > 0.$$

$$(8) \int_0^{\infty} e^{-ax} \cos mx dx = \frac{a}{a^2 + m^2} \text{ if } a > 0.$$

$$(9) \int_0^1 \frac{dx}{1 + 2x \cos \phi + x^2} = \frac{\phi}{2 \sin \phi}.$$

$$(10) \int_0^{\infty} \frac{dx}{1 + 2x \cos \phi + x^2} = \frac{\phi}{\sin \phi}.$$

83. When  $fx$  is finite and single-valued between  $x = a$  and  $x = b$ , but has a *finite* discontinuity at some intermediate value  $x = c$



$$\int_a^b fx dx = \int_a^c fx dx + \int_c^b fx dx,$$

and therefore  $\int_a^b fx dx$  can be found by Art. 82 when the indefinite integral  $\int fx dx$  can be obtained; but when  $fx$  becomes *infinite* for  $x = a$ , or for  $x = b$ , or for some intermediate value  $x = c$ , special care must be exercised, and some special investigation is usually required.

If  $fx$  is infinite when  $x = a$  and  $\int_{a+\epsilon}^b fx dx$  approaches a finite limit as  $\epsilon$  approaches zero, this limit is what we shall mean by  $\int_a^b fx dx$ ; if  $\int_{a+\epsilon}^b fx dx$  increases indefinitely as  $\epsilon$  approaches zero, we shall say that  $\int_a^b fx dx$  is infinite; and if  $\int_{a+\epsilon}^b fx dx$  neither approaches a finite limit nor increases indefinitely as  $\epsilon$

approaches zero, we shall say that  $\int_a^b f x . dx$  is indeterminate. It is in the first case only that  $\int_a^b f x . dx$  can be safely employed in mathematical work.

If  $f x$  is infinite when  $x = b$  and  $\int_a^{b-\epsilon} f x . dx$  approaches a finite limit as  $\epsilon$  approaches zero, that limit is the value of  $\int_a^b f x . dx$ .

If  $f x$  is infinite when  $x = c$ , and each of the expressions  $\int_a^{c-\epsilon} f x . dx$  and  $\int_{c+\epsilon}^b f x . dx$  approaches a finite limit as  $\epsilon$  approaches zero, the sum of these limits is  $\int_a^b f x . dx$ . Should either or both of the expressions,

$$\int_a^{c-\epsilon} f x . dx, \quad \int_{c+\epsilon}^b f x . dx,$$

fail to approach a finite limit as  $\epsilon$  approaches zero,  $\int_a^b f x . dx$  is either infinite or indeterminate, and cannot be safely used.

When the indefinite integral of  $f x . dx$  can be obtained there is little difficulty in deciding on the nature of  $\int_a^b f x . dx$  in any of the cases just considered, or in getting its value when that value is finite and determinate.

For example,

(a)  $\int_0^1 \frac{dx}{x}$  is infinite, since

$$\int \frac{dx}{x} = \log x \text{ and } \int_{\epsilon}^1 \frac{dx}{x} = \log(1) - \log \epsilon = \log \frac{1}{\epsilon},$$

and increases indefinitely as  $\epsilon$  approaches zero.

(b)  $\int_0^{\infty} \frac{dx}{1-x^2}$  is not finite and determinate, for

$$\int \frac{dx}{1-x^2} = \frac{1}{2} \log \frac{1+x}{1-x},$$

$$\int_0^{1-\epsilon} \frac{dx}{1-x^2} = \frac{1}{2} \log \left( \frac{2-\epsilon}{\epsilon} \right),$$

and increases indefinitely as  $\epsilon$  approaches zero.

(c)  $\int_0^a \frac{dx}{\sqrt{a^2 - x^2}}$  is finite and determinate, for

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a},$$

$$\int_0^{a-\epsilon} \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{a-\epsilon}{a} - \sin^{-1} 0 = \sin^{-1} \frac{a-\epsilon}{a},$$

and its limiting value as  $\epsilon$  approaches zero is  $\sin^{-1}(1)$  or  $\frac{\pi}{2}$ .

(d)  $\int_0^2 \frac{x dx}{(1-x)^{\frac{1}{2}}}$  is finite and determinate, for

$$\int \frac{x dx}{(1-x)^{\frac{1}{2}}} = \frac{2}{3}(1-x)^{\frac{3}{2}} - \frac{2}{5}(1-x)^{\frac{5}{2}},$$

$$\int_0^{1-\epsilon} \frac{x dx}{(1-x)^{\frac{1}{2}}} = \frac{2}{3}\epsilon^{\frac{3}{2}} - \frac{2}{5}\epsilon^{\frac{5}{2}} - \frac{2}{3} + \frac{2}{5},$$

and its limiting value as  $\epsilon$  approaches zero is  $\frac{2}{3} - \frac{2}{5}$ .


$$\int_{1+\epsilon}^2 \frac{x dx}{(1-x)^{\frac{1}{2}}} = -\frac{2}{3} - \frac{2}{5} + \frac{2}{3}\epsilon^{\frac{3}{2}} + \frac{2}{5}\epsilon^{\frac{5}{2}},$$

and its limiting value as  $\epsilon$  approaches zero is  $-\frac{2}{3} - \frac{2}{5}$ , and consequently

$$\int_0^2 \frac{x dx}{(1-x)^{\frac{1}{2}}} = \frac{2}{3} - \frac{2}{5} - \frac{2}{3} - \frac{2}{5} = -\frac{4}{5}.$$

84. When, as is sometimes the case, the indefinite integral cannot be obtained, and the function to be integrated becomes infinite at or between the limits of integration, it is only necessary to investigate the limiting value of  $\epsilon f(a + \epsilon)$  as  $\epsilon$  approaches zero if  $fx$  becomes infinite when  $x = a$ ; of  $\epsilon f(b - \epsilon)$  if  $fx$  becomes infinite when  $x = b$ ; and of both  $\epsilon f(c - \epsilon)$  and  $\epsilon f(c + \epsilon)$  if  $fx$  becomes infinite when  $x = c$ . If each of the values in question has zero for its limit,  $\int_a^b fx dx$  is finite and determinate, otherwise it is infinite or indeterminate.

For, in order that the area  $EE'B'B$  should not increase indefinitely as  $\epsilon$  approaches zero, the rectangle  $AA'E'E$ , whose area is  $\epsilon f(a + \epsilon)$ , must approach zero as its limit; and the same reasoning holds good for the other cases considered above.



Let us apply this test to the examples considered in Art. 83.

(a)  $\int_0^1 \frac{dx}{x} = \infty$  because  $\lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon}{\epsilon} \right] = 1$ .

(b)  $\int_0^\infty \frac{dx}{1-x^2}$  is indeterminate, for

$$\lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon}{1-(1-\epsilon)^2} \right] = \lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon}{2\epsilon - \epsilon^2} \right] = \lim_{\epsilon \rightarrow 0} \left[ \frac{1}{2-\epsilon} \right] = \frac{1}{2},$$

and

$$\lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon}{1-(1+\epsilon)^2} \right] = -\frac{1}{2}.$$

(c)  $\int_0^a \frac{dx}{\sqrt{a^2-x^2}}$  is finite and determinate, for

$$\lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon}{\sqrt{a^2-(a-\epsilon)^2}} \right] = \lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon}{\sqrt{2a\epsilon - \epsilon^2}} \right] = \lim_{\epsilon \rightarrow 0} \left[ \frac{1}{\sqrt{\frac{2a}{\epsilon} - 1}} \right] = 0.$$

(d)  $\int_0^1 \frac{x dx}{(1-x)^{\frac{1}{2}}}$  is finite and determinate, for

$$\lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon(1-\epsilon)}{(1-(1-\epsilon))^{\frac{1}{2}}} \right] = \lim_{\epsilon \rightarrow 0} [\epsilon^{\frac{1}{2}}(1-\epsilon)] = 0,$$

and

$$\lim_{\epsilon \rightarrow 0} \left[ \frac{\epsilon(1+\epsilon)}{(1-(1+\epsilon))^{\frac{1}{2}}} \right] = -\lim_{\epsilon \rightarrow 0} [\epsilon^{\frac{1}{2}}(1+\epsilon)] = 0.$$



Even when, as in the examples just given, the indefinite integral can be obtained, there is a decided advantage in using the very simple method of this article. For if the application of the test shows that the definite integral in question is infinite or indeterminate, the labor of finding the indefinite integral is saved; and if the application of the test proves the definite integral finite and determinate, it follows that the indefinite integral does not become infinite for the value of  $x$  which makes the given function infinite, and consequently when the indefinite integral has been obtained, the method of Art. 82 can be used without hesitation.

As an example, where the indefinite integral cannot be obtained, let us consider at some length

$$\int_0^1 \left(\log \frac{1}{x}\right)^n dx.$$

If  $n$  is positive,  $\left(\log \frac{1}{x}\right)^n$  is continuous and single-valued between  $x=0$  and  $x=1$ , but becomes infinite when  $x=0$ . We must then investigate the limiting value of  $\epsilon \left(\log \frac{1}{\epsilon}\right)^n$  as  $\epsilon$  approaches zero.

$\epsilon \left(\log \frac{1}{\epsilon}\right)^n$  is indeterminate when  $\epsilon=0$ , but its *true value* is easily found to be zero if  $n$  is positive, whether  $n$  is whole or fractional. For positive values of  $n$ ,  $\int_0^1 \left(\log \frac{1}{x}\right)^n dx$  is, then, finite and determinate.

If  $n$  is negative, call  $n = -m$ .

$$\text{Then} \quad \int_0^1 \left(\log \frac{1}{x}\right)^n dx = \int_0^1 \frac{dx}{\left(\log \frac{1}{x}\right)^m}.$$

$$\frac{1}{\left(\log \frac{1}{x}\right)^m}$$

is continuous and single-valued from  $x=0$  to  $x=1$ , but becomes infinite when  $x=1$ .

We must, then, find  $\lim_{\epsilon \doteq 0} \left[ \frac{\epsilon}{\left(\log \frac{1}{1-\epsilon}\right)^m} \right]$ ,

which proves to be  $\lim_{\epsilon \doteq 0} \left[ \frac{1-\epsilon}{m \left(\log \frac{1}{1-\epsilon}\right)^{m-1}} \right]$ :

if  $m < 1$ , this is zero; if  $m = 1$ , it is 1; and if  $m > 1$ , it is infinite.

$$\int_0^1 \frac{dx}{\left(\log \frac{1}{x}\right)^m}$$

is, then, finite and determinate if  $m < 1$ , but infinite if  $m =$  or  $m > 1$ ; and we reach the result that

$$\int_0^1 \left(\log \frac{1}{x}\right)^n dx$$

is finite and determinate if  $n > -1$ , but infinite if  $n = -1$  or  $n < -1$ .

#### EXAMPLES.

(1) Prove that

$$\int_0^1 \frac{\log x}{1-x} \cdot dx, \int_0^1 \frac{\log x}{1-x^2} \cdot dx, \int_0^1 \frac{dx}{x} \log \left(\frac{1+x}{1-x}\right),$$

are finite and determinate.

(2) Prove that

$$\int_0^\infty \frac{dx}{1-x^a}, \int_0^\infty \frac{x^m dx}{1-x^m}, \text{ where } m \text{ and } n \text{ are integers, and}$$

$$\int_0^\infty \frac{x^{a-1}}{1-x} \cdot dx, \text{ are not finite and determinate.}$$

(3) Find for what values of  $n$   $\int_0^1 (\log x)^n dx$  is finite and determinate.

(4) Find for what values of  $m$  and  $n$   $\int_0^1 x^m \left(\log \frac{1}{x}\right)^n dx$  is finite and determinate.

(5) Show that  $\int_0^1 x^{m-1} (1-x)^{n-1} dx$  is finite and determinate if  $m$  and  $n$  are positive.

(6) Prove that  $\int_0^{\frac{\pi}{2}} \log \sin x \cdot dx$  is finite and determinate.

(7) Show that the following integrals are finite and determinate, and obtain their values :

$$\int_0^a \frac{dx}{\sqrt{a^2 - x^2}} = \frac{\pi}{2}.$$

$$\int_0^a \frac{dx}{\sqrt{ax - x^2}} = \pi.$$

$$\int_1^2 \frac{dx}{x\sqrt{x^2 - 1}} = \frac{\pi}{3}.$$


85. It was stated in Art. 82 that by  $\int_a^\infty f(x) \cdot dx$  we mean the limit approached by  $\int_a^b f(x) \cdot dx$  as  $b$  is indefinitely increased, and, as we have seen, if the indefinite integral  $\int f(x) \cdot dx$  can be found, there is no difficulty in investigating the nature of  $\int_a^\infty f(x) \cdot dx$  and in obtaining its value if it is finite and determinate. There are, however, many exceedingly important definite integrals of the form  $\int_a^\infty f(x) \cdot dx$  whose values are obtained by ingenious devices without employing the indefinite integral, and these devices are valid only provided that the integral in question is finite and determinate, since an infinite value not recognized and treated

as such, or a value absolutely indeterminate, renders inconclusive any piece of mathematical reasoning into which it enters.

If we construct the curve

$y = fx, \int_a^{\infty} fx \cdot dx$  is the limiting

value approached by the area  $ABB_1A_1$ , as  $OB_1$  is indefinitely increased; and in order that this area should



be finite and determinate, it is clearly necessary that the area  $BCC_1B_1$  should approach zero as its limit as  $OB_1$  and  $OC_1$  are indefinitely increased, however great the amount by which  $OC_1$  exceeds  $OB_1$ .

That is,

$$\lim_{\substack{b = \infty \\ c = \infty}} \left[ \int_b^c fx \cdot dx \right]$$

must be equal to zero no matter how much more rapidly  $c$  increases than  $b$ .

86. The investigation of the limiting value of  $\int_a^c fx \cdot dx$ , as  $b$  and  $c$  are indefinitely increased, is usually made with the aid of the following important theorem known as the *Maximum-Minimum Theorem*.

*If a given function of  $x$  is the product of two functions, one of which  $v$  does not change its sign between  $x = a$  and  $x = b$ , and if  $M$  is algebraically the greatest and  $N$  the least value of the other factor  $u$  between  $x = a$  and  $x = b$ ,  $\int_a^b uv \cdot dx$  lies between  $M \int_a^b v \cdot dx$  and  $N \int_a^b v \cdot dx$ .*

To prove this theorem, let us first suppose that  $v$  is positive between  $x = a$  and  $x = b$ . Now,  $M - u$  is positive for the values of  $x$  considered,  $(M - u)v$  is positive, and therefore

$$\int_a^b (M - u)v \cdot dx > 0 \quad \text{and} \quad M \int_a^b v \cdot dx > \int_a^b uv \cdot dx. \quad (1)$$

$u - N$  also is positive for all values of  $x$  between  $x = a$  and  $x = b$ ,  $(u - N)v$  is positive, and therefore

$$\int_a^b (u - N)v \, dx > 0 \quad \text{and} \quad \int_a^b uv \, dx > N \int_a^b v \, dx. \quad (2)$$

Hence  $\int_a^b uv \, dx$  lies between  $M \int_a^b v \, dx$  and  $N \int_a^b v \, dx$ .

It is easy to modify this proof to meet the case where  $v$  is negative.

(a) As an example of the use of this theorem, we will prove  $\int_0^{\infty} e^{-x^2} dx$  finite and determinate.

$e^{-x^2}$  is single-valued, finite, and continuous for all positive values of  $x$ ; if, then, we can show that  $\int_b^c e^{-x^2} dx$  has the limit zero as  $b$  and  $c$  are indefinitely increased,  $c$  remaining greater than  $b$ , our proof is complete.

$e^{-x^2}$  can be written  $x^2 e^{-x^2} \cdot \frac{1}{x^2}$ , and  $\frac{1}{x^2}$  never changes its sign.

As  $x$  increases  $x^2 e^{-x^2}$  eventually decreases, and continues to decrease toward zero as  $x$  increases indefinitely, as may be proved by determining its value for  $x = \infty$ .

We have 
$$\left[ \frac{x^2}{e^{x^2}} \right]_{x=\infty} = \left[ \frac{2x}{2xe^{x^2}} \right]_{x=\infty} = \left[ \frac{1}{e^{x^2}} \right]_{x=\infty} = 0.$$

Hence, eventually the greatest value of  $x^2 e^{-x^2}$  between  $x = b$  and  $x = c$  is  $b^2 e^{-b^2}$ , and the least value is  $c^2 e^{-c^2}$ .

Therefore, by the Maximum-Minimum Theorem,

$$b^2 e^{-b^2} \int_b^c \frac{dx}{x^2} > \int_b^c x^2 e^{-x^2} \cdot \frac{dx}{x^2} > c^2 e^{-c^2} \int_b^c \frac{dx}{x^2},$$

$$b^2 e^{-b^2} \left( \frac{1}{b} - \frac{1}{c} \right) > \int_b^c e^{-x^2} dx > c^2 e^{-c^2} \left( \frac{1}{b} - \frac{1}{c} \right).$$

As  $b$  and  $c$  are indefinitely increased, the first and third members of our last inequality approach the same limit, zero. Consequently the limiting value of  $\int_b^c e^{-x^2} dx$  is zero, and

$\int_0^\infty e^{-x^2} dx$  is finite and determinate.

(b) Let us consider  $\int_0^\infty \frac{\sin ax}{x} \cdot dx$ .

$\frac{\sin ax}{x}$  is equal to  $a$  when  $x = 0$ , and is single-valued, finite, and continuous for all positive values of  $x$ .

By integration by parts,

$$\int \frac{\sin ax}{x} dx = -\frac{\cos ax}{ax} - \frac{1}{a} \int \frac{\cos ax}{x^2} \cdot dx.$$

Therefore

$$\int_b^c \frac{\sin ax}{x} dx = \frac{\cos ab}{ab} - \frac{\cos ac}{ac} - \frac{1}{a} \int_b^c \frac{\cos ax}{x^2} \cdot dx.$$

As  $b$  and  $c$  are indefinitely increased,  $\frac{\cos ab}{ab}$  and  $\frac{\cos ac}{ac}$  approach the limiting value zero.

$\frac{1}{x^2}$  does not change sign between  $x = b$  and  $x = c$ , and the greatest value of  $\cos ax$  is 1, and its least value is  $-1$ .

$$\text{Hence } 1 \int_b^c \frac{dx}{x^2} > \int_b^c \frac{\cos ax}{x^2} dx > -1 \int_b^c \frac{dx}{x^2},$$

$$\text{or } \frac{1}{b} - \frac{1}{c} > \int_b^c \frac{\cos ax}{x^2} dx > -\left(\frac{1}{b} - \frac{1}{c}\right);$$

and the limiting value of  $\int_b^c \frac{\cos ax}{x^2} dx$  as  $b$  and  $c$  increase is zero. The same thing is true of  $\int_b^c \frac{\sin ax}{x} dx$ ; and therefore

$$\int_0^\infty \frac{\sin ax}{x} dx \text{ is finite and determinate.}$$

(c)  $\int_0^\infty \cos(x^2) dx$  is finite and determinate.

For  $\cos(x^2)$  is single-valued, finite, and continuous for all positive values of  $x$ ; and it is not difficult to prove that  $\int_b^c \cos(x^2) dx$  approaches the limit zero as  $b$  and  $c$  are indefinitely increased.

We have

$$\int_b^c \cos(x^2) dx = \int_b^c \frac{2x \cos(x^2) dx}{2x} = \left[ \frac{\sin(x^2)}{2x} \right]_b^c + \frac{1}{2} \int_b^c \frac{\sin(x^2) dx}{x^2}.$$

The limiting value of  $\frac{\sin(c^2)}{2c} - \frac{\sin(b^2)}{2b}$  as  $b$  and  $c$  are increased indefinitely is zero; and the limiting value of  $\int_b^c \frac{\sin(x^2) dx}{x^2}$  can be proved zero by the method followed in (b).

#### EXAMPLES.

(1) Construct the curves  $y = e^{-x^2}$ ;  $y = \frac{\sin ax}{x}$ ;  $y = \cos(x^2)$ .

(2) Prove that the following integrals are finite and determinate:

$$\int_0^{\infty} \frac{\sin^2 x}{x^3} dx, \quad \int_0^{\infty} \frac{\sin x}{\sqrt{x}} dx, \quad \int_0^{\infty} \frac{e^{-ax} \sin mx}{x} dx,$$

$$\int_0^{\infty} e^{-ax^2} \cos bx dx, \quad \int_0^{\infty} e^{-ax^2} x^{2n} dx, \quad \int_0^{\infty} e^{-x^2} \frac{x^2}{x^2} dx,$$

$$\int_0^{\infty} \log \left( \frac{e^x + 1}{e^x - 1} \right) dx.$$

(3) Show that  $\int_0^{\infty} x^n e^{-x} dx$  is finite and determinate for all values of  $n$  greater than  $-1$ .

87. When we have occasion to use a reduction formula in finding the value of a definite integral, it is often worth while to substitute the limits of integration in the general formula before attempting to apply it to the particular problem.

For example, let us find  $\int_0^a \frac{x^a dx}{\sqrt{a^2 - x^2}}$ .

We can reduce the exponent of  $x$  by [4], Art. 64,

$$\int x^{m-1} z^p dx = \frac{x^{m-n} z^{p+1}}{b(m+np)} - \frac{a(m-n)}{b(m+np)} \int x^{m-n-1} z^p dx.$$

For our example this becomes

$$\int x^{m-1}(a^2-x^2)^{-\frac{1}{2}}dx = \frac{x^m \frac{1}{2}(a^2-x^2)^{\frac{1}{2}}}{-m+1} - \frac{a^2(m-2)}{-m+1} \int x^{m-3}(a^2-x^2)^{-\frac{1}{2}}dx.$$

When  $x = 0$ , and also when  $x = a$ ,  $\frac{x^{m-2}(a^2-x^2)^{\frac{1}{2}}}{-m+1} = 0$ .

Hence

$$\int_0^a x^{m-1}(a^2-x^2)^{-\frac{1}{2}}dx = \frac{a^2(m-2)}{m-1} \int_0^a x^{m-3}(a^2-x^2)^{-\frac{1}{2}}dx;$$

$$\begin{aligned} \int_0^a x^6(a^2-x^2)^{-\frac{1}{2}}dx &= \frac{5}{6} \cdot a^2 \int_0^a x^4(a^2-x^2)^{-\frac{1}{2}}dx \\ &= \frac{5}{6} \cdot \frac{3}{4} \cdot a^4 \int_0^a x^2(a^2-x^2)^{-\frac{1}{2}}dx \\ &= \frac{5}{6} \cdot \frac{3}{4} \cdot \frac{1}{2} \cdot a^6 \int_0^a \frac{dx}{\sqrt{a^2-x^2}}. \end{aligned}$$

Therefore  $\int_0^a \frac{x^6 dx}{\sqrt{a^2-x^2}} = \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot \frac{\pi a^6}{2}$ .

EXAMPLES.

(1)  $\int_0^a \frac{x^5 dx}{\sqrt{a^2-x^2}} = \frac{2}{3} \cdot \frac{4}{5} a^5.$

(2)  $\int_0^a \sqrt{a^2-x^2} \cdot dx = \frac{\pi a^2}{4}.$

(3)  $\int_0^a x^2 \sqrt{a^2-x^2} \cdot dx = \frac{1}{4} \cdot \frac{\pi a^4}{4}.$

(4)  $\int_0^a x^2(a^2-x^2)^{\frac{3}{2}} \cdot dx = \frac{1}{6} \cdot \frac{3}{16} \cdot \pi a^6.$

(5)  $\int_0^{\frac{\pi}{2}} \sin^n x \cdot dx = \frac{1 \cdot 3 \cdot 5 \dots (n-1)}{2 \cdot 4 \cdot 6 \dots n} \cdot \frac{\pi}{2}$  when  $n$  is even.  
 $= \frac{2 \cdot 4 \cdot 6 \dots (n-1)}{3 \cdot 5 \cdot 7 \dots n}$  when  $n$  is odd.



(6) Show that  $\int_0^{\frac{\pi}{2}} \cos^n x \, dx = \int_0^{\frac{\pi}{2}} \sin^n x \, dx$ .

(7)  $\int_0^1 \frac{x^{2n} \, dx}{\sqrt{1-x^2}} = \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} \cdot \frac{\pi}{2}$ .

*Suggestion:* let  $x = \sin \theta$ .

(8)  $\int_0^1 \frac{x^{2n+1} \, dx}{\sqrt{1-x^2}} = \frac{2.4.6 \dots 2n}{3.5.7 \dots (2n+1)}$ .

(9) From Exs. 7 and 8 obtain Wallis's formula

$$\frac{\pi}{2} = \frac{2.2.4.4.6.6.8.8 \dots}{1.3.3.5.5.7.7.9 \dots}$$

*Suggestion:*  $\int_0^1 \frac{x^{2n-1} \, dx}{\sqrt{1-x^2}} > \int_0^1 \frac{x^{2n} \, dx}{\sqrt{1-x^2}} > \int_0^1 \frac{x^{2n+1} \, dx}{\sqrt{1-x^2}}$ .

88. When in finding  $\int_a^b f(x) \, dx$  the method of *integration by substitution* is used, and  $y = Fx$  is introduced in place of  $x$ , we can regard the new integral as a definite integral, the limits of integration being  $Fa$  and  $Fb$ , and thus avoid the labor of replacing  $y$  by its value in terms of  $x$  in the result of the indefinite integration.

Let us find  $\int_{-\infty}^0 e^{ax} \sqrt{1-e^{2ax}} \, dx$ .

Substitute  $y = e^{ax}$ .

$$dy = ae^{ax} \, dx.$$

Hence  $\int e^{ax} \sqrt{1-e^{2ax}} \, dx = \frac{1}{a} \int \sqrt{1-y^2} \, dy$ .

When  $x = -\infty$ ,  $y = 0$ , and when  $x = 0$ ,  $y = 1$ .

Therefore  $\int_{-\infty}^0 e^{ax} \sqrt{1-e^{2ax}} \, dx = \frac{1}{a} \int_0^1 \sqrt{1-y^2} \, dy = \frac{\pi}{4a}$ .

There is one rather rare class of cases where special care is needed in using the method just described. It is when  $y$  has a maximum or a minimum value between  $x = a$  and  $x = b$ , say for  $x = c$ , and  $x$  is a multiple-valued function of  $y$ .

For suppose  $y$  a maximum when  $x = c$ , then as  $x$  increases from  $a$  to  $b$ ,  $y$  increases to the value  $Fc$ , and then decreases to the value  $Fb$ , instead of simply increasing or decreasing from  $Fa$  to  $Fb$ . If  $x$  is a single-valued function of  $y$ , and  $\phi(y)dy$  is the result of substituting  $y$  for  $x$  in  $fx.dx$ ,  $\phi y$  is a single-valued function of  $y$ , and  $\int_{Fa}^{Fb} \phi y . dy = \int_{Fa}^{Fc} \phi y . dy + \int_{Fc}^{Fb} \phi y . dy$ , and there is no error in using  $\int_{Fa}^{Fb} \phi y . dy$  for  $\int_a^b fx . dx$ . But if  $x$  is a multiple-valued function of  $y$ , it will always happen that when  $y$  passes through a maximum, we pass from one set of values of  $x$  to another, and therefore from one set of values of  $\phi y$  to another, and in that case it is necessary to express our required integral as  $\int_{Fa}^{Fc} \phi y . dy + \int_{Fc}^{Fb} \phi y . dy$ , taking pains to select the correct set of values for  $\phi y$  in each integral.

If  $y$  is a minimum between  $x = a$  and  $x = b$ , essentially the same reasoning holds good.

A couple of examples will make this clearer.

(a) Take 
$$\int_0^{2a} \frac{x . dx}{\sqrt{2ax - x^2}} .$$

Let  $y = 2ax - x^2$ . Then  $\frac{dy}{dx} = 2(a - x) = 0$  when  $x = a$ .

$\frac{d^2y}{dx^2} = -2$ , and  $y$  is a maximum when  $x = a$ .

$$x = a \pm \sqrt{a^2 - y} ,$$

$$dx = \mp \frac{dy}{2\sqrt{a^2 - y}} .$$

Since  $\frac{dy}{dx}$  is positive from  $x = 0$  to  $x = a$ , and negative from  $x = a$  to  $x = 2a$ ,  $dx = \frac{dy}{2\sqrt{a^2 - y}}$  and  $x = a - \sqrt{a^2 - y}$  from  $x = 0$  to  $x = a$ , and  $dx = -\frac{dy}{2\sqrt{a^2 - y}}$ , and  $x = a + \sqrt{a^2 - y}$  from  $x = a$  to  $x = 2a$ .

Hence

$$\begin{aligned} \int_0^{2a} \frac{x dx}{\sqrt{2ax-x^2}} &= \int_0^a \frac{x dx}{\sqrt{2ax-x^2}} + \int_a^{2a} \frac{x dx}{\sqrt{2ax-x^2}} \\ &= \frac{1}{2} \int_0^{a^2} \frac{a - \sqrt{a^2-y}}{\sqrt{a^2y-y^2}} \cdot dy - \frac{1}{2} \int_a^0 \frac{a + \sqrt{a^2-y}}{\sqrt{a^2y-y^2}} \cdot dy \\ &= \frac{1}{2} \int_0^{a^2} \frac{a - \sqrt{a^2-y}}{\sqrt{a^2y-y^2}} \cdot dy + \frac{1}{2} \int_0^{a^2} \frac{a + \sqrt{a^2-y}}{\sqrt{a^2y-y^2}} \cdot dy \\ &= \int_0^{a^2} \frac{ady}{\sqrt{a^2y-y^2}} = \pi a. \quad (\text{Ex. 7, Art. 84}) \end{aligned}$$

$$(b) \quad \int_0^{\frac{\pi}{2}} \frac{dx}{(\sin x + \cos x)^2}$$

Let  $y = \sin x + \cos x$ .  $\frac{dy}{dx} = \cos x - \sin x = 0$  when  $x = \frac{\pi}{4}$ .

$\frac{d^2y}{dx^2} = -\sin x - \cos x = -\sqrt{2}$  when  $x = \frac{\pi}{4}$ . Therefore  $y$  has a maximum value  $\sqrt{2}$  when  $x = \frac{\pi}{4}$ .

$$y = \sin x + \cos x = \sqrt{2} \cdot \cos\left(\frac{\pi}{4} - x\right),$$

$$x = \frac{\pi}{4} - \cos^{-1} \frac{y}{\sqrt{2}}, \quad dx = \pm \frac{dy}{\sqrt{2-y^2}}.$$

Since  $\frac{dy}{dx} = 0$  and  $\frac{d^2y}{dx^2} < 0$  when  $x = \frac{\pi}{4}$ , it follows that  $\frac{dy}{dx}$  is positive from  $x = 0$  to  $x = \frac{\pi}{4}$ , and negative from  $x = \frac{\pi}{4}$  to  $x = \frac{\pi}{2}$ .

Hence we have

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \frac{dx}{(\sin x + \cos x)^2} &= \int_0^{\frac{\pi}{4}} \frac{dx}{(\sin x + \cos x)^2} + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{dx}{(\sin x + \cos x)^2} \\ &= \int_1^{\sqrt{2}} \frac{dy}{y^2 \sqrt{2-y^2}} - \int_{\sqrt{2}}^1 \frac{dy}{y^2 \sqrt{2-y^2}} = 2 \int_1^{\sqrt{2}} \frac{dy}{y^2 \sqrt{2-y^2}}. \end{aligned}$$

Let  $\frac{y}{\sqrt{2}} = \sin \theta;$

$$\int_1^{\sqrt{2}} \frac{dy}{y^2 \sqrt{2-y^2}} = \frac{1}{2} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \csc^2 \theta d\theta = \frac{1}{2},$$

and

$$\int_0^{\frac{\pi}{2}} \frac{dx}{(\sin x + \cos x)^2} = 1.$$

EXAMPLE.

(1) Show that  $\int_0^{\pi} \frac{dx}{(\sin x + \cos x)^2} = \infty.$

89. *Differentiation of a definite integral.*

We have seen in Art. 51 that a definite integral is a function of the *limits of integration*, and not of the variable with respect to which we integrate; that is, that  $\int_a^b f x . dx$  is a function of  $a$  and  $b$ , and not a function of  $x$ . Strictly speaking,  $\int_a^b f x . dx$  is a function of  $a$  and  $b$ , and of any constants that  $f x$  may contain, where by *constant* we mean any quantity that is independent of  $x$ .

If the limits  $a$  and  $b$  are variables, they are always independent of the  $x$  with respect to which the integration is performed, which must from the nature of the case disappear when the definite integral is formed, as it always may be in theory, from the indefinite integral; and this assertion holds good even when the same letter which is used for the variable with respect to which the integration is performed appears explicitly in the limits of integration.

Thus if we write  $\int_0^x \sin x . dx$ , the  $x$  in  $\sin x . dx$  and the  $x$  which is the upper limit of integration do not represent the same variable, and are entirely unconnected. Indeed, the former  $x$

may be replaced by any other letter without affecting the value of the integral. For

$$\int_0^{\pi} \sin x \cdot dx = \int_0^{\pi} \sin z \cdot dz = 1 - \cos x.$$

Let us now consider the possibility of differentiating a definite integral.

Required  $D_a \int_a^b f(x, a) dx$ , where  $a$  is independent of  $x$ , and  $a$  and  $b$  do not depend upon  $a$ .

We have

$$\begin{aligned} D_a \int_a^b f(x, a) dx &= \lim_{\Delta a \doteq 0} \left[ \frac{\int_a^b f(x, a + \Delta a) dx - \int_a^b f(x, a) dx}{\Delta a} \right] \\ &= \lim_{\Delta a \doteq 0} \left[ \int_a^b \frac{f(x, a + \Delta a) - f(x, a)}{\Delta a} dx \right] \\ &= \int_a^b \left( \lim_{\Delta a \doteq 0} \left[ \frac{f(x, a + \Delta a) - f(x, a)}{\Delta a} \right] \right) dx. \end{aligned}$$

$$\text{Hence } D_a \int_a^b f(x, a) dx = \int_a^b [D_a f(x, a)] dx, \quad [1]$$

and we find that we have merely to differentiate under the sign of integration.

The truth of the converse of the last proposition can be easily established, and we have

$$\int \left[ \int_a^b f(x, a) dx \right] da = \int_a^b \left[ \int f(x, a) da \right] dx, \quad [2]$$

or even

$$\int_c^d \left[ \int_a^b f(x, a) dx \right] da = \int_a^b \left[ \int_c^d f(x, a) da \right] dx, \quad [3]$$

if  $a$ ,  $b$ ,  $c$ , and  $d$  are entirely independent.

Suppose now that we are dealing with variable limits of integration.

Let us find first  $\frac{d}{dz} \int_a^z f(x).dx$ .

Let  $\int f(x).dx = Fx$ , then  $\int_a^z f(x).dx = Fz - Fa$ ; and since by definition  $\frac{dFx}{dx} = fx$ , it follows that  $\frac{dFz}{dz} = fz$ .

$$\text{Hence } \frac{d}{dz} \int_a^z f(x).dx = \frac{d(Fz - Fa)}{dz} = fz. \quad [4]$$

In the same way it may be shown that

$$\frac{d}{dz} \int_z^b f(x).dx = -fz. \quad [5]$$

Let us now take the most complicated case, namely, to find  $\frac{d}{da} \int_a^b f(x, a) dx$ , where  $a$  and  $b$  are functions of  $a$ .

$$\text{Let } \int f(x, a) dx = F(x, a);$$

$$\text{then } u = \int_a^b f(x, a) dx = F(b, a) - F(a, a),$$

$$\text{and } \frac{du}{da} = \frac{dF(b, a)}{da} - \frac{dF(a, a)}{da};$$

but as  $b$  and  $a$  are functions of  $a$ ,

$$\frac{dF(b, a)}{da} = D_b F(b, a) \frac{db}{da} + D_a F(b, a),$$

$$\text{and } \frac{dF(a, a)}{da} = D_a F(a, a) \frac{da}{da} + D_a F(a, a),$$

by I. Art. 200.

$$D_b F(b, a) = f(b, a),$$

$$D_a F(a, a) = f(a, a).$$

$$\text{Hence } \frac{du}{da} = D_a [F(b, a) - F(a, a)] + f(b, a) \frac{db}{da} - f(a, a) \frac{da}{da},$$

or

$$\frac{d}{da} \int_a^b f(x, a) dx = \int_a^b (D_a f(x, a)) dx + f(b, a) \frac{db}{da} - f(a, a) \frac{da}{da}. \quad [6]$$

## EXAMPLES.

$$(1) \frac{d}{dy} \int_0^{\pi} \sin(x+y) dx = (x+1) \sin(xy+y) - \sin y.$$

$$(2) \frac{d}{dx} \int_0^{\sqrt{x}} x^3 dx = \frac{1}{3}.$$

$$(3) \frac{d}{d\phi} \int_0^{e^{\phi}} \sqrt{1 - \cos \phi} \cdot d\phi = e^{\phi} \sqrt{1 - \cos e^{\phi}}$$

90. When the indefinite integral cannot be found, the problem of obtaining the value of the definite integral usually becomes a more or less difficult mathematical puzzle, which can be solved, if solved at all, only by the exercise of great ingenuity. Some of the results arrived at, however, are so important, and some of the devices employed so interesting, that we shall present them briefly here. But we must repeat the warning that most of the methods are valid only in case the definite integral is finite and determinate; and erroneous results have more than once been obtained and published when a little attention to the precautions described in Articles 83–86 would have prevented the mistake.

91. *Integration by development in series.*

$$(a) \int_0^1 \frac{\log x}{1-x} dx. \quad (\text{v. Art. 84, Ex. 1.})$$

$$\frac{1}{1-x} = (1-x)^{-1} = 1 + x + x^2 + x^3 + \dots, \text{ if } x < 1.$$

$$\int_0^1 \frac{\log x}{1-x} dx = \int_0^1 (\log x + x \log x + x^2 \log x + \dots) dx.$$

$$\int_0^1 x^n \log x dx = -\frac{1}{(n+1)^2}. \quad (\text{v. Art. 55 (a).})$$

Therefore

$$\int_0^1 \frac{\log x}{1-x} dx = -\left(\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots\right) = -\frac{\pi^2}{6}.$$

(v. Todhunter's Trigonometry, Chap. XXIII., Ex. 1.)

$$(b) \int_0^{\infty} \log \left( \frac{e^x + 1}{e^x - 1} \right) dx. \quad (\text{v. Art. 86, Ex. 2.})$$

$$\begin{aligned} \log \left( \frac{e^x + 1}{e^x - 1} \right) &= \log \left( \frac{1 + e^{-x}}{1 - e^{-x}} \right) = \log (1 + e^{-x}) - \log (1 - e^{-x}) \\ &= e^{-x} - \frac{e^{-2x}}{2} + \frac{e^{-3x}}{3} - \frac{e^{-4x}}{4} \dots - \left( -e^{-x} - \frac{e^{-2x}}{2} - \frac{e^{-3x}}{3} - \frac{e^{-4x}}{4} \dots \right). \end{aligned}$$

(I. Art. 130.)

Hence

$$\begin{aligned} \int_0^{\infty} \log \left( \frac{e^x + 1}{e^x - 1} \right) dx &= 2 \int_0^{\infty} \left( e^{-x} + \frac{e^{-3x}}{3} + \frac{e^{-5x}}{5} + \frac{e^{-7x}}{7} + \dots \right) dx \\ &= 2 \left( 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots \right). \end{aligned}$$

But  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots = \frac{\pi^2}{8}.$

(v. Todhunter's Trig., Chap. XXIII., Ex. 1.)

Therefore  $\int_0^{\infty} \log \left( \frac{e^x + 1}{e^x - 1} \right) dx = \frac{\pi^2}{4}.$

EXAMPLES.

(1)  $\int_0^1 \frac{\log x}{1+x} dx = -\frac{\pi^2}{12}.$

(2)  $\int_0^1 \frac{\log x}{1-x^2} dx = -\frac{\pi^2}{8}.$

(3)  $\int_0^1 \frac{dx}{x} \cdot \log \left( \frac{1+x}{1-x} \right) = \frac{\pi^2}{4}.$

(4)  $\int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} = \frac{\pi}{2} \left[ 1 + \left( \frac{1}{2} \right)^2 k^2 + \left( \frac{1.3}{2.4} \right)^2 k^4 + \left( \frac{1.3.5}{2.4.6} \right)^2 k^6 + \dots \right]$  if  $k^2 < 1.$



$$(5) \int_0^{\frac{\pi}{2}} \sqrt{1 - k^2 \sin^2 \phi} \cdot d\phi = \frac{\pi}{2} \left[ 1 - \left(\frac{1}{2}\right)^2 \frac{k^2}{1} - \left(\frac{1.3}{2.4}\right)^2 \frac{k^4}{3} - \left(\frac{1.3.5}{2.4.6}\right)^2 \frac{k^6}{5} - \dots \right] \quad \text{if } k^2 < 1.$$

92. *Integration by ingenious devices.*

$$(a) \int_0^{\frac{\pi}{2}} \log \sin x \cdot dx. \quad (\text{v. Art. 84, Ex. 6.})$$

Let  $u = \int_0^{\frac{\pi}{2}} \log \sin x \cdot dx.$

Substitute  $y = \frac{\pi}{2} - x.$

$$u = - \int_{\frac{\pi}{2}}^0 \log \cos y \cdot dy = \int_0^{\frac{\pi}{2}} \log \cos x \cdot dx.$$

$$2u = \int_0^{\frac{\pi}{2}} (\log \sin x + \log \cos x) dx = \int_0^{\frac{\pi}{2}} \log (\sin x \cos x) dx$$

$$= \int_0^{\frac{\pi}{2}} \log \left( \frac{\sin 2x}{2} \right) \cdot dx$$

$$= -\frac{\pi}{2} \log (2) + \int_0^{\frac{\pi}{2}} \log \sin 2x \cdot dx$$

$$= -\frac{\pi}{2} \log (2) + \frac{1}{2} \int_0^{\pi} \log \sin x \cdot dx.$$

$$\int_0^{\pi} \log \sin x \cdot dx = \int_0^{\frac{\pi}{2}} \log \sin x \cdot dx + \int_{\frac{\pi}{2}}^{\pi} \log \sin x \cdot dx$$

$$= u + \int_{\frac{\pi}{2}}^{\pi} \log \sin x \cdot dx.$$

Substitute  $y = \pi - x,$

and

$$\int_{\frac{\pi}{2}}^{\pi} \log \sin x . dx = - \int_{\frac{\pi}{2}}^{\pi} \log \sin y . dy = \int_0^{\frac{\pi}{2}} \log \sin x . dx = u.$$

Hence  $2u = -\frac{\pi}{2} \log(2) + u,$

and

$$u = \int_0^{\frac{\pi}{2}} \log \sin x . dx = \int_0^{\frac{\pi}{2}} \log \cos x . dx = -\frac{\pi}{2} \log(2). \quad [1]$$

(b)  $\int_0^{\infty} e^{-x^2} dx.$  (v. Art. 86 (a).)

Let  $u = \int_0^{\infty} e^{-x^2} dx,$  and let  $x = ay;$

then  $u = \int_0^{\infty} ae^{-a^2y^2} dy = \int_0^{\infty} ae^{-a^2x^2} dx,$

$$ue^{-a^2} = \int_0^{\infty} ae^{-(1+a^2)x^2} dx,$$

$$u \int_0^{\infty} e^{-a^2} da = u^2 = \int_0^{\infty} \left( \int_0^{\infty} ae^{-(1+a^2)x^2} da \right) dx, \text{ by [3], Art. 89.}$$

But

$$\int_0^{\infty} ae^{-(1+a^2)x^2} da = \frac{1}{2} \frac{1}{1+x^2}.$$

Hence  $u^2 = \frac{1}{2} \int_0^{\infty} \frac{dx}{1+x^2} = \frac{\pi}{4},$

and  $\int_0^{\infty} e^{-x^2} dx = \frac{1}{2} \sqrt{\pi}.$  [2]

(c)  $\int_0^{\infty} \frac{\sin mx}{x} . dx.$  (v. Art. 86 (b).)

We have  $\frac{1}{x} = \int_0^{\infty} e^{-ax} da$  if  $x > 0.$  (Art. 82, Ex. 6.)

Hence

$$\begin{aligned} \int_0^{\infty} \frac{\sin mx}{x} \cdot dx &= \int_0^{\infty} \left( \sin mx \int_0^{\infty} e^{-ax} da \right) dx \\ &= \int_0^{\infty} \left( \int_0^{\infty} e^{-ax} \sin mx \cdot da \right) dx \\ &= \int_0^{\infty} \left( \int_0^{\infty} e^{-ax} \sin mx \cdot dx \right) da, \text{ by [3], Art. 89.} \\ &= \int_0^{\infty} \frac{m da}{a^2 + m^2}. \quad (\text{Art. 82, Ex. 7.}) \end{aligned}$$

Therefore

$$\left. \begin{aligned} \int_0^{\infty} \frac{\sin mx}{x} \cdot dx &= \frac{\pi}{2} \text{ if } m > 0 \\ &= -\frac{\pi}{2} \text{ if } m < 0 \\ &= 0 \text{ if } m = 0 \end{aligned} \right\}, \quad [3]$$

by Art. 82, Ex. 5.

EXAMPLES.

$$(1) \int_0^{\pi} x \log \sin x \cdot dx = -\frac{\pi^2}{2} \log(2).$$

$$(2) \int_0^{\infty} \log \left( x + \frac{1}{x} \right) \frac{dx}{1+x^2} = \pi \log(2).$$

*Suggestion:* let  $x = \tan \theta$ .

$$(3) \int_0^{\infty} e^{-ax^2} \cdot dx = \frac{1}{2a} \sqrt{\pi}.$$

$$(4) \int_0^1 \frac{dx}{\sqrt{\log \frac{1}{x}}} = \sqrt{\pi}.$$

$$\begin{aligned} (5) \int_0^{\infty} \frac{\sin x \cos mx}{x} \cdot dx &= 0 \text{ if } m < -1 \text{ or } m > 1 \\ &= \frac{\pi}{4} \text{ if } m = -1 \text{ or } m = 1 \\ &= \frac{\pi}{2} \text{ if } -1 < m < 1. \end{aligned}$$

$$(6) \int_0^{\infty} \frac{\sin^2 x}{x^2} dx = \frac{\pi}{2}. \quad \text{Suggestion: integrate by parts.}$$

93. *Differentiation or integration with respect to a quantity which is independent of x.* (v. Art. 89.)

$$(a) \text{ We have } \int_0^{\infty} e^{-ax} dx = \frac{1}{a}. \quad (\text{Art. 82, Ex. 6.})$$

Differentiate both members with respect to  $a$ ,

$$\int_0^{\infty} (-xe^{-ax}) dx = -\frac{1}{a^2} \quad \text{or} \quad \int_0^{\infty} xe^{-ax} dx = \frac{1}{a^2}.$$

Differentiate again,

$$\int_0^{\infty} x^2 e^{-ax} dx = \frac{2}{a^3}.$$

Differentiating  $n$  times,

$$\int_0^{\infty} x^n e^{-ax} dx = \frac{n!}{a^{n+1}}. \quad [1] \quad (\text{v. Art. 86, Ex. 3.})$$

$$(b) \text{ We have } \int_0^{\infty} e^{-ax^2} dx = \frac{1}{2} \frac{\sqrt{\pi}}{a^{\frac{1}{2}}}. \quad (\text{Art. 92, Ex. 3.})$$

Differentiating  $n$  times with respect to  $a$ ,

$$\int_0^{\infty} x^{2n} e^{-ax^2} dx = \frac{1.3.5 \dots (2n-1)}{2^{n+1} a^n} \sqrt{\frac{\pi}{a}}. \quad [2] \\ (\text{v. Art. 86, Ex. 2.})$$

$$(c) \text{ We have } \int_0^{\infty} e^{-cx} dx = \frac{1}{c}. \quad (\text{Art. 82, Ex. 6.})$$

Multiply by  $dc$ , and integrate from  $a$  to  $b$ ,

$$\int_0^{\infty} \left( \int_a^b e^{-cx} dc \right) dx = \int_a^b \frac{dx}{c}.$$

$$\text{Hence } \int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx = \log \frac{b}{a}. \quad [3]$$

$$(d) \quad \int_0^1 x^a dx = \frac{1}{a+1}.$$

Multiply by  $da$ , and integrate from  $b$  to  $a$ ,

$$\int_b^a \left( \int_0^1 x^a da \right) dx = \int_b^a \frac{da}{a+1}.$$

$$\text{Hence} \quad \int_0^1 \frac{x^a - x^b}{\log x} dx = \log \left( \frac{a+1}{b+1} \right). \quad [4]$$

EXAMPLES.

$$(1) \text{ From } \int_0^\infty \frac{dx}{x^2+a} = \frac{\pi}{2} \frac{1}{\sqrt{a}} \text{ obtain}$$

$$\int_0^\infty \frac{dx}{(x^2+a)^{n+1}} = \frac{\pi}{2} \frac{1 \cdot 3 \cdot 5 \dots (2n-1)}{2 \cdot 4 \cdot 6 \dots 2n} \cdot \frac{1}{\sqrt{a^{2n+1}}}.$$

$$(2) \text{ From } \int_0^1 x^n dx = \frac{1}{n+1} \text{ obtain}$$

$$\int_0^1 x^n (\log x)^m dx = (-1)^m \frac{m!}{(n+1)^{m+1}}.$$

$$(3) \text{ From } \int_0^\infty e^{-ax} \cos mx \cdot dx = \frac{a}{a^2+m^2} \text{ obtain}$$

$$\int_0^\infty \frac{e^{-ax} - e^{-bx}}{x} \cos mx \cdot dx = \frac{1}{2} \log \left( \frac{b^2+m^2}{a^2+m^2} \right).$$

$$(4) \text{ From } \int_0^\infty e^{-ax} \sin mx \cdot dx = \frac{m}{a^2+m^2} \text{ obtain}$$

$$\int_0^\infty \frac{e^{-ax} - e^{-bx}}{x} \sin mx \cdot dx = \tan^{-1} \frac{b}{m} - \tan^{-1} \frac{a}{m}.$$

94. The method illustrated in Art. 93 can be applied to much more complicated forms.

$$(a) \quad \int_0^\infty e^{-x-\frac{x^2}{2}} \cdot dx. \quad (\text{v. Art. 86, Ex. 2.})$$

Let 
$$u = \int_0^{\infty} e^{-x^2 - \frac{a^2}{x^2}} dx;$$

then 
$$\frac{du}{da} = -2 \int_0^{\infty} \frac{adx}{x^2} \cdot e^{-x^2 - \frac{a^2}{x^2}}.$$

Substitute 
$$z = \frac{a}{x},$$

and 
$$\frac{du}{da} = -2 \int_0^{\infty} e^{-x^2 - \frac{a^2}{x^2}} dz = -2 \int_0^{\infty} e^{-x^2 - \frac{a^2}{x^2}} dx = -2u.$$

Hence 
$$\frac{du}{u} = -2 da.$$

Integrating, 
$$\log u = -2a + C,$$

and 
$$u = C_1 e^{-2a}.$$

When  $a = 0$ , 
$$u = \int_0^{\infty} e^{-x^2} dx = \frac{1}{2} \sqrt{\pi}. \quad (\text{Art. 92 (b) [2].})$$

Therefore 
$$C_1 = \frac{1}{2} \sqrt{\pi},$$

and 
$$u = \int_0^{\infty} e^{-x^2 - \frac{a^2}{x^2}} dx = \frac{e^{-2a} \sqrt{\pi}}{2}. \quad [1]$$

(b) 
$$\int_0^{\infty} e^{-a^2 x^2} \cos bx \cdot dx. \quad (\text{v. Art. 86, Ex. 2.})$$

Let 
$$u = \int_0^{\infty} e^{-a^2 x^2} \cos bx \cdot dx,$$

then 
$$\frac{du}{db} = - \int_0^{\infty} x e^{-a^2 x^2} \sin bx \cdot dx.$$

Integrating by parts,

$$\int_0^{\infty} x e^{-a^2 x^2} \sin bx \cdot dx = \frac{b}{2a^2} \int_0^{\infty} e^{-a^2 x^2} \cos bx \cdot dx = \frac{b}{2a^2} u.$$

Therefore 
$$\frac{du}{db} = - \frac{b}{2a^2} u,$$

or 
$$\frac{du}{u} = - \frac{b}{2a^2} db.$$

Integrating, we have

$$\log u = -\frac{b^2}{4a^2} + C,$$

or 
$$u = C_1 e^{-\frac{b^2}{4a^2}}.$$

When  $b = 0$ , 
$$u = \int_0^\infty e^{-a^2 x^2} dx = \frac{\sqrt{\pi}}{2a}. \quad (\text{Art. 92, Ex. 3.})$$

Hence 
$$u = \int_0^\infty e^{-a^2 x^2} \cos bx \cdot dx = \frac{\sqrt{\pi}}{2a} e^{-\frac{b^2}{4a^2}}. \quad [2]$$

EXAMPLES.

(1) 
$$\int_0^\infty \frac{e^{-ax} \sin mx}{x} \cdot dx = \tan^{-1} \frac{m}{a}.$$

(2) 
$$\int_0^\infty \frac{\cos mx}{1+x^2} \cdot dx = \frac{\pi}{2} e^{-m}.$$

Suggestion: 
$$\frac{1}{1+x^2} = 2 \int_0^\infty a e^{-a^2(1+x^2)} da.$$

95. *Introduction of imaginary constants.*

$$\int_0^\infty \cos(x^2) dx. \quad (\text{v. Art. 86 (c.)})$$

We have 
$$\int_0^\infty e^{-a^2 x^2} dx = \frac{1}{2a} \sqrt{\pi}. \quad (\text{Art. 92, Ex. 3.})$$

Let 
$$a^2 = c^2 \sqrt{-1} = c^2 \left( \cos \frac{\pi}{2} + \sqrt{-1} \sin \frac{\pi}{2} \right).$$

Then 
$$a = c \left( \cos \frac{\pi}{4} + \sqrt{-1} \sin \frac{\pi}{4} \right) = \frac{c\sqrt{2}}{2} (1 + \sqrt{-1}), \quad (\text{Art. 25.})$$

and 
$$\frac{1}{2a} = \frac{1}{c\sqrt{2}(1 + \sqrt{-1})} = \frac{1}{2c\sqrt{2}} (1 - \sqrt{-1}).$$

Hence 
$$\int_0^{\infty} e^{-c^2 x^2 \sqrt{-1}} dx = \frac{1}{2c} \sqrt{\frac{\pi}{2}} \cdot (1 - \sqrt{-1}).$$

But 
$$e^{-c^2 x^2 \sqrt{-1}} = \cos(c^2 x^2) - \sqrt{-1} \sin(c^2 x^2). \tag{[5] Art. 31.}$$

Therefore

$$\int_0^{\infty} \cos(c^2 x^2) dx - \sqrt{-1} \int_0^{\infty} \sin(c^2 x^2) dx = \frac{1}{2c} \sqrt{\frac{\pi}{2}} (1 - \sqrt{-1}),$$

and

$$\int_0^{\infty} \cos(c^2 x^2) dx = \int_0^{\infty} \sin(c^2 x^2) dx = \frac{1}{2c} \sqrt{\frac{\pi}{2}}. \tag{[1] Art. 17.}$$

Let 
$$c = 1,$$

and 
$$\int_0^{\infty} \cos(x^2) dx = \int_0^{\infty} \sin(x^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2}}. \tag{[2]}$$

If we substitute  $y = x^2$  in [2], we get

$$\int_0^{\infty} \frac{\cos y}{\sqrt{y}} dy = \int_0^{\infty} \frac{\sin y}{\sqrt{y}} dy = \sqrt{\frac{\pi}{2}}. \tag{[3]}$$

*Gamma Functions.*

96. It was shown in Art. 84 that  $\int_0^1 \left(\log \frac{1}{x}\right)^n dx$  is finite and determinate for all values of  $n$  greater than  $-1$ , and infinite when  $n$  is equal to or less than  $-1$ . The substitution of  $y = \log \frac{1}{x}$  reduces this integral to  $\int_0^{\infty} y^n e^{-y} dy$ , or, what is the same thing, to  $\int_0^{\infty} x^n e^{-x} dx$ ; and in Art. 86, Ex. 3, the student has been required to show that this integral is finite and determinate for all values of  $n$  greater than  $-1$ .

$$\int x^n e^{-x} dx = -x^n e^{-x} + n \int x^{n-1} e^{-x} dx,$$

by integration by parts.



If  $n$  is greater than zero,

$$x^n e^{-x} = 0 \quad \text{when } x = 0,$$

and  $x^n e^{-x}$  is indeterminate when  $x = \infty$ . Its true value when  $x = \infty$ , obtained by the method of I. Art. 141, is, however, zero.

$$\text{Therefore} \quad \int_0^{\infty} x^n e^{-x} dx = n \int_0^{\infty} x^{n-1} e^{-x} dx \quad [1]$$

for all positive values of  $n$ .

If  $n$  is an integer, a repeated use of [1] gives

$$\int_0^{\infty} x^n e^{-x} dx = n! \int_0^{\infty} e^{-x} dx;$$

$$\text{but} \quad \int_0^{\infty} e^{-x} dx = 1,$$

$$\text{and we have} \quad \int_0^{\infty} x^n e^{-x} dx = n! \quad [2]$$

provided that  $n$  is a positive whole number.

If  $n$  is not a positive integer, but is greater than  $-1$ ,  $\int_0^{\infty} x^n e^{-x} dx$  is a finite and determinate function of  $n$ , and its value can be computed to any required degree of accuracy by methods which we have not space to consider here.

$\int_0^{\infty} x^{n-1} e^{-x} dx$  is generally represented by  $\Gamma(n)$ , and has been very carefully studied under the name of the *Gamma Function*.

If  $n$  is a positive integer, we have from [2]

$$\Gamma(n+1) = n!. \quad [3]$$

$$\text{From [3],} \quad \Gamma(2) = 1. \quad [4]$$

$$\text{Since} \quad \Gamma(1) = \int_0^{\infty} x^0 e^{-x} dx = \int_0^{\infty} e^{-x} dx,$$

$$\Gamma(1) = 1. \quad [5]$$

We have always from [1]

$$\Gamma(n+1) = n\Gamma(n), \quad [6]$$

if  $n$  is greater than zero.

Since  $\int_0^\infty x^n e^{-x} dx$  is infinite when  $n$  is equal to or less than  $-1$ , it follows from the definition of  $\Gamma(n)$  that  $\Gamma(n) = \infty$  if  $n$  is equal to or less than zero. It has, however, been found convenient to adopt formula [6] as the definition of  $\Gamma(n)$  when  $n$  is equal to or less than zero, and to restrict the original definition to positive values of  $n$ . The result easily deduced is that  $\Gamma(n)$  is infinite when  $n$  is equal to zero or to a negative integer, but is finite and determinate for all other values of  $n$ .

97. We may regard the formula

$$\Gamma(n + 1) = n\Gamma(n)$$

as a sort of *reduction formula*; and since each time we apply it we can raise or lower the value of  $n$  by unity, we can obtain any required *Gamma Function* by the aid of a table containing the values of  $\Gamma(n)$  corresponding to the values of  $n$  between any two arbitrarily chosen consecutive whole numbers.

Such tables have been computed, and we give one here containing the *common logarithms* of the values of  $\Gamma(n)$  from  $n = 1$  to  $n = 2$ . The table is carried out to four decimal places, and each logarithm is printed with the characteristic 9, which, of course, is ten units too large, the true characteristic being  $-1$ .

$10 + \log \Gamma(n)$ .

$n$	0	1	2	3	4	5	6	7	8	9
1.0	..	9975	9951	9928	9905	9883	9862	9841	9821	9802
1.1	9.9783	9765	9748	9731	9715	9699	9684	9669	9655	9642
1.2	9.9629	9617	9605	9594	9583	9573	9564	9554	9546	9538
1.3	9.9530	9523	9516	9510	9505	9500	9495	9491	9487	9483
1.4	9.9481	9478	9476	9475	9473	9473	9472	9473	9473	9474
1.5	9.9475	9477	9479	9482	9485	9488	9492	9496	9501	9506
1.6	9.9511	9517	9523	9529	9536	9543	9550	9558	9566	9575
1.7	9.9584	9593	9603	9613	9623	9633	9644	9656	9667	9679
1.8	9.9691	9704	9717	9730	9743	9757	9771	9786	9800	9815
1.9	9.9831	9846	9862	9878	9895	9912	9929	9946	9964	9982

Such a table enables us to compute with Gamma Functions as readily as with Trigonometric Functions, and consequently the problem of obtaining the value of a definite integral is practically solved if the integral in question can be expressed in terms of Gamma Functions.

For example, let us consider

$$(a) \quad \int_0^{\infty} x^n e^{-ax} dx.$$

$$\text{Let} \quad y = ax;$$

$$\text{then} \quad \int_0^{\infty} x^n e^{-ax} dx = \frac{1}{a^{n+1}} \int_0^{\infty} y^n e^{-y} dy = \frac{1}{a^{n+1}} \int_0^{\infty} x^n e^{-x} dx.$$

$$\text{Hence} \quad \int_0^{\infty} x^n e^{-ax} dx = \frac{\Gamma(n+1)}{a^{n+1}}, \quad [1]$$

provided that  $a$  is positive and  $n > -1$ .

$$(b) \quad \int_0^1 x^m \left(\log \frac{1}{x}\right)^n dx. \quad (\text{v. Art. 84, Ex. 4.})$$

$$\text{Let} \quad y = -\log x.$$

$$\text{then} \quad \int_0^1 x^m \left(\log \frac{1}{x}\right)^n dx = \int_0^{\infty} y^n e^{-(m+1)y} dy.$$

Hence, by [1],

$$\int_0^1 x^m \left(\log \frac{1}{x}\right)^n dx = \frac{\Gamma(n+1)}{(m+1)^{n+1}}, \quad [2]$$

if  $m > -1$  and  $n > -1$ .

$$(c) \quad \int_0^{\infty} e^{-x^2} dx.$$

$$\text{Let} \quad y = x^2;$$

$$\text{then} \quad \int_0^{\infty} e^{-x^2} dx = \frac{1}{2} \int_0^{\infty} \frac{e^{-y}}{\sqrt{y}} dy = \frac{1}{2} \int_0^{\infty} x^{-\frac{1}{2}} e^{-x} dx.$$

$$\text{Hence} \quad \int_0^{\infty} e^{-x^2} dx = \frac{1}{2} \Gamma\left(\frac{1}{2}\right). \quad [3]$$

$$98. \quad \int_0^1 x^{m-1} (1-x)^{n-1} dx = B(m, n) \quad [1]$$

is an exceedingly important integral that can be expressed in terms of Gamma Functions; it is known as the *Beta Function*, or the *First Eulerian Integral*,  $\Gamma(n)$  being sometimes called the *Second Eulerian Integral*.

In the Beta Function,  $m$  and  $n$  are *positive*, and  $B(m, n)$  is always finite and determinate. (v. Art. 84, Ex. 5.)

In  $\int_0^1 x^{m-1} (1-x)^{n-1} dx$  let  $y = 1-x$ ,

and we get

$$\int_0^1 x^{m-1} (1-x)^{n-1} dx = \int_0^1 y^{n-1} (1-y)^{m-1} dy,$$

or

$$B(m, n) = B(n, m). \quad [2]$$

In  $\int_0^1 x^{m-1} (1-x)^{n-1} dx$  let  $x = \frac{y}{1+y}$ ,

and we get

$$\int_0^1 x^{m-1} (1-x)^{n-1} dx = \int_0^\infty \frac{y^{m-1} dy}{(1+y)^{m+n}} = \int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} dx.$$

Hence  $\int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} dx = B(m, n). \quad [3]$

We have seen in [1] Art. 97 (a) that

$$\int_0^\infty x^n e^{-ax} dx = \frac{\Gamma(n+1)}{a^{n+1}}.$$

Hence  $\Gamma(m) = \int_0^\infty a^m x^{m-1} e^{-ax} dx,$

$$\Gamma(m) a^{n-1} e^{-a} = \int_0^\infty a^{m+n-1} x^{m-1} e^{-a(1+x)} dx,$$

$$\Gamma(m) \int_0^\infty a^{n-1} e^{-a} da = \int_0^\infty x^{m-1} \left( \int_0^\infty a^{m+n-1} e^{-a(1+x)} da \right) dx,$$

$$\Gamma(m) \Gamma(n) = \int_0^\infty x^{m-1} \frac{\Gamma(m+n)}{(1+x)^{m+n}} dx.$$

Therefore 
$$\frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)} = \int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} \cdot dx; \quad [4]$$

or by [9],

$$B(m, n) = \int_0^1 x^{m-1}(1-x)^{n-1} dx = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}. \quad [5]$$

If  $n = 1 - m$ , then since  $\Gamma(1) = 1$ ,

$$\int_0^1 \frac{x^{m-1}}{(1-x)^m} dx = \int_0^\infty \frac{x^{m-1}}{1+x} dx = \Gamma(m)\Gamma(1-m). \quad [6]$$

Formula [6] leads to an interesting confirmation of Art. 92 (b).

Let  $m = \frac{1}{2}$ , and we have from [6]

$$[\Gamma(\frac{1}{2})]^2 = \int_0^\infty \frac{dx}{(1+x)\sqrt{x}}.$$

Substitute

$$y = \sqrt{x},$$

and we have 
$$\int_0^\infty \frac{dx}{(1+x)\sqrt{x}} = 2 \int_0^\infty \frac{dy}{1+y^2} = \pi.$$

Hence

$$\Gamma(\frac{1}{2}) = \sqrt{\pi}; \quad [7]$$

and since by Art. 97 (c)

$$\begin{aligned} \int_0^\infty e^{-x^2} dx &= \frac{1}{2} \Gamma(\frac{1}{2}), \\ \int_0^\infty e^{-x^2} dx &= \frac{1}{2} \sqrt{\pi}. \end{aligned}$$

99. By the aid of formulas [4], [5], and [7] of Art. 98 a number of important integrals can be obtained.

For example, let us consider

$$\int_0^{\frac{\pi}{2}} \sin^n x \cdot dx, \quad \text{where } n \text{ is greater than } -1.$$

Let

$$y = \sin x,$$

and we have 
$$\int_0^{\frac{\pi}{2}} \sin^n x \cdot dx = \int_0^1 y^n (1-y^2)^{-\frac{1}{2}} dy.$$

Let, now,  $z = y^2$ ,

and

$$\begin{aligned} \int_0^1 y^n (1 - y^2)^{-\frac{1}{2}} dy &= \frac{1}{2} \int_0^1 z^{\frac{n}{2}-\frac{1}{2}} (1 - z)^{-\frac{1}{2}} dz \\ &= \frac{1}{2} \int_0^1 x^{\frac{n+1}{2}-1} (1 - x)^{\frac{1}{2}-1} dx = \frac{1}{2} B\left(\frac{n+1}{2}, \frac{1}{2}\right). \end{aligned}$$

But

$$B\left(\frac{n+1}{2}, \frac{1}{2}\right) = \frac{\Gamma\left(\frac{n+1}{2}\right)\Gamma\left(\frac{1}{2}\right)}{\Gamma\left(\frac{n+1}{2} + \frac{1}{2}\right)} \quad \text{by [5] Art. 98.}$$

$$= \sqrt{\pi} \cdot \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2} + 1\right)} \quad \text{by [7] Art. 98.}$$

Hence

$$\int_0^{\frac{\pi}{2}} \sin^n x \cdot dx = \frac{\sqrt{\pi}}{2} \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2} + 1\right)} \quad [1]$$

If  $n$  is a whole number, this will reduce to the result given in Art. 87, Ex. 5.

EXAMPLES.

$$(1) \int_0^1 \frac{x^{2n} dx}{\sqrt{1-x^2}} = \frac{\sqrt{\pi}}{2} \frac{\Gamma\left(n + \frac{1}{2}\right)}{\Gamma(n+1)}.$$

$$(2) \int_0^{\frac{\pi}{2}} \sin^m x \cos^n x \cdot dx = \frac{\Gamma\left(\frac{m+1}{2}\right)\Gamma\left(\frac{n+1}{2}\right)}{2\Gamma\left(\frac{m+n}{2} + 1\right)}.$$

$$(3) \int_0^1 \frac{dx}{\sqrt{1-x^n}} = \frac{\sqrt{\pi}}{n} \frac{\Gamma\left(\frac{1}{n}\right)}{\Gamma\left(\frac{1}{n} + \frac{1}{2}\right)}.$$

$$(4) \int_0^1 x^p (1-x^n)^m dx = \frac{1}{n} \frac{\Gamma(p+1) \Gamma\left(\frac{m+1}{n}\right)}{\Gamma\left(p+1 + \frac{m+1}{n}\right)}.$$

## CHAPTER IX.

## LENGTHS OF CURVES.

100. If we use rectangular coördinates, we have seen (I. Art. 27) that

$$\tan \tau = \frac{dy}{dx}, \quad [1]$$

and (I. Arts. 52 and 181) that

$$ds^2 = dx^2 + dy^2. \quad [2]$$

From these we get  $\sin \tau = \frac{dy}{ds},$  [3]

$$\cos \tau = \frac{dx}{ds}, \quad [4]$$

by the aid of a little elementary Trigonometry.

These formulas are of great importance in dealing with all properties of curves that concern in any way the lengths of arcs.

We have already considered the use of [2] in the first volume of the Calculus, and we have worked several examples by its aid in rectification of curves. Before going on to more of the same sort we shall find it worth while to obtain the equations of two very interesting transcendental curves, the *catenary* and the *tractrix*.

*The Catenary.*


101. The common *catenary* is the curve in which a uniform heavy flexible string hangs when its ends are supported.

As the string is flexible, the only force exerted by one portion of the string on an adjacent portion is a pull along the string, which we shall call the tension of the string, and shall represent by  $T$ .  $T$  of course has different values at different points of the string, and is some function of the coördinates of the point in question.



The tension at any point has to support the weight of the portion of the string below the point, and a certain amount of side pull, due to the fact that the string would hang vertically were it not that its ends are forcibly held apart.

Let the origin be taken at the lowest point of the curve, and suppose the string fastened at that point.



Let  $s$  be the arc  $OP$ ,  $P$  being any point of the string. As the string is uniform, the weight of  $OP$  is proportional to its length; we shall call this weight  $ms$ .

This weight acts vertically downward, and must be balanced by the vertical effect of  $T$ , which, by I. Art. 112, is  $T \sin \tau$ .

$$\text{Hence} \quad T \sin \tau = ms. \quad (1)$$

As there is no *external* horizontal force acting, the horizontal effect of the tension at one end of any portion of the string must be the same as the horizontal effect at the other end. In other words,

$$T \cos \tau = c \quad (2)$$

where  $c$  is a constant. Dividing (1) by (2) we get

$$s = \frac{c}{m} \tan \tau,$$

$$\text{or} \quad s = a \tan \tau, \quad (3)$$

where  $a$  is some constant. From this we want to get an equation in terms of  $x$  and  $y$ .

$$\tan \tau = \sqrt{\sec^2 \tau - 1} = \sqrt{\frac{ds^2}{dx^2} - 1};$$

$$\text{hence} \quad s^2 = a^2 \left( \frac{ds^2}{dx^2} - 1 \right),$$

$$\text{or} \quad a^2 ds^2 = (a^2 + s^2) dx^2,$$

$$\text{and} \quad \frac{ads}{(a^2 + s^2)^{\frac{1}{2}}} = dx. \quad \text{Integrate both members.}$$

$$a \log(s + \sqrt{a^2 + s^2}) = x + C;$$

when  $x = 0, s = 0,$

hence

$$C = a \log a,$$

and

$$\log(s + \sqrt{a^2 + s^2}) = \frac{x}{a} + \log a,$$

$$s + \sqrt{a^2 + s^2} = ae^{\frac{x}{a}},$$

$$\sqrt{a^2 + s^2} = ae^{\frac{x}{a}} - s,$$

$$a^2 = a^2 e^{\frac{2x}{a}} - 2ae^{\frac{x}{a}}s,$$

$$s = \frac{a}{2}(e^{\frac{2x}{a}} - e^{-\frac{2x}{a}}) = a \tan \tau \quad \text{by (3).}$$

Hence

$$a \frac{dy}{dx} = \frac{a}{2}(e^{\frac{2x}{a}} - e^{-\frac{2x}{a}}),$$

and

$$y = \frac{a}{2}(e^{\frac{2x}{a}} + e^{-\frac{2x}{a}}) + C.$$

If we change our axes, taking the origin at a point  $a$  units below the lowest point of the curve,  $y = a$  when  $x = 0,$  and therefore  $C = 0,$  and we get, as the equation of the catenary,

$$y = \frac{a}{2}(e^{\frac{2x}{a}} + e^{-\frac{2x}{a}}). \tag{4}$$

EXAMPLE.

Find the curve in which the cables of a suspension-bridge must hang.

Ans. A parabola.

*The Tractrix.*


102. If two particles are attached to a string, and rest on a rough horizontal plane, and one, starting with the string stretched, moves in a straight line at right angles with the initial position of the string, dragging the other particle after it, the path of the second particle is called the *tractrix*.

Take as the axis of  $X$  the path of the first particle, and as the axis of  $Y$  the initial position of the string, and let  $a$  be

the length of the string. From the nature of the curve the string is always a tangent, and we shall have for any point  $P$

$$\frac{y}{a} = -\sin \tau, \quad [1]$$

for  $\tau$  lying in the fourth quadrant has a negative sine.



$$\frac{y^2}{a^2} = \sin^2 \tau = \frac{dy^2}{ds^2} = \frac{dy^2}{dx^2 + dy^2};$$

hence

$$y^2(dx^2 + dy^2) = a^2 dy^2,$$

$$y^2 dx^2 = (a^2 - y^2) dy^2,$$

and

$$dx = \pm \frac{(a^2 - y^2)^{\frac{1}{2}} dy}{y}$$

is the differential equation of the *tractrix*.

On the right-hand half of the curve  $\tau$  is in the fourth quadrant,  $\frac{dy}{dx}$  or  $\tan \tau$  is negative, and we shall write the equation

$$dx = -\frac{(a^2 - y^2)^{\frac{1}{2}} dy}{y}. \quad [2]$$

If we allow the radical to be ambiguous in sign we shall get also the curve that would be described if the first particle went to the left instead of to the right. The *tractrix* curve, generally considered, includes these two portions.

Integrating both members of [2], and determining the arbitrary constant, we get

$$x = -\sqrt{a^2 - y^2} + a \log \frac{a + \sqrt{a^2 - y^2}}{y} \quad [3]$$

as the equation of the *tractrix*.

EXAMPLES.

(1) Show by Art. 102 (1) that in the tractrix  $s = a \log \frac{a}{y}$  if  $s$  is measured from the starting-point.

(2) Find the evolute of the tractrix. (I. Art. 93.)

*Rectification of Curves.*

103. In finding the length of an arc of a given curve we can regard it as the limit of the sum of the differentials of the arc, and express it by a definite integral.

We shall have 
$$s = \int_{x=x_0}^{x=x_1} \sqrt{dx^2 + dy^2}.$$

Of course in using this formula we must express  $\sqrt{dx^2 + dy^2}$  in terms of  $x$  only, or of  $y$  only, or of some single variable on which  $x$  and  $y$  depend, before we can integrate.

For example; let us find the length of an arc of the circle

$$x^2 + y^2 = a^2.$$

$$2x \cdot dx + 2y \cdot dy = 0,$$

$$dy = -\frac{x \cdot dx}{y},$$

$$dx^2 + dy^2 = \frac{x^2 + y^2}{y^2} dx^2 = \frac{a^2}{y^2} dx^2 = \frac{a^2}{a^2 - x^2} dx^2,$$

$$s = a \int_{x_0}^{x_1} \frac{dx}{\sqrt{a^2 - x^2}} = a \left( \sin^{-1} \frac{x_1}{a} - \sin^{-1} \frac{x_0}{a} \right).$$

The length of a quadrant  $= a \int_0^a \frac{dx}{\sqrt{a^2 - x^2}} = \frac{\pi a}{2};$

$\therefore$  the length of a circumference  $= 2\pi a.$

*Length of Arc of Cycloid.*

104. For the cycloid we have

$$\left. \begin{aligned} x &= a\theta - a \sin \theta \\ y &= a - a \cos \theta \end{aligned} \right\} \quad (\text{I. Art. 99.})$$

$$dx = a(1 - \cos \theta) d\theta = y d\theta,$$

$$\theta = \text{vers}^{-1} \frac{y}{a},$$

$$d\theta = \frac{1}{a} \cdot \frac{dy}{\sqrt{2\frac{y}{a} - \frac{y^2}{a^2}}} = \frac{dy}{\sqrt{2ay - y^2}},$$

$$dx = \frac{y dy}{\sqrt{2ay - y^2}},$$

$$ds^2 = dx^2 + dy^2 = \frac{2ay dy^2}{2ay - y^2} = \frac{2ady^2}{2a - y},$$

$$ds = \sqrt{2a} \cdot \frac{dy}{\sqrt{2a - y}},$$

$$s = \sqrt{2a} \int_{y_0}^{y_1} \frac{dy}{\sqrt{2a - y}} = 2\sqrt{2a} (\sqrt{2a - y_0} - \sqrt{2a - y_1}).$$

If the arc is measured from the cusp,  $y_0 = 0$ ,

$$s = 4a - 2\sqrt{2a}\sqrt{2a - y_1}. \quad [1]$$

If the arc is measured to the highest point,  $y_1 = 2a$ ,

$$s = 4a.$$

The whole arch =  $8a$ .**EXAMPLE.**Taking the origin at the vertex, and taking the direction downward as the positive direction for  $y$ , the equations become

$$\left. \begin{aligned} x &= a\theta + a \sin \theta \\ y &= a - a \cos \theta \end{aligned} \right\} \quad (\text{I. Art. 100.})$$

Show that  $s = 2\sqrt{2ay}$  when the arc is measured from the summit of the curve.

105. We can rectify the cycloid without eliminating  $\theta$ .

$$\left. \begin{aligned} x &= a\theta - a \sin \theta \\ y &= a - a \cos \theta \end{aligned} \right\},$$

$$dx = a(1 - \cos \theta) d\theta,$$

$$dy = a \sin \theta d\theta,$$

$$dx^2 + dy^2 = 2a^2 d\theta^2 (1 - \cos \theta),$$

and  $s = a \sqrt{2} \int_{\theta_0}^{\theta_1} (1 - \cos \theta)^{\frac{1}{2}} d\theta,$

$$s = a \sqrt{2} \int_{\theta_0}^{\theta_1} \left[ 2 \sin^2 \frac{\theta}{2} \right]^{\frac{1}{2}} d\theta = 4a \int_{\theta_0}^{\theta_1} \sin \frac{\theta}{2} d\frac{\theta}{2} = 4a \left( \cos \frac{\theta_0}{2} - \cos \frac{\theta_1}{2} \right).$$

If  $\theta_0 = 0$  and  $\theta_1 = 2\pi$ , we get  $s = 8a$  as the whole curve.

106. Let us find the length of an arch of the epicycloid.

$$\left. \begin{aligned} x &= (a + b) \cos \theta - b \cos \frac{a+b}{b} \theta \\ y &= (a + b) \sin \theta - b \sin \frac{a+b}{b} \theta \end{aligned} \right\}, \text{(I. Art. 109 [1].)}$$

$$dx = \left[ -(a + b) \sin \theta + (a + b) \sin \frac{a+b}{b} \theta \right] d\theta,$$

$$dy = \left[ (a + b) \cos \theta - (a + b) \cos \frac{a+b}{b} \theta \right] d\theta.$$

$$ds^2 = (a + b)^2 d\theta^2 \left[ 2 - 2 \left( \cos \frac{a+b}{b} \theta \cos \theta + \sin \frac{a+b}{b} \theta \sin \theta \right) \right]$$

$$= 2(a + b)^2 d\theta^2 \left( 1 - \cos \frac{a}{b} \theta \right).$$

$$s = (a + b) \sqrt{2} \int_{\theta_0}^{\theta_1} \left( 1 - \cos \frac{a}{b} \theta \right)^{\frac{1}{2}} d\theta,$$

$$s = \frac{4b(a + b)}{a} \left[ \cos \frac{a}{2b} \theta_0 - \cos \frac{a}{2b} \theta_1 \right]. \quad [1]$$

To get a complete arch we must let  $\theta_0 = 0$  and  $\theta_1 = \frac{2b}{a} \pi$ .  
Hence, for a whole arch,

$$s = \frac{8b(a + b)}{a}.$$

## EXAMPLES.

(1) Find the length of an arch of a hypocycloid.

$$\text{Ans. } s = \frac{8b(a-b)}{a}.$$

(2) Find the length of an arch of the curve  $x^{\frac{1}{2}} + y^{\frac{1}{2}} = a^{\frac{1}{2}}$ , and show that it agrees with the result of Ex. 1.

(v. I. Art. 109, Ex. 2.)

107. Let us attempt to find the length of an arc of the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

We have

$$\frac{2x dx}{a^2} + \frac{2y dy}{b^2} = 0,$$

$$dy = -\frac{b^2 x}{a^2 y} dx,$$

$$ds^2 = \frac{b^4 x^2 + a^4 y^2}{a^4 y^2} dx^2 = \frac{a^2 - \frac{a^2 - b^2}{a^2} x^2}{a^2 - x^2} dx^2 = \frac{a^2 - e^2 x^2}{a^2 - x^2} dx^2,$$

where  $e$  is the eccentricity of the ellipse.

$$\text{Hence } s = \int_{x_0}^{x_1} \left[ \frac{a^2 - e^2 x^2}{a^2 - x^2} \right]^{\frac{1}{2}} dx. \quad [1]$$

The length of the elliptic quadrant is

$$s_q = \int_0^a \left[ \frac{a^2 - e^2 x^2}{a^2 - x^2} \right]^{\frac{1}{2}} dx. \quad [2]$$

These integrals cannot be obtained directly, but

$$\left[ \frac{a^2 - e^2 x^2}{a^2 - x^2} \right]^{\frac{1}{2}} = \left[ 1 + \frac{(1 - e^2)x^2}{a^2 - x^2} \right]^{\frac{1}{2}}$$

can be expanded by the Binomial Theorem, and the terms of the result can be integrated separately, and we shall have the required length expressed by a series.

A more convenient way of dealing with the problem is to use an auxiliary angle. Instead of  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  we can use the pair of equations

$$\left. \begin{aligned} x &= a \sin \phi \\ y &= b \cos \phi \end{aligned} \right\}, \quad (\text{I. Art. 150}),$$

$$\begin{aligned} dx &= a \cos \phi \cdot d\phi, \\ dy &= -b \sin \phi \cdot d\phi, \\ ds^2 &= (a^2 \cos^2 \phi + b^2 \sin^2 \phi) d\phi^2 = [a^2 - (a^2 - b^2) \sin^2 \phi] d\phi^2 \\ &= a^2 \left( 1 - \frac{a^2 - b^2}{a^2} \sin^2 \phi \right) d\phi^2 = a^2 (1 - e^2 \sin^2 \phi) d\phi^2, \end{aligned}$$

where  $e$  is the eccentricity of the ellipse.

$$\begin{aligned} s &= a \int_{\phi_0}^{\phi_1} (1 - e^2 \sin^2 \phi)^{\frac{1}{2}} d\phi \tag{3} \\ &= a \int_{\phi_0}^{\phi_1} [1 - \frac{1}{2} e^2 \sin^2 \phi - \frac{1}{2} \cdot \frac{1}{4} e^4 \sin^4 \phi - \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{3}{8} e^6 \sin^6 \phi \dots] d\phi. \end{aligned}$$

For the arc of a quadrant we have

$$s_q = a \int_0^{\frac{\pi}{2}} [1 - e^2 \sin^2 \phi]^{\frac{1}{2}} d\phi. \tag{4}$$

EXAMPLES.

(1) Obtain  $s_q$  as a series from [2], and also from [4], and compare the results with Art. 91, Ex. 5.

(2) Show that the length of an arc of the hyperbola is

$$s = b \int_{\phi_0}^{\phi_1} \left[ 1 + \frac{a^2 e^2}{b^2} \sinh^2 \phi \right]^{\frac{1}{2}} d\phi.$$

*Polar Formulæ.*

108. If we use polar coördinates we have

$$\begin{aligned} ds &= \sqrt{dr^2 + r^2 d\phi^2}, \quad (\text{I. Art. 207, Ex. 2.}) \\ \tan \epsilon &= \frac{r d\phi}{dr}, \quad (\text{I. Art. 207.}) \end{aligned}$$

From these we get, by Trigonometry,

$$\sin \epsilon = \frac{r d\phi}{ds}, \quad \cos \epsilon = \frac{dr}{ds}.$$

109. Let us find the equation of the curve which crosses all its radii vectores at the same angle. Here

$$\tan \epsilon = \alpha, \quad \text{a constant,} \quad \frac{r d\phi}{dr} = \alpha, \quad \frac{adr}{r} = d\phi,$$



$$a \log r = \phi + C, \quad r = e^{\frac{\phi}{a} + \frac{c}{a}} = e^{\frac{\phi}{a}} e^{\frac{c}{a}},$$

$$r = be^{\frac{\phi}{a}}. \quad (1)$$

where  $b$  is some constant depending upon the position of the origin. This curve is known as the *Logarithmic* or *Equiangular Spiral*.

110. To rectify the *Logarithmic Spiral*. We have, from 109 (1),

$$\frac{\phi}{a} = \log \frac{r}{b};$$

$$d\phi = a \frac{dr}{r},$$

$$rd\phi = adr,$$

$$ds^2 = dr^2 + r^2 d\phi^2 = (1 + a^2) dr^2;$$

$$s = \int_{r_0}^{r_1} (1 + a^2)^{\frac{1}{2}} dr = (1 + a^2)^{\frac{1}{2}} (r_1 - r_0).$$

#### EXAMPLES.

(1) Find the length of an arc of the parabola from its polar equation

$$r = \frac{m}{1 + \cos \phi}.$$

(2) Find the length of an arc of the Spiral of Archimedes

$$r = a\phi.$$

111. To rectify the *Cardioid*. We have

$$r = 2a(1 - \cos \phi), \quad (\text{I. Art. 109, Ex. 1}).$$

$$dr = 2a \sin \phi \cdot d\phi,$$

$$ds^2 = 4a^2 \sin^2 \phi \cdot d\phi^2 + 4a^2(1 - \cos \phi)^2 d\phi^2$$

$$= 8a^2 d\phi^2 (1 - \cos \phi),$$

$$s = 2\sqrt{2} \cdot a \int_{\phi_0}^{\phi_1} (1 - \cos \phi)^{\frac{1}{2}} d\phi = 8a \left[ \cos \frac{\phi_0}{2} - \cos \frac{\phi_1}{2} \right]$$

$$= 16a \text{ for the whole perimeter.}$$

*Involutes.*

112. If we can express the length of the arc of a given curve, measured from a fixed point, in terms of the coördinates of its variable extremity, we can find the equation of the *involute* of the curve.

We have found the equations of the evolute of  $y = fx$  in the form

$$\left. \begin{aligned} x' &= x - \rho \cos \nu \\ y' &= y - \rho \sin \nu \end{aligned} \right\}, \quad (\text{I. Art. 91}).$$

We have proved that  $\tan \nu = \tan \tau'$ , (I. Art. 95),

and that  $\frac{ds'}{d\rho} = 1$ , (I. Art. 96);

$$\left. \begin{aligned} \sin \tau' &= \frac{dy'}{ds'} \\ \cos \tau' &= \frac{dx'}{ds'} \end{aligned} \right\}, \quad (\text{Art. 100}).$$

Since  $\tan \nu = \tan \tau'$ ,  $\nu = \tau'$  or  $\nu = 180^\circ + \tau'$ .

As normal and radius of curvature have opposite directions, we shall consider  $\nu = 180^\circ + \tau'$ .

Then  $\sin \nu = -\sin \tau'$  and  $\cos \nu = -\cos \tau'$ .

$$\text{Hence} \quad x' = x + \rho \frac{dx'}{ds'}, \quad (1)$$

$$y' = y + \rho \frac{dy'}{ds'}. \quad (2)$$

$$\begin{aligned} \text{Since} \quad d\rho &= ds', \\ \rho &= s' + l \end{aligned} \quad (3)$$

where  $l$  is an arbitrary constant. Since  $x$  and  $y$  are the coördinates of any point of the *involute*, it is only necessary to eliminate  $x'$ ,  $y'$ , and  $\rho$  by combining equations (1), (2), and (3) with the equation of the evolute.

As we are supposed to start with the equation of the evolute and work towards the equation of the involute, it will be more natural to accent the letters belonging to the latter curve instead

of those going with the former; and our equations may be written

$$x = x' + \rho' \frac{dx}{ds}; \quad y = y' + \rho' \frac{dy}{ds}; \quad \rho' = s + l. \quad (4)$$

Since  $\rho' = l$  when  $s = 0$ , it follows that  $l$  is the free portion of the string with which we start. (I. Art. 97.) By varying  $l$  we may get different involutes of the same curve.

To test our method, let us find the involute of the curve

$$y^2 = \frac{8}{27m} (x - m)^3, \quad (5)$$

for which  $l = m$ . We must first find  $s$ .

$$2y dy = \frac{8}{9m} (x - m)^2 dx,$$

$$dy = \frac{4}{9m} \frac{(x - m)^2}{y} dx,$$

$$ds^2 = \frac{2x + m}{3m} dx^2,$$

$$s = \frac{1}{\sqrt{3m}} \int_m^x (2x + m)^{\frac{1}{2}} dx = \frac{1}{3\sqrt{3m}} (2x + m)^{\frac{3}{2}} - m,$$

$$\rho' = s + m = \frac{1}{3\sqrt{3m}} (2x + m)^{\frac{3}{2}},$$

$$x = x' + \frac{2x + m}{3},$$

$$y = y' + \frac{4}{27m} \frac{(2x + m)(x - m)^2}{y},$$

$$x' = \frac{x - m}{3},$$

$$y' = -\frac{4}{9y} (x - m)^2 = -\frac{4x'^2}{y},$$

$$x = 3x' + m,$$

$$y = -\frac{4x'^2}{y'}.$$

Substituting in (5) the values of  $x$  and  $y$  just obtained, we have

$$y'^2 = 2mx'$$

as the equations of the required involute.

EXAMPLE.

Find an involute of  $ay^2 = x^2$ .

113. An involute of the cycloid is easily found. Take equations I. Art. 100 (C).

$$\left. \begin{aligned} x &= a\theta + a \sin \theta \\ y &= -a + a \cos \theta \end{aligned} \right\}$$

Let

$$\rho' = s,$$

$$dx = a(1 + \cos \theta) d\theta = 2a \cos^2 \frac{\theta}{2} d\theta,$$

$$dy = -a \sin \theta d\theta = -2a \sin \frac{\theta}{2} \cos \frac{\theta}{2} d\theta,$$

$$ds^2 = 2a^2 d\theta^2 (1 + \cos \theta) = 4a^2 d\theta^2 \cos^2 \frac{\theta}{2},$$

$$s = 2a \int_0^\theta \cos \frac{\theta}{2} d\theta = 4a \sin \frac{\theta}{2},$$

$$x = x' + 4a \sin \frac{\theta}{2} \cos \frac{\theta}{2} = x' + 2a \sin \theta,$$

$$y = y' - 4a \sin^2 \frac{\theta}{2} = y' - 2a(1 - \cos \theta),$$

$$\left. \begin{aligned} x' &= a\theta - a \sin \theta \\ y' &= a - a \cos \theta \end{aligned} \right\}$$

a cycloid with its cusp at the summit of the given cycloid.

EXAMPLE.

From the equations of a circle

$$\left. \begin{aligned} x &= a \cos \phi \\ y &= a \sin \phi \end{aligned} \right\}$$

obtain the equations of the involute of the circle. Let  $l = 0$ .

$$\text{Ans. } \left. \begin{aligned} x' &= a(\cos \phi + \phi \sin \phi) \\ y' &= a(\sin \phi - \phi \cos \phi) \end{aligned} \right\}$$

*Intrinsic Equation of a Curve.*

114. An equation connecting the length of the arc, measured from a fixed point of any curve to a variable point, with the angle between the tangent at the fixed point and the tangent at the variable point, is the *intrinsic equation* of the curve. If the fixed point is the origin and the fixed tangent the axis of  $X$ , the variables in the *intrinsic equation* are  $s$  and  $\tau$ .

We have already such an equation for the catenary

$$s = a \tan \tau, \quad \text{Art. 101 (3), [1]}$$

the origin being the lowest point of the curve.

The intrinsic equation of a circle is obviously

$$s = a\tau, \quad [2]$$

whatever origin we may take.

The intrinsic equation of the tractrix is easily obtained. We have

$$y = -a \sin \tau, \quad \text{Art. 102 (1),}$$

and 
$$s = a \log \frac{a}{y}; \quad \text{Art. 102, Ex. 1.}$$

hence 
$$s = a \log(-\csc \tau)$$

where  $\tau$  is measured from the axis of  $X$ , and  $s$  is measured from the point where the curve crosses the axis of  $Y$ . As the curve is tangent to the axis of  $Y$ , we must replace  $\tau$  by  $\tau - 90^\circ$ , and we get

$$s = a \log \sec \tau \quad [3]$$

as the intrinsic equation of the tractrix.

## EXAMPLE.

Show that the intrinsic equation of an inverted cycloid, when the vertex is origin, is

$$s = 4a \sin \tau; \quad (1)$$

when the cusp is origin, is

$$s = 4a(1 - \cos \tau). \quad (2)$$

115. To find the intrinsic equation of the epicycloid we can use the results obtained in Art. 106.

$$dx = (a+b) \left( \sin \frac{a+b}{b} \theta - \sin \theta \right) d\theta = 2(a+b) \cos \frac{a+2b}{2b} \theta \sin \frac{a}{2b} \theta \cdot d\theta,$$

$$dy = (a+b) \left( \cos \theta - \cos \frac{a+b}{b} \theta \right) d\theta = 2(a+b) \sin \frac{a+2b}{2b} \theta \sin \frac{a}{2b} \theta \cdot d\theta,$$

by the formulas of Trigonometry

$$\sin \alpha - \sin \beta = 2 \cos \frac{1}{2}(\alpha + \beta) \sin \frac{1}{2}(\alpha - \beta),$$

$$\cos \beta - \cos \alpha = 2 \sin \frac{1}{2}(\alpha + \beta) \sin \frac{1}{2}(\alpha - \beta);$$

$$\tan \tau = \frac{dy}{dx} = \tan \frac{a+2b}{2b} \theta,$$

hence  $\tau = \frac{a+2b}{2b} \theta;$

$$s = \frac{4b(a+b)}{a} \left( 1 - \cos \frac{a}{2b} \theta \right) \text{ by Art. 106[1];}$$

therefore  $s = \frac{4b(a+b)}{a} \left( 1 - \cos \frac{a}{a+2b} \tau \right) \quad [1]$

is the intrinsic equation of the epicycloid, with the *cusp* as origin.

If we take the origin at a vertex instead of at a cusp

$$s = \frac{4b(a+b)}{a} + s',$$

$$\tau = \frac{\pi(a+2b)}{2a} + \tau';$$

hence  $s' = \frac{4b(a+b)}{a} \sin \frac{a}{a+2b} \tau';$

or  $s = \frac{4b(a+b)}{a} \sin \frac{a}{a+2b} \tau \quad [2]$

is the intrinsic equation of an epicycloid referred to a vertex.

## EXAMPLE.

Obtain the intrinsic equation of the hypocycloid in the forms

$$s = \frac{4b(a-b)}{a} \left( 1 - \cos \frac{a}{a-2b} \tau \right), \quad (1)$$

$$s = \frac{4b(a-b)}{a} \sin \frac{a}{a-2b} \tau. \quad (2)$$

116. The intrinsic equation of the Logarithmic Spiral is found without difficulty.

We have  $r = be^{\frac{\phi}{a}}$ , (Art. 109),

and  $s = \sqrt{1+a^2} (r_1 - r_0)$ . (Art. 110).

If we measure the arc from the point where the spiral crosses the initial line,  $r_0 = b$ , and we have

$$s = b\sqrt{1+a^2}(e^{\frac{s}{a}} - 1).$$

In polar coördinates  $\tau = \phi + \epsilon$ , and in this case  $\epsilon = \tan^{-1}a$ ; if we measure our angle from the tangent at the beginning of the arc we must subtract  $\epsilon$  from the value just given, and we have

$$s = b(\sqrt{1+a^2})(e^{\frac{\tau}{a}} - 1);$$

or, more briefly,  $s = k(e^{\tau} - 1)$ ,  $k$  and  $c$  being constants.

117. If we wish to get the intrinsic equation of a curve directly from the equation in rectangular coördinates, the following method will serve:

Let the axis of  $X$  be tangent to the curve at the point we take as origin.

$$\tan \tau = \frac{dy}{dx}; \quad (1)$$

and as the equation of the curve enables us to express  $y$  in terms of  $x$ , (1) will give us  $x$  in terms of  $\tau$ , say  $x = F\tau$ ;

then  $dx = F' \tau \cdot d\tau$ , divide by  $ds$ ;

$$\frac{dx}{ds} = F' \tau \frac{d\tau}{ds}, \text{ but } \frac{dx}{ds} = \cos \tau;$$

hence  $ds = \sec \tau F' \tau \cdot d\tau$ . (2)

Integrating both members we shall have the required intrinsic equation.

For example, let us take  $x^2 = 2my$ , which is tangent to the axis of  $X$  at the origin.

$$2x dx = 2m dy,$$

$$\frac{dy}{dx} = \tan \tau = \frac{x}{m},$$

$$dx = m \sec^2 \tau \cdot d\tau,$$

$$\frac{dx}{ds} = \cos \tau = m \sec^2 \tau \frac{d\tau}{ds},$$

$$ds = m \sec^3 \tau \cdot d\tau, \tag{1}$$

$$s = m \int \frac{d\tau}{\cos^3 \tau} = \frac{m}{2} \left[ \frac{\sin \tau}{\cos^2 \tau} + \log \tan \left( \frac{\pi}{4} + \frac{\tau}{2} \right) \right] + C,$$

$$s = 0 \text{ when } \tau = 0; \quad \therefore C = 0;$$

$$s = \frac{m}{2} \left[ \frac{\sin \tau}{\cos^2 \tau} + \log \tan \left( \frac{\pi}{4} + \frac{\tau}{2} \right) \right]. \tag{2}$$

#### EXAMPLES.


(1) Devise a method when the curve is tangent to the axis of  $Y$ , and apply it to  $y^2 = 2mx$ .

(2) Obtain the intrinsic equation of  $y^2 = \frac{8}{27m} (x - m)^3$ .

(3) Obtain the intrinsic equation of the involute of a circle. (Art. 113, Ex.)



118. The *evolute* or the *involute* of a curve is easily found from its intrinsic equation.



If the curvature of the given curve decreases as we pass along the curve,  $\rho$  increases, and

$$s' = \rho - \rho_0 \quad (\text{I. Art. 96}).$$

If the curvature increases,  $\rho$  decreases, and

$$s' = \rho_0 - \rho.$$

Hence always  $s' = \pm (\rho - \rho_0)$ ; [1]

$$\rho = \frac{ds}{d\tau}, \quad (\text{I. Arts. 86 and 90}).$$

We see from the figure that  $\tau' = \tau$ .

Hence  $s' = \pm \left[ \left( \frac{ds}{d\tau} \right)_{\tau=\tau'} - \left( \frac{ds}{d\tau} \right)_{\tau=0} \right]$ ,

or, as we shall write it for brevity,

$$s = \pm \left. \frac{ds}{d\tau} \right|_0^{\tau}. \quad [2]$$

119. The evolute of the *tractrix*  $s = a \log \sec \tau$  is

$$s = a \left. \frac{d \log \sec \tau}{d\tau} \right|_0^{\tau} = a \tan \tau, \quad \text{the catenary.}$$

The evolute of the circle  $s = a\tau$  is

$$s = a \left. \frac{d\tau}{d\tau} \right|_0^{\tau} = 0, \quad \text{a point.}$$

The evolute of the cycloid  $s = 4a(1 - \cos \tau)$  is

$$s = 4a \left. \frac{d(1 - \cos \tau)}{d\tau} \right|_0^\tau = 4a \sin \tau,$$

an equal cycloid, with its vertex at the origin.

EXAMPLES.

- (1) Prove that the evolute of the logarithmic spiral is an equal logarithmic spiral.
- (2) Find the evolute of a parabola.
- (3) Find the evolute of the catenary.

120. The evolute of an epicycloid is a similar epicycloid, with each vertex at a cusp of the given curve.

Take the equation

$$s = \frac{4b(a+b)}{a} \left( 1 - \cos \frac{a}{a+2b} \tau \right). \quad \text{Art. 115 [1].}$$

For the evolute,

$$s = \frac{4b(a+b)}{a} \left. \frac{d \left( 1 - \cos \frac{a}{a+2b} \tau \right)}{d\tau} \right|_0^\tau,$$

$$s = \frac{4b(a+b)}{a+2b} \sin \frac{a}{a+2b} \tau. \quad [1]$$

The form of [1] is that of an epicycloid referred to a vertex as origin; let us find  $a'$  and  $b'$ , the radii of the fixed and rolling circles.

$$s = \frac{4b'(a'+b')}{a'} \sin \frac{a'}{a'+2b'} \tau, \quad \text{by Art. 115 [2];}$$

hence, 
$$\frac{4b'(a'+b')}{a'} = \frac{4b(a+b)}{a+2b},$$

$$\frac{a'}{a'+2b'} = \frac{a}{a+2b}.$$

Solving these equations, we get

$$a' = \frac{a^2}{a + 2b}.$$

$$b' = \frac{ab}{a + 2b}.$$

$$\frac{a'}{b'} = \frac{a}{b}.$$

and the radii of the fixed and rolling circles have the same ratio in the evolute as in the original epicycloid: therefore the two curves are similar.

#### EXAMPLE.

Show that the evolute of a hypocycloid is a similar hypocycloid.

121. We have seen that in involute and evolute  $r$  has the same value: that is,  $r = r'$ .

If  $s'$  and  $r'$  refer to the evolute, and  $s$  and  $r$  to the involute, we have found that

$$s' = \frac{ds}{d\tau}.$$

or

$$s' = \frac{ds}{d\tau} - l, \quad l \text{ being a constant.}$$

the length of the radius of curvature at the origin.

$$(s' + l)d\tau' = ds.$$

$$s = \int (s' + l)d\tau'$$

is the equation of the involute.

The involute of the catenary  $s = a \tan \tau$  is, when  $l = 0$ ,

$$s = a \int \tan \tau d\tau = a \log \sec \tau, \quad \text{the tractrix.}$$

The involute of the cycloid  $s = 4a \sin \tau$  when  $l = 0$  is

$$s = 4a \int_0^\tau \sin \tau . d\tau = 4a(1 - \cos \tau),$$

an equal cycloid referred to its cusp as origin.

The involute of a cycloid referred to its cusp  $s = 4a(1 - \cos \tau)$  when  $l = 0$  is

$$s = 4a \int_0^\tau (1 - \cos \tau) d\tau = 4a(\tau + \sin \tau),$$

a curve we have not studied.

The involute of a circle  $s = a\tau$  when  $l = 0$  is

$$s = a \int_0^\tau \tau . d\tau = \frac{a\tau^2}{2}.$$

122. While any given curve has but one evolute, it has an infinite number of involutes, since the equation of the involute

$$s' = \int_0^\tau (s + l) d\tau$$

contains an arbitrary constant  $l$ ; and the nature of the involute will in general be different for different values of  $l$ .

If we form the involute of a given curve, taking a particular value for  $l$ , and form the involute of this involute, taking the same value of  $l$ , and so on indefinitely, the curves obtained will continually approach the logarithmic spiral.

Let (1)  
 $s = f\tau$

be the given curve.

$$s = \int_0^\tau (l + f\tau) d\tau = l\tau + \int_0^\tau f\tau . d\tau$$

is the first involute ;

$$s = \int_0^\tau (l + l\tau + \int_0^\tau f\tau . d\tau) d\tau = l\tau + \frac{l\tau^2}{2} + \int_0^\tau \int_0^\tau f\tau . d\tau^2$$

is the second involute ;

$$s = l\tau + \frac{l\tau^2}{2} + \frac{l\tau^3}{3!} + \dots + \frac{l\tau^n}{n!} + \int_0^\tau \int_0^\tau \dots \int_0^\tau f\tau . d\tau^n \tag{2}$$

is the  $n$ th involute.

By Maclaurin's Theorem,

$$f\tau = f_0 + \tau f'_0 + \frac{\tau^2}{2!} f''_0 + \frac{\tau^3}{3!} f'''_0 + \dots$$

But  $s = 0$  when  $\tau = 0$ ; hence  $f_0 = 0$ , and

$$\begin{aligned} f\tau &= A_1\tau + \frac{A_2}{2!}\tau^2 + \frac{A_3}{3!}\tau^3 + \dots, \\ \int_0^\tau f\tau.d\tau &= \frac{A_1}{2}\tau^2 + \frac{A_2}{3!}\tau^3 + \frac{A_3}{4!}\tau^4 + \dots, \\ \int_0^{\tau^2} f\tau.d\tau^2 &= \frac{A_1}{3!}\tau^3 + \frac{A_2}{4!}\tau^4 + \frac{A_3}{5!}\tau^5 + \dots, \\ \int_0^{\tau^n} f\tau.d\tau^n &= \frac{A_1\tau^{n+1}}{(n+1)!} + \frac{A_2\tau^{n+2}}{(n+2)!} + \frac{A_3\tau^{n+3}}{(n+3)!} + \dots; \quad (3) \end{aligned}$$

as  $n$  increases indefinitely all the terms of (3) approach zero (I. Art. 133), and the limiting form of (2) is

$$\begin{aligned} s &= l\tau + \frac{l\tau^2}{2!} + \frac{l\tau^3}{3!} + \dots \\ &= l\left(1 + \frac{\tau}{1} + \frac{\tau^2}{2!} + \frac{\tau^3}{3!} + \dots - 1\right), \\ s &= l(e^\tau - 1) \qquad \text{by I. Art. 133 [2],} \end{aligned}$$

which is a logarithmic spiral.

123. The equation of a curve in rectangular coördinates is readily obtained from the intrinsic equation.

Given  $s = f\tau$ ,

we know that  $\sin \tau = \frac{dy}{ds}$ ,

and  $\cos \tau = \frac{dx}{ds}$ ;

hence 
$$\left. \begin{aligned} ds &= \cos \tau ds = \cos \tau f' \tau.d\tau, \\ dy &= \sin \tau ds = \sin \tau f' \tau.d\tau, \\ s &= \int \cos \tau f' \tau.d\tau \\ y &= \int \sin \tau f' \tau.d\tau \end{aligned} \right\}$$

The elimination of  $\tau$  between these equations will give us the equation of the curve in terms of  $x$  and  $y$ . Let us apply this method to the catenary.

$$s = a \tan \tau,$$

$$ds = a \sec^2 \tau d\tau,$$

$$x = a \int_0^\tau \sec \tau d\tau = a \log \sqrt{\frac{1 + \sin \tau}{1 - \sin \tau}},$$

$$y = a \int_0^\tau \sec \tau \tan \tau d\tau = a(\sec \tau - 1),$$

$$e^{\frac{2x}{a}} = \frac{1 + \sin \tau}{1 - \sin \tau},$$

$$\sin \tau = \frac{e^{\frac{2x}{a}} - 1}{e^{\frac{2x}{a}} + 1} = \frac{e^{\frac{x}{a}} - e^{-\frac{x}{a}}}{e^{\frac{x}{a}} + e^{-\frac{x}{a}}},$$

$$\sec \tau = \frac{1}{2}(e^{\frac{x}{a}} + e^{-\frac{x}{a}}),$$

$$y = \frac{a}{2}(e^{\frac{x}{a}} + e^{-\frac{x}{a}}) - a,$$

the equation of the catenary referred to its lowest point as origin.

#### *Curves in Space.*

124. The length of the arc of a curve of double curvature is the limit of the sum of the chords of smaller arcs into which the given arc may be broken up, as the number of these smaller arcs is indefinitely increased. Let  $(x, y, z)$ ,  $(x + dx, y + \Delta y, z + \Delta z)$  be the coördinates of the extremities of any one of the small arcs in question;  $dx, \Delta y, \Delta z$  are infinitesimal;  $\sqrt{dx^2 + \Delta y^2 + \Delta z^2}$  is the length of the chord of the arc. In dealing with the limit of the sum of these chords, any one may be replaced by a quantity differing from it by infinitesimals of higher order than the first.  $\sqrt{dx^2 + dy^2 + dz^2}$  is such a value;

hence 
$$s = \int_{x=x_0}^{x=x_1} \sqrt{dx^2 + dy^2 + dz^2}.$$

Let us rectify the helix.

$$\left. \begin{aligned} x &= a \cos \theta \\ y &= a \sin \theta \\ z &= k\theta \end{aligned} \right\}. \quad (\text{I. Art. 214.})$$

$$dx = -a \sin \theta \cdot d\theta,$$

$$dy = a \cos \theta \cdot d\theta,$$

$$dz = k d\theta,$$

$$ds^2 = (a^2 + k^2) d\theta^2,$$

$$s = (a^2 + k^2)^{\frac{1}{2}} \int_{\theta_0}^{\theta_1} d\theta = \sqrt{a^2 + k^2} (\theta_1 - \theta_0).$$

EXAMPLES.

(1) Find the length of the curve  $\left( y = \frac{x^2}{2a}, z = \frac{x^3}{6a^2} \right)$ .

*Ans.*  $s = x + z + l$ .

(2)  $y = 2\sqrt{ax} - x, z = x - \frac{2}{3}\sqrt{\frac{x^3}{a}}$ . *Ans.*  $s = x + y - z + l$ .


## CHAPTER X.

### AREAS.

125. We have found and used a formula for the area bounded by a given curve, the axis of  $X$ , and a pair of ordinates.

$$A = \int y dx.$$

We can readily get this formula as a definite integral. The area in the figure is the sum of the slices into which it is divided by the ordinates; if  $\Delta x$ , the base of each slice, is indefinitely decreased, the slice is infinitesimal. The area of any slice differs from  $y\Delta x$  by less than  $\Delta y\Delta x$ , which is of the second order if  $\Delta x$  is the principal infinitesimal. We have then



$$A = \lim_{\Delta x \doteq 0} \sum_{x_0}^{x_1} y \Delta x \quad \text{by I. Art. 161.}$$

Hence 
$$A = \int_{x_0}^{x_1} y dx. \quad [1]$$

If the curve in question lies above the axis of  $X$ , and  $x_0$  is less than  $x_1$ , each ordinate is positive, each  $\Delta x$  is positive, each term of the sum whose limit is required is positive, the sum is positive, and the limit of the sum or the area sought is positive. If, however, the curve lies below the axis of  $X$ , and  $x_0$  is less than  $x_1$ , each ordinate is negative, each  $\Delta x$  is positive, each term of the sum is negative, the sum is negative, and the limit



of the sum or the area sought is negative. If, then, the curve happens to cross the axis between  $x_0$  and  $x_1$ , formula [1] gives us the difference between the portion of the area above the axis of  $X$  and the portion below the axis of  $X$ , but throws no light upon the magnitudes of the separate portions. Consequently, in any actual geometrical problem it is usually necessary to find the portion of the required area above the axis of  $X$  and the portion below the axis of  $X$  separately; and for this purpose it is essential to know at what points the curve crosses the axis. Indeed, if the problem is in the least complicated, it is necessary to begin by carefully tracing the given curve from its equation, and then to keep its form and position in mind during the whole process of solution.

#### EXAMPLES.

(1) Show that  $\int_{x_0}^{x_1} x dy$  is the area bounded by a curve, the axis of  $Y$ , and perpendiculars let fall from the ends of the bounding arc upon the axis of  $Y$ .

(2) If the axes are inclined at the angle  $\omega$ , show that these formulas become


$$A = \sin \omega \int_{x_0}^{x_1} y dx = \sin \omega \int_{y_0}^{y_1} x dy.$$

(3) Find the area bounded by the axis of  $X$ , the curve  $x^2 + 4y = 0$ , and the ordinate of the point corresponding to the abscissa 4. Ans.  $5\frac{1}{3}$ .

(4) Find the area bounded by the axis of  $X$ , the curve  $y = x^3$ , and the ordinates corresponding to the abscissæ  $-2$  and  $2$ . Ans. 8.

(5) Find the area bounded by the axis of  $X$ , the axis of  $Y$ , the curve  $y = \cos x$ , and the ordinate corresponding to the abscissa  $3\pi$ . Ans. 6.

126. In polar coördinates we can regard the area between two radii vectores and the curve as the limit of the sum of sectors.



The area in question is the sum of the smaller sectorial areas, any one of which differs from  $\frac{1}{2}r^2\Delta\phi$  by less than the difference between the two circular sectors  $\frac{1}{2}(r + \Delta r)^2\Delta\phi$  and  $\frac{1}{2}r^2\Delta\phi$ ; that is, by less than  $r\Delta r\Delta\phi + \frac{(\Delta r)^2\Delta\phi}{2}$ , which is of the

second order if  $\Delta\phi$  is the principal infinitesimal.

Hence 
$$A = \lim_{\Delta\phi \rightarrow 0} \left[ \frac{1}{2} \sum_{\phi=\phi_0}^{\phi=\phi_1} r^2 \Delta\phi \right],$$

$$A = \frac{1}{2} \int_{\phi_0}^{\phi_1} r^2 d\phi.$$

127. Let us find the area between the catenary, the axis of X, the axis of Y, and any ordinate.

$$A = \int_0^x y dx = \frac{a}{2} \int_0^x (e^{\frac{x}{a}} + e^{-\frac{x}{a}}) dx,$$

$$A = \frac{a^2}{2} (e^{\frac{x}{a}} - e^{-\frac{x}{a}}),$$

but  $\frac{a}{2} (e^{\frac{x}{a}} - e^{-\frac{x}{a}}) = s,$  by Art. 101.

Hence  $A = as,$

and the area in question is the length of the arc multiplied by the distance of the lowest point of the curve from the origin.

128. Let us find the area between the tractrix and the axis of X.

We have  $dx = -\frac{dy}{y} \sqrt{a^2 - y^2}.$  (Art. 102.)

$$A = \int y dx = - \int dy \sqrt{a^2 - y^2}.$$

The area in question is

$$A = - \int_a^0 dy \sqrt{a^2 - y^2} = \frac{\pi a^2}{4},$$

which is the area of the quadrant of a circle with  $a$  as radius.

EXAMPLE.

Give, by the aid of infinitesimals, a geometric proof of the result just obtained for the tractrix.

129. In the last section we found the area between a curve and its asymptote, and obtained a finite result. Of course this means that, as our second bounding ordinate recedes from the origin, the area in question, instead of increasing indefinitely, approaches a finite limit, which is the area obtained. Whether the area between a curve and its asymptote is finite or infinite will depend upon the nature of the curve.

Let us find the area between an hyperbola and its asymptote.

The equation of the hyperbola referred to its asymptotes as axes is

$$xy = \frac{a^2 + b^2}{4}.$$

Let  $\omega$  be the angle between the asymptotes; then

$$A = \sin \omega \int_0^\infty y dx = \frac{a^2 + b^2}{4} \sin \omega \int_0^\infty \frac{dx}{x} = \infty.$$


Take the curve  $y^2 x = 4 a^2 (2 a - x)$ ,

or  $y^2 = 4 a^2 \cdot \frac{2 a - x}{x}$ ;

any value of  $x$  will give two values of  $y$  equal with opposite signs; therefore the axis of  $x$  is an axis of symmetry of the curve.

When  $x = 2 a$ ,  $y = 0$ ; as  $x$  decreases,  $y$  increases; and when  $x = 0$ ,  $y = \infty$ . If  $x$  is negative, or greater than  $2 a$ ,  $y$  is imaginary. The shape of the curve is something like that in the

figure, the axis of  $Y$  being an asymptote. The area between the curve and the asymptote is then either



$$A = 2 \int_0^{2a} y dx \quad \text{or} \quad A = 2 \int_0^{\infty} x dy;$$

by the first formula,

$$A = 4a \int_0^{2a} \sqrt{\frac{2a-x}{x}} \cdot dx = 4a^2 \pi;$$

by the second,

$$A = 16a^3 \int_0^{\infty} \frac{d\eta}{y^2 + 4a^2} = 4a^2 \pi.$$

EXAMPLES.

(1) Find the area between the curve  $y^2(x^2 + a^2) = a^2 x^2$  and its asymptote  $y = a$ . Ans.  $A = 2a^2$ .

(2) Find the area between  $y^2(2a-x) = x^3$  and its asymptote  $x = 2a$ . Ans.  $A = 3\pi a^2$ .

(3) Find the area bounded by the curve  $y^2 = \frac{x^2(a+x)}{a-x}$  and its asymptote  $x = a$ . Ans.  $A = 2a^2 \left(1 + \frac{\pi}{4}\right)$ .

130. If the coördinates of the points of a curve are expressed in terms of an auxiliary variable, no new difficulty is presented.

Take the case of the circle  $x^2 + y^2 = a^2$ , which may be written

$$\left. \begin{aligned} x &= a \cos \phi \\ y &= a \sin \phi \end{aligned} \right\};$$

$$dy = a \cos \phi d\phi.$$

The whole area  $A = a^2 \int_0^{2\pi} \cos^2 \phi d\phi = \pi a^2$ .

## EXAMPLES.

- (1) The whole area of an ellipse  $\left. \begin{array}{l} x = a \cos \phi \\ y = b \sin \phi \end{array} \right\}$  is  $\pi ab$ .
- (2) The area of an arch of the cycloid is  $3\pi a^2$ .
- (3) The area of an arch of the companion to the cycloid  $x = a\theta$ ,  $y = a(1 - \cos \theta)$  is  $2\pi a^2$ .

131. If we wish to find the area between two curves, or the area bounded by a closed curve, the altitude of our elementary rectangle is the difference between the two values of  $y$ , which correspond to a single value of  $x$ . If the area between two curves is required, we must find the abscissas of their points of intersection, and they will be our limits of integration; if the whole area bounded by a closed curve is required, we must find the values of  $x$  belonging to the points of contact of tangents parallel to the axis of  $Y$ .

Let us find the whole area of the curve

$$a^4 y^2 + b^2 x^4 = a^2 b^2 x^2,$$

or 
$$a^4 y^2 = b^2 x^2 (a^2 - x^2).$$

The curve is symmetrical with reference to the axis of  $X$ , and passes through the origin. It consists of two loops whose areas must be found separately. Let us find where the tangents are parallel to the axis of  $Y$ .

$$y = \frac{b}{a^2} x \sqrt{a^2 - x^2},$$

$$\frac{dy}{dx} = \frac{b}{a^2} \cdot \frac{a^2 - 2x^2}{\sqrt{a^2 - x^2}} = \tan \tau.$$

$\tau = \frac{\pi}{2}$  when  $\tan \tau = \infty$ , that is, when  $x = \pm a$ .

$$A = 2 \frac{b}{a^2} \int_{-a}^a x \sqrt{a^2 - x^2} dx + 2 \frac{b}{a^2} \int_0^a x \sqrt{a^2 - x^2} dx = \frac{4}{3} ab.$$

Again; find the whole area of  $(y-x)^2 = a^2 - x^2$ .

$$y = x \pm \sqrt{a^2 - x^2},$$

$$A = \int (y' - y'') dx = \int 2\sqrt{a^2 - x^2} \cdot dx.$$

To find the limits of integration, we must see where  $r = \frac{\pi}{2}$ .

$$\frac{dy}{dx} = \frac{\sqrt{a^2 - x^2} \mp x}{\sqrt{a^2 - x^2}} = \infty \text{ when } x = \pm a.$$

$$A = 2 \int_{-a}^a \sqrt{a^2 - x^2} \cdot dx = \pi a^2.$$

EXAMPLES.

(1) Find the area of the loop of the curve  $y^2 = \frac{x^2(a+x)}{a-x}$ .

$$\text{Ans. } 2a^2 \left(1 - \frac{\pi}{4}\right).$$

(2) Find the area between the curves  $y^2 - 4ax = 0$  and  $x^2 - 4ay = 0$ .

$$\text{Ans. } \frac{16a^2}{3}.$$

(3) Find the whole area of the curve  $x^3 + y^3 = a^3$ .  $\text{Ans. } \frac{8}{3}\pi a^2$ .

(4) Find the area of a loop of  $a^2 y^4 = x^4(a^2 - x^2)$ .  $\text{Ans. } \frac{4a^2}{5}$ .

(5) Find the whole area of the curve

$$2y^2(a^2 + x^2) - 4ay(a^2 - x^2) + (a^2 - x^2)^2 = 0.$$

$$\text{Ans. } a^2\pi \left(4 - \frac{5\sqrt{2}}{2}\right).$$

132. We have seen that in polar coordinates

$$A = \frac{1}{2} \int_{\phi_0}^{\phi_1} r^2 d\phi.$$

Let us try one or two examples.

(a) To find the whole area of a circle.

The polar equation is  $r = a$ .

$$A = \frac{1}{2} \int_0^{2\pi} a^2 d\phi = \pi a^2.$$

(b) To find the area of the cardioid  $r = 2a(1 - \cos \phi)$ .

$$A = \frac{1}{2} \int_0^{2\pi} 4a^2(1 - \cos \phi)^2 d\phi = 2a^2 \int_0^{2\pi} (1 - 2\cos \phi + \cos^2 \phi) d\phi,$$

$$A = 6a^2\pi.$$

(c) To find the area between an arch of the epicycloid and the circumference of the fixed circle.

$$\left. \begin{aligned} x &= (a+b)\cos \theta - b \cos \frac{a+b}{b} \theta \\ y &= (a+b)\sin \theta - b \sin \frac{a+b}{b} \theta \end{aligned} \right\}.$$

We can get the area bounded by two radii vectores and the arch in question, and subtract the area of the corresponding sector of the fixed circle.

Changing to polar coördinates,

$$x = r \cos \phi,$$

$$y = r \sin \phi.$$

We want  $\frac{1}{2} \int r^2 d\phi$ .

$$\tan \phi = \frac{y}{x},$$

$$\sec^2 \phi d\phi = \frac{x dy - y dx}{x^2};$$

but, since

$$x = r \cos \phi, \quad \sec \phi = \frac{r}{x};$$

hence

$$\frac{r^2 d\phi}{x^2} = \frac{x dy - y dx}{x^2},$$

and

$$r^2 d\phi = x dy - y dx;$$

$$dx = (a+b) \left( -\sin \theta + \sin \frac{a+b}{b} \theta \right) d\theta,$$

$$dy = (a+b) \left( \cos \theta - \cos \frac{a+b}{b} \theta \right) d\theta.$$

$$x dy - y dx = (a+b)(a+2b) \left( 1 - \cos \frac{a}{b} \theta \right) d\theta = r^2 d\phi.$$

Our limits of integration are obviously 0 and  $\frac{2b\pi}{a}$ .

Hence 
$$A = \frac{1}{2}(a+b)(a+2b) \int_0^{2\pi} \left(1 - \cos \frac{a}{b} \theta\right) d\theta,$$

$$A = \frac{b\pi}{a} (a+b)(a+2b),$$

is the area of the sector of the epicycloid. Subtract the area of the circular sector  $\pi ab$ , and we get

$$A = \frac{b^2(3a+2b)}{a} \pi$$

as the area in question.

(d) To find the area of a loop of the curve  $r^2 = a^2 \cos 2\phi$ .

For any value of  $\phi$  the values of  $r$  are equal with opposite signs. Hence the origin is a centre.

When  $\phi = 0$ ,  $r = \pm a$ ; as  $\phi$  increases,  $r$  decreases in length till  $\phi = \frac{\pi}{4}$ , when  $r = 0$ ; as soon as  $\phi > \frac{\pi}{4}$ ,  $r$  is imaginary. If  $\phi$  decreases from 0,  $r$  decreases in length until  $\phi = -\frac{\pi}{4}$ , when  $r = 0$ ; and when  $\phi < -\frac{\pi}{4}$ ,  $r$  is imaginary. To get the area of a loop, then, we must integrate from  $\phi = -\frac{\pi}{4}$  to  $\phi = \frac{\pi}{4}$ .

$$A = \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} r^2 d\phi = \frac{1}{2} a^2 \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \cos 2\phi \cdot d\phi = \frac{a^2}{2}.$$

EXAMPLES.

(1) Find the area of a sector of the parabola  $r = \frac{m}{1 + \cos \phi}$ .

(2) Find the area of a loop of the curve  $r^2 \cos \phi = a^2 \sin 3\phi$ .

*Ans.*  $\frac{3a^2}{4} - \frac{a^2}{2} \log 2.$

(3) Find the whole area of the curve  $r = a(\cos 2\phi + \sin 2\phi)$ .

*Ans.*  $\pi a^2.$

(4) Find the area of a loop of the curve  $r \cos \phi = a \cos 2\phi$ .

*Ans.*  $\left(2 - \frac{\pi}{2}\right) a^2.$

(5) Find the area between  $r = a(\sec \phi + \tan \phi)$  and its asymptote  $r \cos \phi = 2a$ .

*Ans.*  $\left(\frac{\pi}{2} + 2\right) a^2.$



133. When the equation of a curve is given in rectangular coördinates, we can often simplify the problem of finding its area by transforming to polar coördinates.

For example, let us find the area of

$$(x^2 + y^2)^2 = 4a^2x^2 + 4b^2y^2.$$

Transform to polar coördinates.

$$r^4 = 4r^2(a^2 \cos^2 \phi + b^2 \sin^2 \phi),$$

$$r^2 = 4(a^2 \cos^2 \phi + b^2 \sin^2 \phi),$$

$$A = 2 \int_0^{2\pi} (a^2 \cos^2 \phi + b^2 \sin^2 \phi) d\phi = 2\pi(a^2 + b^2).$$

#### EXAMPLES.

- (1) Find the area of a loop of the curve  $(x^2 + y^2)^3 = 4a^2x^2y^2$ .

$$\text{Ans. } \frac{\pi a^2}{8}.$$

- (2) Find the whole area of the curve  $\frac{x^2}{a^4} + \frac{y^2}{b^4} = \frac{1}{c^2} \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} \right)^2$ .

$$\text{Ans. } \frac{\pi c^2}{2ab} (a^2 + b^2).$$


- (3) Find the area of a loop of the curve  $y^3 - 3axy + x^3 = 0$ .

$$\text{Ans. } \frac{3a^2}{2}.$$

134. The area between a curve and its evolute can easily be found from the intrinsic equation of the curve.

It is easily seen that the area bounded by the radii of curvature at two points infinitely near, by the curve and by the evolute, differs from  $\frac{1}{2}\rho^2 d\tau$  by an infinitesimal of higher order. The area bounded by two given radii vectores, the curve and the evolute, is then

$$A = \frac{1}{2} \int_{\tau_0}^{\tau_1} \rho^2 d\tau.$$



$$\rho = \frac{ds}{d\tau}$$

Hence 
$$A = \frac{1}{2} \int_{\tau_0}^{\tau_1} \left( \frac{ds}{d\tau} \right)^2 d\tau.$$

For example, the area between a cycloid and its evolute is

$$\begin{aligned} A &= \frac{1}{2} \int_{\tau_0}^{\tau_1} \left( \frac{d(4a \sin \tau)}{d\tau} \right)^2 d\tau \\ &= 8a^2 \int_{\tau_0}^{\tau_1} \cos^2 \tau d\tau. \end{aligned}$$

Let  $\tau_0 = 0$  and  $\tau_1 = \frac{\pi}{2}$ ;


$$A = 8a^2 \int_0^{\frac{\pi}{2}} \cos^2 \tau d\tau = 2\pi a^2.$$

EXAMPLES.

- (1) Find the area between a circle and its evolute.
- (2) Find the area between the circle and its involute.

*Holditch's Theorem.*

135. If a line of fixed length move with its ends on any closed curve which is always concave toward it, the area between the curve and the locus of a given point of the moving line is equal to the area of an ellipse, of which the segments into which the line is divided by the given point are the semi-axes.




Let the figure represent the given curve, the locus of  $P$ , and the envelope of the moving line.

Let  $AP = a$  and  $PB = b$ , and let  $CB = \rho$ ,  $C$  being the point of contact of the moving line with its envelope. Let  $AB = a + b = c$ .

The area between the first curve and the second is the area between the first curve and the envelope, minus the area between the second curve and the envelope.

Let  $\theta$  be the angle which the moving line makes at any instant with some fixed direction. Let the figure represent two near positions of the moving line;  $\Delta\theta$ , the angle between these positions, being the principal infinitesimal.



$$PB = \rho, P'B' = \rho + \Delta\rho.$$

The area  $PBB'P'P$  differs from  $\frac{1}{2}\rho^2 d\theta$  by an infinitesimal of higher order than the first.

$\frac{1}{2}\rho^2 d\theta$  is the area of  $PBMP$ , and differs from  $PP'NB$  by less than the rectangle on  $PM$  and  $PQ$ , which is of higher order than the first, by I. Art. 153. But  $PP'NB$  differs from  $PP'B'B$  by less than the rectangle on  $BN$  and  $NB'$ , which is of higher order than the first, since  $NB'$ , which is less than  $PP' + \Delta\rho$ , is infinitesimal and  $\Delta\theta$  is infinitesimal.

The area between the first curve and the envelope is then  $\frac{1}{2} \int_0^{2\pi} \rho^2 d\theta$ ; or, since we can take  $PP'A'A$  just as well for our elementary area,  $\frac{1}{2} \int_0^{2\pi} (c - \rho)^2 d\theta$ .

$$\text{Hence} \quad \frac{1}{2} \int_0^{2\pi} \rho^2 d\theta = \frac{1}{2} \int_0^{2\pi} (c - \rho)^2 d\theta;$$

$$\text{whence} \quad 2c \int_0^{2\pi} \rho d\theta = 2c^2 \pi,$$

$$\text{or} \quad \int_0^{2\pi} \rho d\theta = \pi c. \tag{1}$$

The area between the second curve and the envelope is  $\frac{1}{2} \int_0^{2\pi} (\rho - \lambda)^2 d\theta$ .

The area between the first curve and the second is then

$$\begin{aligned}
 A &= \frac{1}{2} \int_0^{2\pi} \rho^2 d\theta - \frac{1}{2} \int_0^{2\pi} (\rho - b)^2 d\theta \\
 &= b \int_0^{2\pi} \rho d\theta - b^2 \pi \\
 &= \pi bc - b^2 \pi && \text{by (1),} \\
 &= \pi b(a + b) - b^2 \pi, \\
 A &= \pi ab, && (2)
 \end{aligned}$$

which is the area of an ellipse of which  $a$  and  $b$  are semi-axes.

Q. E. D.

EXAMPLES.


(1) If a line of fixed length move with its extremities on two lines at right angles with each other, the area of the locus of a given point of the line is that of an ellipse on the segments of the line as semi-axes.

(2) The result of (1) holds even when the fixed lines are not perpendicular.

*Areas by Double Integration.*

136. If we take  $x$  and  $y$  as the coördinates of any point  $P$  within our area,  $x$  and  $y$  will be independent variables, and

we can find the area bounded by two given curves,  $y = fx$  and  $y = Fx$ , by a double integration. Suppose the area in question divided into slices by lines drawn parallel to the axis of  $Y$ , and these slices subdivided into parallelograms by lines drawn parallel to the axis of  $X$ . The area of any one of the small parallelograms is  $\Delta y \Delta x$ . If we keep  $x$  constant, and take the sum



of these rectangles from  $y = fx$  to  $y = Fx$ , we shall get a result differing from the area of the corresponding slice by less than

$2 \Delta x \Delta y$ , which is infinitesimal of the second order if  $\Delta x$  and  $\Delta y$  are of the first order.

Hence 
$$\int_x^{x_1} \Delta x \cdot dy = \Delta x \int_x^{x_1} dy$$

is the area of the slice in question. If now we take the limit of the sum of all these slices, choosing our initial and final values of  $x$ , so that we shall include the whole area, we shall get the area required.

Hence 
$$A = \int_{x_0}^{x_1} \left( \int_x^{x_1} dy \right) dx.$$

In writing a double integral, the parentheses are usually omitted for the sake of conciseness, and this formula is given as

$$A = \int_{x_0}^{x_1} \int_x^{x_1} dy dx,$$


the order in which the integrations are to be performed being the same as if the parentheses were actually written.

If we begin by keeping  $y$  constant, and integrating with respect to  $x$ , we shall get the area of a slice formed by lines parallel to the axis of  $X$ , and we shall have to take the limit of the sum of these slices varying  $y$  in such a way as to include the whole area desired. In that case we should use the formula

$$A = \int_y^{y_1} \int_{f^{-1}y}^{F^{-1}y} dx dy.$$

137. For example, let us find the area bounded by the parabolas  $y^2 = 4ax$  and  $x^2 = 4ay$ .

The parabolas intersect at the origin and at the point  $(4a, 4a)$ .



$$A = \int_0^{4a} \int_{\frac{x^2}{4a}}^{\sqrt{4ax}} dy dx, \text{ or } A = \int_0^{4a} \int_{\frac{y^2}{4a}}^{\sqrt{4ay}} dx dy;$$

$$\int_{\frac{x^2}{4a}}^{\sqrt{4ax}} dy = \sqrt{4ax} - \frac{x^2}{4a};$$

$$\int_0^{4a} \int_{\frac{x^2}{4a}}^{\sqrt{4ax}} dy dx = \int_0^{4a} \left( \sqrt{4ax} - \frac{x^2}{4a} \right) dx = \frac{16}{3} a^2.$$

The second formula gives the same result.

EXAMPLES.

(1) Find the area of a rectangle by double integration; of a parallelogram; of a triangle.


(2) Find the area between the parabola  $y^2 = ax$  and the circle  $y^2 = 2ax - x^2$ .

*Ans.*  $2\left(\frac{\pi a^2}{4} - \frac{2a^2}{3}\right)$ .

(3) Find the whole area of the curve  $(y - mx - c)^2 = a^2 - x^2$ .

*Ans.*  $\pi a^2$ .

138. If we use polar coördinates we can still find our areas by double integration.



Let  $r = f\phi$  and  $r = F\phi$  be two curves. Divide the area between them into slices by drawing radii vectores; then subdivide these slices by drawing arcs of circles, with the origin as centre.

Let  $P$ , with coördinates  $r$  and  $\phi$ , be any point within the space whose area is sought. The curvilinear rectangle at  $P$  has the base  $r\Delta\phi$  and the altitude  $\Delta r$ ; its area differs from  $r\Delta\phi\Delta r$  by an infinitesimal of higher order than  $r\Delta\phi\Delta r$ .

The area of any slice as  $ab'a'b'$  is  $\int_{f\phi}^{F\phi} r\Delta\phi dr$ ,  $\phi$  and  $\Delta\phi$  being constant, that is  $\Delta\phi \int_{f\phi}^{F\phi} r dr$ . The whole area, the limit of the sum of such slices is  $A = \int_{\phi_0}^{\phi_1} \int_{f\phi}^{F\phi} r dr d\phi$ . (1)

Or we may first sum our rectangles, keeping  $r$  unchanged, and we get as the area of  $ef'e'f'$

$$r\Delta r \int_{f^{-1}\phi}^{F^{-1}\phi} d\phi, \text{ and } A = \int_{r_0}^{r_1} \int_{f^{-1}\phi}^{F^{-1}\phi} r d\phi dr. \quad (2)$$

It must be kept in mind that  $r$  in (1) and (2) is the radius vector of any point within the area sought, and not of a point on the boundary.

For example, the area between two concentric circles,  $r = a$  and  $r = b$ , is

$$A = \int_b^a \int_0^{2\pi} r d\phi dr = \int_0^{2\pi} \int_b^a r dr d\phi = \pi(a^2 - b^2).$$


Again, let us find the area between two tangent circles and a diameter through the point of contact.

Let  $a$  and  $b$  be the two radii,

$$r = 2a \cos \phi \quad (1)$$

and  $r = 2b \cos \phi \quad (2)$

are the equations of the two circles.



$$A = \int_0^{\frac{\pi}{2}} \int_{2b \cos \phi}^{2a \cos \phi} r dr d\phi = 2(a^2 - b^2) \int_0^{\frac{\pi}{2}} \cos^2 \phi d\phi = \frac{\pi}{2}(a^2 - b^2).$$

If we wish to reverse the order of our integrations we must break our area into two parts by an arc described from the origin as a centre, and with  $2b$  as a radius; then we have

$$\begin{aligned} A &= \int_0^{2b} \int_{\cos^{-1} \frac{r}{2b}}^{\cos^{-1} \frac{r}{2a}} r d\phi dr + \int_{2b}^{2a} \int_0^{\cos^{-1} \frac{r}{2a}} r d\phi dr \\ &= \int_0^{2b} r \left( \cos^{-1} \frac{r}{2a} - \cos^{-1} \frac{r}{2b} \right) dr + \int_{2b}^{2a} r \cos^{-1} \frac{r}{2a} dr \\ &= \frac{\pi}{2}(a^2 - b^2). \end{aligned}$$

#### EXAMPLE.


Find the area between the axis of  $X$  and two coils of the spiral  $r = a\phi$ .

CHAPTER XI.

AREAS OF SURFACES.

*Surfaces of Revolution.*

139. If a plane curve  $y = fx$  revolves about the axis of  $X$ , the area of the surface generated is the limit of the sum of the areas generated by the chords of the infinitesimal arcs into which the whole arc may be broken up. Each of these chords will generate the surface of the frustum of a cone of revolution if it revolves completely around the axis; and the area of the surface of a frustum of a cone of revolution is, by elementary Geometry, one-half the sum of the circumferences of the bases multiplied by the slant height. The frustum generated by the chord in the figure will have an area differing by infinitesimals of higher order from  $\pi(y + y + \Delta y)\Delta s$  or from  $2\pi y ds$ . The area generated by any given arc is then



$$S = 2\pi \int_{x_0}^{x_1} y ds. \tag{1}$$

If the arc revolves through an angle  $\theta$  instead of making a complete revolution, the surface generated is

$$S = \theta \int_{x_0}^{x_1} y ds. \tag{2}$$

It must be noted that [1] and [2] will give a positive value for  $S$  if the generating curve lies wholly above the axis of  $X$  at the start, and a negative value for  $S$  if it lies wholly below the axis of  $X$  at the start. If the curve happens to cross the axis of  $X$  between the points whose ordinates are  $y_0$  and  $y_1$ , [1] and [2] give not the area of the surface generated by the curve in question, but the difference between the areas generated by the




portion originally above the axis, and the portion originally below the axis.

EXAMPLE.

Show that if the arc revolves about the axis of  $Y$ ,  $S = 2\pi \int_{x_0}^{x_1} x ds$ .

140. To find the area of a cylinder of revolution.

Take the axis of the cylinder as the axis of  $X$ . Let  $a$  be the altitude and  $b$  the radius of the base of the cylinder. The equation of the revolving line is



$$y = b;$$

$$dy = 0,$$


$$ds = \sqrt{dx^2 + dy^2} = dx;$$

$$S = 2\pi \int_0^a y dx = 2\pi ab,$$

or the product of the altitude by the circumference of the base.

Again, let us find the surface of a zone.

The equation of the generating circle is



$$x^2 + y^2 = a^2;$$

$$ds = \frac{adx}{y};$$

$$S = 2\pi \int_{x_0}^{x_1} adx = 2a\pi(x_1 - x_0).$$

If  $x_0 = -a$  and  $x_1 = a$ ,  $S = 4a^2\pi$ .

Hence the surface of a zone is the altitude of the zone multiplied by the circumference of a great circle, and the surface of a sphere is equal to the areas of four great circles.

Again, take the surface generated by the revolution of a cycloid about its base.

$$\left. \begin{aligned} x &= a\theta - a \sin \theta \\ y &= a - a \cos \theta \end{aligned} \right\};$$

$$ds = a d\theta \sqrt{2(1 - \cos \theta)}, \quad \text{by Art. 105;}$$

$$S = 2\pi \int_0^{2\pi} a^2 \sqrt{2} \cdot (1 - \cos \theta)^{\frac{3}{2}} d\theta = \frac{64}{3} \pi a^2.$$

EXAMPLES.

(1) The area of the surface generated by the revolution of the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

about the axis of  $X$  is  $2\pi ab \left( \sqrt{1 - e^2} + \frac{\sin^{-1} e}{e} \right)$ ;

about the axis of  $Y$  is  $2\pi a^2 \left( 1 + \frac{1 - e^2}{2e} \log \frac{1 + e}{1 - e} \right)$ ,

where  $e^2 = \frac{a^2 - b^2}{a^2}$ .

(2) Find the area of the surface generated by the revolution of the catenary about the axis of  $X$ ; about the axis of  $Y$ .

(3) The whole surface generated by the revolution of the tractrix about its asymptote is  $4\pi a^2$ .


(4) The area generated by the revolution of a cycloid about its vertical axis is  $8\pi a^2 \left( \pi - \frac{4}{3} \right)$ .

(5) The area generated by the revolution of a cycloid about the tangent at its vertex is  $\frac{32}{3}\pi a^2$ .

(6) The area generated by the revolution of the curve  $x^3 + y^3 = a^3$  about its axis is  $\frac{16}{5}\pi a^2$ .

141. If we know the area generated by the revolution of a curve about any axis, we can get the area generated by the revolution about any parallel axis by an easy transformation of coördinates.

Given the surface generated by the arc from  $s_0$  to  $s_1$  about  $OX$ , to find the area generated by the same arc when it revolves about  $O'X'$ .



Let  $S$  be the surface about  $OX$ , and  $S'$  about  $O'X'$ .

We have

$$S = 2\pi \int_{s_0}^{s_1} y ds, \quad S' = 2\pi \int_{s_0}^{s_1} y' ds'.$$

By Anal. Geom.,  $x = x'$ ,  
 $y = y_0 + y'$ .

Hence  $dx = dx'$ ,  $dy = dy'$ ,  $ds = ds'$ .

$$\text{and } S = 2\pi \int_{s_0}^{s_1} (y_0 + y') ds = 2\pi y_0 (s_1 - s_0) + 2\pi \int_{s_0}^{s_1} y' ds,$$

$$= 2\pi y_0 (s_1 - s_0) + S'.$$

Therefore  $S' = S - 2\pi y_0 (s_1 - s_0)$ . [1]

$s_1 - s_0$  is the length of the revolving curve;  $2\pi y_0$  is the circumference of a circle of which  $y_0$  is the radius. Hence the new area is equal to the old area minus the area of a cylinder whose length is the length of the given arc and whose base is a circle of which the distance between the two lines is radius.

In using this principle careful attention must be paid to the sign of  $y_0$ , and it must be noted that the original formula  $S = 2\pi \int_{s_0}^{s_1} y ds$  will always give a negative value for the area of the surface generated, if the revolving arc starts from below the axis; and hence, that the surface generated by the revolution of any curve about an axis of symmetry will come out zero.

As an example of the use of the principle, let us find the surface of a ring.


Let  $a$  be the distance of the centre of the circle from the axis, and  $b$  the radius of the circle. Since the area generated by the revolution of the circle about a diameter is zero, the required area is

$$2\pi b \cdot 2\pi a = 4\pi^2 ab.$$

#### EXAMPLE.

Find the area of the ring generated by the revolution of a cycloid about any axis parallel to its base.

$$\text{Ans. } S = 4ab\pi \left( \pi + \frac{16a + 12b}{3b} \right).$$



142. If we use polar coördinates,

$$S = 2\pi \int_{s_0}^{s_1} y ds$$

becomes

$$S = 2\pi \int_{s_0}^{s_1} r \sin \phi \cdot ds.$$

where

$$ds = \sqrt{dr^2 + r^2 d\phi^2}.$$

For example; let us find the area of the surface generated by the revolution of the upper half of a cardioide about the horizontal axis.

$$r = 2a(1 - \cos \phi);$$

$$dr = 2a \sin \phi \cdot d\phi,$$

$$ds^2 = 8a^2(1 - \cos \phi)d\phi^2,$$

$$S = 2\pi \int_0^\pi \sqrt{4} \sqrt{2a^2(1 - \cos \phi)}^{\frac{1}{2}} \sin \phi \cdot d\phi.$$

$$S = \frac{16}{3} \pi a^2.$$

#### EXAMPLES.

- (1) Find the surface of a sphere from the polar equation.
- (2) Find the surface of a paraboloid of revolution from the polar equation of the parabola

$$r = \frac{m}{1 - \cos \phi}.$$

#### *Cylindrical Surfaces.*

143. If a cylindrical surface is generated by a line which is always parallel to the axis of  $Z$ , the area of the portion bounded by two positions of the generating line, the plane of  $XY$ , and any curve whose projection on the plane of  $XZ$  is given, is easily found.


Let  $ABCD$  be the cylindrical area required.

Let  $y = fx$  (1)

be the equation of  $AB$ , the line of intersection of the surface with the plane  $XY$ ; and let

$z = Fx$  (2)

be the equation of  $C_1D_1$ , the projection of  $CD$  on the plane of  $XZ$ .



If  $x, y, z$  are the coördinates of any point  $P$  of  $CD$ , the required area is evidently the limit of the sum of rectangles, of which  $PP'P''P'''$  is any one. The area of  $PP'P''P'''$  differs by an infinitesimal of higher order than  $ds$  from  $zds$ , and therefore the required area  $S = \int_{x_0}^{x_1} zds$ .

$x, z$  are the coördinates of  $P_1$ , and satisfy (2), and  $ds = \sqrt{dx^2 + dy^2}$  where  $x, y$  are the coördinates of  $P'$  and satisfy (1).

We have, then,  $S = \int_{x_0}^{x_1} z \sqrt{dx^2 + dy^2}$ . [3]

For example, let  $AB$  be the quadrant of a circle, and let the projection of the required area on the plane of  $XZ$  be the quadrant of an equal circle, so that the surface required is one-eighth of the surface of a *groin*.

Here  $x^2 + y^2 = a^2$ , (4)

and  $x^2 + z^2 = a^2$ ; (5)

$ds = \sqrt{dx^2 + dy^2} = \frac{a}{y} dx = \frac{adx}{\sqrt{a^2 - x^2}}$ ,

and  $z = \sqrt{a^2 - x^2}$ .

Therefore  $S = \int_0^a \sqrt{a^2 - x^2} \cdot \frac{adx}{\sqrt{a^2 - x^2}} = a \int_0^a dx = a^2.$

Again, let us find the area of the curved surface of the portion of a cylinder of revolution included within a spherical surface, whose centre lies on the surface of the cylinder, and whose radius is equal to the diameter of the cylinder.

If the centre of the sphere is taken as the origin, and a diametral plane of the cylinder as the plane of  $XZ$ , the surface required is four times that indicated in the figure.

The equation of the cylinder is

$$x^2 - ax + y^2 = 0, \tag{6}$$


and of the sphere

$$x^2 + y^2 + z^2 - a^2 = 0. \tag{7}$$

Subtract (6) from (7), and we get

$$z^2 + ax - a^2 = 0 \tag{8}$$

as the equation of a cylindrical surface perpendicular to the plane  $XZ$ , and passing through all the points of intersection of (6) and (7). (8) is, then, the equation of the projection on the plane of  $XZ$  of the line of intersection of the given spherical surface and the given cylindrical surface.



From (6),  $ds = \sqrt{dx^2 + dy^2} = \frac{a}{2y} dx = \frac{adx}{2\sqrt{ax - x^2}}.$

From (8),  $z = \sqrt{a^2 - ax}.$

Hence  $S = \int_0^a \sqrt{a^2 - ax} \cdot \frac{adx}{2\sqrt{ax - x^2}} = \frac{a\sqrt{a}}{2} \int_0^a \frac{\sqrt{a-x} \cdot dx}{\sqrt{x}\sqrt{a-x}}$   
 $= \frac{a\sqrt{a}}{2} \int_0^a \frac{dx}{\sqrt{x}} = a^2;$

and the whole area required,

$$4S = 4a^2.$$

## EXAMPLES.

(1) Find the area cut from the cylindrical surface whose base in the plane  $XY$  is a quadrant of the curve  $x^2 + y^2 = a^2$  by the plane  $x = z$ . *Ans.*  $\frac{2}{3}a^2$ .

(2) Find the area of that portion of a cylindrical surface whose base in the plane of  $XY$  is a quadrant of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , and whose projection on the plane of  $XZ$  is bounded by the curve  $a^2 z^2 = b^2 x^2 (a^2 - x^2)$ . *Ans.*  $S = \frac{ab(a^2 + ab + b^2)}{3(a + b)}$ .

(3) Let the base of the cylindrical surface be a tractrix, whose vertex lies at a distance  $a$  to the left of the origin, and whose asymptote is the axis of  $Y$ , while its projection on the plane of  $XZ$  is bounded by the parabola  $z^2 = -2mx$ .

$$\text{Ans. } S = 2a\sqrt{2ma}.$$

(4) Let the base of the cylindrical surface be the upper half of a cycloid, having its vertex at the origin and its base parallel to the axis of  $Y$ , and at a distance  $2a$  from the origin, while its projection on the plane of  $XZ$  is bounded by the parabola  $z^2 = 2mx$ .

$$\text{Ans. } S = 4a\sqrt{am}.$$

*Any Surface.*


144. Let  $x, y, z$  be the coördinates of any point  $P$  of the surface, and  $x + \Delta x, y + \Delta y, z + \Delta z$  the coördinates of a second point  $Q$  infinitely near the first. Draw planes through  $P$  and  $Q$  parallel to the planes of  $XY$  and  $YZ$ . These planes will intercept a curved quadrilateral  $PQ$  on the surface; its projection  $pq$ , a rectangle, on the plane of  $XZ$ ; and a parallelogram  $p'q'$  not shown in the figure, on the tangent plane at  $P$ , of which  $pq$  is the projection.  $PQ$  will differ from  $p'q'$  by an infinitesimal of higher order, and therefore our required surface will be the limit of the sum of the parallelograms of which  $p'q'$  is any one.

If  $\beta$  is the angle the tangent plane at  $P$  makes with  $XZ$ ,  $p'q' \cos \beta = pq$  or  $p'q' = pq \sec \beta = \Delta x \Delta z \sec \beta$ , and  $\sigma$ , our surface required, is equal to the double integral

$$\sigma = \iint \sec \beta dx dz$$

taken between limits so chosen as to embrace the whole surface.

The limit of the sum of the parallelograms, of which  $p'q'$  is a type, will be the required surface if the limit of the sum of the rectangles, of which  $pq$  is a type, is the projection of the surface in question on the plane of  $XZ$ ; so that the values of  $x$  and  $z$



between which we integrate in  $\sigma = \iint \sec \beta dx dz$  are precisely those we should use if we were finding the area of the projection of  $\sigma$  by the double integration  $\iint dx dz$ . (v. Art. 136.)

The equation of the tangent plane at  $P$  is  $(x - x_0)D_{x_0}f + (y - y_0)D_{y_0}f + (z - z_0)D_{z_0}f = 0$ , by I. Art. 217,  $(x_0, y_0, z_0)$  standing for the coördinates of the point of contact, and  $f(x, y, z) = 0$  being the equation of the surface.

The direction cosines of the perpendicular from the origin upon the plane are

$$\begin{aligned} \cos \alpha &= \frac{D_{x_0}f}{\sqrt{(D_{x_0}f)^2 + (D_{y_0}f)^2 + (D_{z_0}f)^2}}, \\ \cos \beta &= \frac{D_{y_0}f}{\sqrt{(D_{x_0}f)^2 + (D_{y_0}f)^2 + (D_{z_0}f)^2}}, \\ \cos \gamma &= \frac{D_{z_0}f}{\sqrt{(D_{x_0}f)^2 + (D_{y_0}f)^2 + (D_{z_0}f)^2}} \end{aligned}$$

by Anal. Geom. of Three Dimensions.



Hence, dropping the accents,

$$\sigma = \int \int \frac{\sqrt{(D_x f)^2 + (D_y f)^2 + (D_z f)^2}}{D_x f} dx dz. \quad [1]$$

By considering the projections upon the other coördinate planes we shall find

$$\sigma = \int \int \frac{\sqrt{(D_x f)^2 + (D_y f)^2 + (D_z f)^2}}{D_y f} dy dz; \quad [2]$$

$$\sigma = \int \int \frac{\sqrt{(D_x f)^2 + (D_y f)^2 + (D_z f)^2}}{D_z f} dx dy. \quad [3]$$

In each of the formulas the derivatives are partial derivatives.

Let us find the area of the portion of the surface of the sphere

$$x^2 + y^2 + z^2 = a^2$$

intercepted by the three coördinate planes.

$$D_x f = 2x,$$

$$D_y f = 2y,$$

$$D_z f = 2z,$$

$$\sqrt{(D_x f)^2 + (D_y f)^2 + (D_z f)^2} = 2a.$$

$$\sigma = \int_0^a \int_0^{\sqrt{a^2 - z^2}} \frac{a}{x} dy dz; \quad (1)$$

or

$$\sigma = \int_0^a \int_0^{\sqrt{a^2 - x^2}} \frac{a}{y} dz dx; \quad (2)$$

or

$$\sigma = \int_0^a \int_0^{\sqrt{a^2 - y^2}} \frac{a}{z} dx dy. \quad (3)$$

For, in the second one, which agrees best with the figure, we must take our limits so that the limit of the sum of the projections may be the quadrant in which the sphere is cut by the

plane  $XZ$ ; and the equation of this section is obtained by letting  $y = 0$  in the equation of the sphere, and is

$$x^2 + z^2 = a^2,$$

whence

$$z = \sqrt{a^2 - x^2}.$$

If we take as our limits in the integral  $\int \frac{a}{y} dz$  zero and  $\sqrt{a^2 - x^2}$  we shall get the area whose projection is a strip running from the axis of  $X$  to the curve; then, taking  $\int \left( \int \frac{a}{y} dz \right) dx$  from 0 to  $a$ , we shall get the area whose projection is the sum of all these strips, and that is our required surface.

$$y = \sqrt{a^2 - x^2 - z^2},$$

$$\sigma = a \int_0^a \int_0^{\sqrt{a^2 - x^2}} \frac{dz dx}{\sqrt{a^2 - x^2 - z^2}};$$

$$\int \frac{dz}{\sqrt{a^2 - x^2 - z^2}} = \sin^{-1} \frac{z}{\sqrt{a^2 - x^2}}$$


if we regard  $x$  as constant;

$$\int_0^{\sqrt{a^2 - x^2}} \frac{dz}{\sqrt{a^2 - x^2 - z^2}} = \frac{\pi}{2};$$

$$\sigma = a \int_0^a \frac{\pi}{2} dx = \frac{\pi a^2}{2},$$

the required area. Formulas (1) and (3) give the same result.

145. Suppose two cylinders of revolution drawn tangent to each other, and perpendicular to the plane of a great circle of a sphere, each having the radius of the great circle as a diameter; required the surface of the sphere not included by the cylinders.



The surface required is eight times the surface of which the shaded portion of the figure is the projection.

If we take the plane of the great circle as the plane of  $XY$ ,

$$x^2 - ax + y^2 = 0 \quad (1)$$

is the equation of the cylinder, and

$$x^2 + y^2 + z^2 = a^2 \quad (2)$$

of the sphere.

$$\text{We have } \sigma = \iint \frac{\sqrt{(D_x f)^2 + (D_y f)^2 + (D_z f)^2}}{D_x f} dy dx.$$

$$\text{From (2)} \quad D_x f = 2x,$$

$$D_y f = 2y,$$

$$D_z f = 2z;$$

$$(D_x f)^2 + (D_y f)^2 + (D_z f)^2 = 4a^2.$$

$$\text{Hence } \sigma = \int \int \frac{a}{z} dy dx = a \int \int \frac{dy dx}{\sqrt{a^2 - x^2 - y^2}}.$$

Our limits of integration for  $y$  are  $\sqrt{ax - x^2}$  and  $\sqrt{a^2 - x^2}$ ; for  $x$  are 0 and  $a$ .

$$\sigma = a \int_0^a \int_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} \frac{dy dx}{\sqrt{a^2-x^2-y^2}}.$$

$$\int_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} \frac{dy}{\sqrt{a^2-x^2-y^2}} = \sin^{-1} \frac{y}{\sqrt{a^2-x^2}} \Big|_{\sqrt{ax-x^2}}^{\sqrt{a^2-x^2}} = \frac{\pi}{2} - \sin^{-1} \sqrt{\frac{x}{a+x}}.$$

To find  $\int_0^a \sin^{-1} \sqrt{\frac{x}{a+x}} dx$  we must integrate by parts.

$$\text{Let} \quad u = \sin^{-1} \sqrt{\frac{x}{a+x}},$$

$$\text{and} \quad dv = dx;$$

$$v = x,$$

$$du = \frac{1}{2(a+x)} \sqrt{\frac{a}{x}} dx;$$

$$\int \sin^{-1} \sqrt{\frac{x}{a+x}} dx = x \sin^{-1} \sqrt{\frac{x}{a+x}} - \frac{\sqrt{a}}{2} \int \frac{\sqrt{x}}{a+x} dx.$$

Let  $w = \sqrt{x}$ ;  $2w dw = dx$

and  $\int \frac{\sqrt{x} \cdot dx}{a+x} = 2 \int \frac{w^2 dw}{a+w^2} = 2 \int \left(1 - \frac{a}{a+w^2}\right) dw.$

$$\int \frac{\sqrt{x} \cdot dx}{a+x} = 2 \left( w - \sqrt{a} \tan^{-1} \frac{w}{\sqrt{a}} \right),$$

$$\int_0^a \sin^{-1} \sqrt{\frac{x}{a+x}} \cdot dx$$

$$= a \sin^{-1} \frac{\sqrt{2}}{2} + a \tan^{-1} 1 - a = \frac{a\pi}{4} + \frac{a\pi}{4} - a = \frac{a\pi}{2} - a,$$

$$\sigma = a \left( \frac{a\pi}{2} - \frac{a\pi}{2} + a \right) = a^2.$$

$8\sigma = 8a^2$  is the whole surface in question.

146. Let us find the area of the curved surface of a right cone whose base is the curve  $x^2 + y^2 = a^2$ , and whose altitude is  $c$ .

If we take the vertex of the cone as the origin of coördinates, and its axis as the axis of  $Z$ , the equation of its curved surface is

$$x^2 + y^2 = \left(\frac{az}{c}\right)^2, \tag{1}$$

and the projection of the surface on the plane of  $XY$  is bounded by the curve

$$x^2 + y^2 = a^2. \tag{2}$$

From (1) we get

$$\frac{\sqrt{(D_x f)^2 + (D_y f)^2 + (D_z f)^2}}{D_z f} = \sqrt{1 + \frac{c^2}{a^2} \frac{(x^2 + y^2)^2}{x^2 y^2}},$$

where  $x, y$  are the coördinates of any point within the projection of the base of the cone.

Since the four faces of the cone are equal, the required surface

$$\sigma = \frac{4}{a} \int_0^a \int_0^{(a^2-x^2)^{\frac{1}{2}}} \sqrt{a^2 x^2 y^2 + c^2 (x^2 + y^2)^2} \cdot dy dx. \tag{3}$$

Let us substitute  $v^2 = x$  and  $w^2 = y$ , whence  $dx = 3v^2 dv$  and  $dy = 3w^2 dw$ , and we have

$$\sigma = \frac{36}{a} \int_0^{a^{\frac{1}{2}}} \int_0^{(a^{\frac{1}{2}} - v^2)^{\frac{1}{2}}} vw \sqrt{a^2 v^2 w^2 + c^2 (v^2 + w^2)^2} \cdot dw dv;$$

or, since in a definite integral it makes no difference what letters we use for the variables,

$$\sigma = \frac{36}{a} \int_0^{a^{\frac{1}{2}}} \int_0^{(a^{\frac{1}{2}} - x^2)^{\frac{1}{2}}} xy \sqrt{a^2 x^2 y^2 + c^2 (x^2 + y^2)^2} \cdot dy dx. \quad (4)$$

The  $x$  and  $y$  in (4), however, must not be confounded with the  $x$  and  $y$  in (3).

The integral in (4) is precisely that which we should have to find if we sought the area of a surface of such a nature that its projection on the plane of  $XY$  was a quadrant of the circle  $x^2 + y^2 = a^{\frac{1}{2}}$ , and the secant of the angle made by the tangent plane at any point  $(x, y, z)$  of the surface with the plane of  $XY$  was  $xy \sqrt{a^2 x^2 y^2 + c^2 (x^2 + y^2)^2}$ .

In the latter problem there is nothing to prevent our replacing  $x$  and  $y$  in  $xy \sqrt{a^2 x^2 y^2 + c^2 (x^2 + y^2)^2}$  by their values in terms of  $r$  and  $\phi$ , the polar coördinates of any point of the projection  $x^2 + y^2 = a^{\frac{1}{2}}$ , and dividing this projection into polar elements instead of rectangular elements, and then integrating between the limits which we should use if we were finding the area of the projection by the formula  $A = \iint r d\phi dr$ .

We have, then,

$$\sigma = \frac{36}{a} \int_0^{\frac{\pi}{2}} \int_0^{a^{\frac{1}{2}}} r^2 \sin \phi \cos \phi \sqrt{a^2 r^4 \sin^2 \phi \cos^2 \phi + c^2 r^4} \cdot r dr d\phi,$$

or

$$\sigma = \frac{36}{a} \int_0^{\frac{\pi}{2}} \int_0^{a^{\frac{1}{2}}} r^5 \sin \phi \cos \phi \sqrt{a^2 \sin^2 \phi \cos^2 \phi + c^2} \cdot dr d\phi,$$

$$\sigma = 6a \int_0^{\frac{\pi}{2}} \sin \phi \cos \phi \sqrt{a^2 \sin^2 \phi \cos^2 \phi + c^2} \cdot d\phi.$$

Substitute  $u = \sin^2 \phi$ , and

$$\sigma = 3a \int_0^1 \sqrt{a^2 u (1-u) + c^2} \cdot du,$$

$$\sigma = \frac{3}{4} \left[ 2ac + (a^2 + 4c^2) \tan^{-1} \frac{a}{2c} \right].$$

EXAMPLES.

(1) Find the area included by the cylinders described in Art. 145 by direct integration.

(2) A square hole is cut through a sphere, the axis of the hole coinciding with a diameter of the sphere; find the area of the surface removed.

(3) A cylinder is constructed on a single loop of the curve  $r = a \cos n\phi$ , having its generating lines perpendicular to the plane of this curve; determine the area of the portion of the surface of the sphere  $x^2 + y^2 + z^2 = a^2$  which the cylinder intercepts.

$$\text{Ans. } \frac{4a^2}{n} \left( \frac{\pi}{2} - 1 \right).$$

(4) Find the area of the portion of the surface of the cone described in Art. 146 included by the cylinder  $x^2 + y^2 = b^2$ .

$$\text{Ans. } \frac{2b^2}{a} \left[ 2\sqrt{a^2 + 3c^2} \tan^{-1} \left( \frac{\sqrt{a^2 + 3c^2}}{c} \right) - a \tan^{-1} \frac{a}{2c} \right].$$

(5) Find the area of the portion of the surface of the sphere  $x^2 + y^2 + z^2 = 2ay$  cut out by one nappe of the cone  $Ax^2 + Bz^2 = y^2$ .

$$\text{Ans. } \frac{4\pi a^2}{\sqrt{(1+A)(1+B)}}.$$

(6) Find the area of the portion of the surface of the sphere  $x^2 + y^2 + z^2 = 2ay$  lying within the paraboloid  $y = Ax^2 + Bz^2$ .

$$\text{Ans. } \frac{2\pi a}{\sqrt{AB}}.$$

(7) The centre of a regular hexagon moves along a diameter of a given circle (radius =  $a$ ), the plane of the hexagon being perpendicular to this diameter, and its magnitude varying in such a manner that one of its diagonals always coincides with a chord of the circle; find the surface generated.

$$\text{Ans. } a^2(2\pi + 3\sqrt{3}).$$

## CHAPTER XII.


## VOLUMES.

*Single Integration.*

147. If sections of a solid are made by parallel planes, and a set of cylinders drawn, each having for its base one of the sections, and for its altitude the distance between two adjacent cutting planes, the limit of the sum of the volumes of these cylinders, as the distance between the sections is indefinitely decreased, is the volume of the solid.

We shall take as established by Geometry the fact that the volume of a cylinder or prism is the product of the area of its base by its altitude.

It follows from what has just been said, that if, in a given solid, all of a set of parallel sections are equal, the volume of the solid is its base by its altitude, no matter how irregular its form.



Let us find the volume of a pyramid having  $b$  for the area of its base, and  $a$  for its altitude.

Divide the pyramid by planes parallel to the base, and let  $z$  be the area of a section at the distance  $x$  from the vertex.

We know from Geometry that  $\frac{z}{b} = \frac{x^2}{a^2}$ .

Hence  $z = \frac{b}{a^2}x^2$ .

Let the distance between two adjacent sections be  $dx$ ; then the volume of the cylinder on  $z$  is

$$\frac{b}{a^2}x^2dx,$$

and  $V$ , the required volume of the pyramid, is


$$V = \frac{b}{a^2} \int_0^a x^2 dx = \frac{ab}{3}.$$

Precisely the same reasoning applies to any cone, which will therefore have for its volume one-third the product of its base by its altitude.

EXAMPLE.

Find the volume of the frustum of a pyramid or of a cone.

148. If a line move keeping always parallel to a given plane, and touching a plane curve and a straight line parallel to the plane of the curve, the surface generated is called a *conoid*. Let us find the volume of a *conoid* when the director line and curve are perpendicular to the given plane.



Divide the conoid into laminae by planes parallel to the fixed plane.

Let  $\Delta y$  be the distance between two adjacent sections, and let  $x$  be the length of the line in which any section cuts the base of the conoid; let  $a$  be the altitude and  $b$  the area of the base of the figure. Any one of our elementary cylinders will have for its volume  $\frac{1}{2}ax\Delta y$ , since the area of its triangular base is  $\frac{1}{2}ax$ , and we have  $V = \frac{1}{2}a \int xdy$ , the limits of integration being so taken as to embrace the whole solid.  $\int xdy$  between the limits in question is the area of the base of the conoid; hence its volume,

$$V = \frac{1}{2}ab.$$

EXAMPLES.

(1) Find the volume of a conoid when the director line and curve are not perpendicular to the given plane.

(2) A woodman fells a tree 2 feet in diameter, cutting half-way through from each side. The lower face of each cut is horizontal, and the upper face makes an angle of  $45^\circ$  with the horizontal. How much wood does he cut out?



149. To find the volume of an ellipsoid.

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

Take the cutting planes parallel to the plane of  $XY$ . A section at the distance  $z$  from the origin will have

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 - \frac{z^2}{c^2} = \frac{c^2 - z^2}{c^2}$$

for its equation, and  $\frac{a}{c}\sqrt{c^2 - z^2}$  and  $\frac{b}{c}\sqrt{c^2 - z^2}$  for its semi-axes; hence its area will be  $\frac{\pi ab}{c^2}(c^2 - z^2)$ .

Any of the elementary cylinders will have for its volume  $\frac{\pi ab}{c^2}(c^2 - z^2)\Delta z$ , and we shall have for the whole solid

$$V = \frac{\pi ab}{c^2} \int_{-c}^c (c^2 - z^2) dz.$$

$$V = \frac{4}{3} \pi abc.$$

If  $a$ ,  $b$ , and  $c$  are equal, the ellipsoid is a sphere, and

$$V = \frac{4}{3} \pi a^3.$$

#### EXAMPLES.

(1) Find the volume included between an hyperboloid of one sheet

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1,$$

and its asymptotic cone

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0.$$

*Ans.* It is equal to a cylinder of the same altitude as the solid in question, and having for a base the section made by the plane of  $XY$ .

(2) Find the whole volume of the solid bounded by the surface

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^4}{c^4} = 1.$$

$$\text{Ans. } \frac{8\pi abc}{5}.$$

- (3) Find the volume cut from the surface

$$\frac{x^2}{c} + \frac{y^2}{b} = 2x$$

by a plane parallel to the plane of ( $YZ$ ) at a distance  $a$  from it.

$$\text{Ans. } \pi a^2 \sqrt{bc}.$$

(4) The centre of a regular hexagon moves along a diameter of a given circle (radius =  $a$ ), the plane of the hexagon being perpendicular to this diameter, and its magnitude varying in such a manner that one of its diagonals always coincides with a chord of the circle; find the volume generated.

$$\text{Ans. } 2\sqrt{3}.a^3.$$

(5) A circle (radius =  $a$ ) moves with its centre on the circumference of an equal circle, and keeps parallel to a given plane which is perpendicular to the plane of the given circle; find the volume of the solid it will generate.

$$\text{Ans. } \frac{2a^3}{3}(3\pi + 8).$$

*Solids of Revolution. Single Integration.*

150. If a solid is generated by the revolution of a plane curve  $y = fx$  about the axis of  $x$ , sections made by planes perpendicular to the axis are circles. The area of any such circle is  $\pi y^2$ , the volume of the elementary cylinder is  $\pi y^2 \Delta x$ , and

$$V = \pi \int_{x_0}^x y^2 dx$$

is the volume of the solid generated.

For example; let us find the volume of the solid generated by the revolution of one branch of the tractrix about the axis of  $X$ . Here we must integrate from  $x = 0$  to  $x = \infty$ .

$$V = \pi \int_0^{\infty} y^2 dx.$$

We have  $dx = -\frac{(a^2 - y^2)^{\frac{1}{2}}}{y} dy$  (Art. 102 [2].)  
in the case of the tractrix;

hence 
$$V = -\pi \int_{x=0}^{x=\infty} y(a^2 - y^2)^{\frac{1}{2}} dy.$$

When  $x = 0$ ,  $y = a$ , and when  $x = \infty$ ,  $y = 0$ .

Therefore 
$$V = -\pi \int_a^0 y(a^2 - y^2)^{\frac{1}{2}} dy = \frac{\pi a^3}{3}.$$

#### EXAMPLES.

(1) If the plane curve revolves about the axis of  $Y$ ,

$$V = \pi \int_{y_0}^{y_1} x^2 dy.$$

(2) The volume of a sphere is  $\frac{4}{3} \pi a^3$ .

(3) The volume of the solid formed by the revolution of a cycloid about its base is  $5 \pi^2 a^3$ .

(4) The curve  $y^2(2a - x) = x^2$  revolves about its asymptote; show that the volume generated is  $2 \pi^2 a^3$ .

(5) The curve  $x^{\frac{1}{2}} + y^{\frac{1}{2}} = a^{\frac{1}{2}}$  revolves about the axis of  $X$ ; show that the volume generated is  $\frac{8}{15} \pi a^3$ .

#### *Solids of Revolution. Double Integration.*

151. If we suppose the area of the revolving curve broken up into infinitesimal rectangles as in Art. 137, the element  $\Delta x \Delta y$  at any point  $P$ , whose coördinates are  $x$  and  $y$ , will generate a ring the volume of which will differ from  $2 \pi y \Delta x \Delta y$  by an amount which will be an infinitesimal of higher order than the second if we regard  $\Delta x$  and  $\Delta y$  as of the first order. For the ring in question is obviously greater than a prism having the same cross-section  $\Delta x \Delta y$ , and having an altitude equal to the inner circumference  $2 \pi y$  of the ring, and is less than a prism having  $\Delta x \Delta y$  for its base and  $2 \pi (y + \Delta y)$ , the outer circumference of the ring, for its altitude; but these two prisms differ by  $2 \pi \Delta x (\Delta y)^2$ , which is of the third order.

$\Delta x \int_0^y 2\pi y dy$ , where the upper limit of integration is the ordinate of the point of the curve immediately above  $P$ , and must be expressed in terms of  $x$  by the aid of the equation of the revolving curve, will give us the elementary cylinder used in Art. 150.

The whole volume required will be the limit of the sum of these cylinders; that is,

$$V = 2\pi \int_{x_0}^{x_1} \int_0^y y dy dx. \quad [1]$$

If the figure revolved is bounded by two curves, the required volume can be found by the formula just obtained, if the limits of integration are suitably chosen.

Let us consider the following example:

A paraboloid of revolution has its axis coincident with the diameter of a sphere, and its vertex in the surface of the sphere; required the volume between the two surfaces.

Let  $y^2 = 2mx$  (1)

be the parabola, and  $x^2 + y^2 - 2ax = 0$  (2)

be the circle, which form the paraboloid and the sphere by their revolution. The abscissas of their points of intersection are 0 and  $2(a - m)$ .

We have  $V = 2\pi \int \int y dy dx,$

and, in performing our first integration, our limits must be the values of  $y$  obtained from equations (1) and (2).

We get  $V = \pi \int [2(a - m)x - x^2] dx,$

and here our limits of integration are 0 and  $2(a - m)$ .

Hence  $V = \frac{4}{3} \pi (a - m)^3 = \frac{\pi h^3}{6},$

if  $h$  is the altitude of the solid in question.

EXAMPLES.

(1) A cone of revolution and a paraboloid of revolution have the same vertex and the same base; required the volume between them.

*Ans.*  $\frac{\pi m h^2}{3}$ , where  $h$  is the altitude of the cone.

(2) Find the volume included between a right cone, whose vertical angle is  $30^\circ$ , and a sphere of given radius touching it along a circle.

$$\text{Ans. } \frac{\pi r^3}{6}.$$

*Solids of Revolution. Polar Formula.*

152. If we use polar coördinates, and suppose the revolving area broken up, as in Art. 138, into elements of which  $rd\phi dr$  is the one at any point  $P$  whose coördinates are  $r$  and  $\phi$ , the element  $rd\phi dr$  will generate a ring whose volume will differ from  $2\pi r^2 \sin \phi d\phi dr$  by an infinitesimal of higher order than the second, if we regard  $d\phi$  and  $dr$  as of the first order; for it will be less than a prism having for its base  $rd\phi dr$ , and for its altitude  $2\pi(r+dr) \sin(\phi+d\phi)$ , and greater than a prism having the same base and the altitude  $2\pi r \sin \phi$ ; and these prisms differ by an amount which is infinitesimal of higher order than the second.

We shall have then

$$V = 2\pi \int \int r^2 \sin \phi dr d\phi, \quad [1]$$

the limits being so taken as to bring in the whole of the generating area.

For example; let us find the volume generated by the revolution of a cardioide about its axis.

$$r = 2a(1 - \cos \phi)$$

is the equation of the cardioide;

$$V = 2\pi \int \int r^2 \sin \phi dr d\phi.$$

Our first integral must be taken between the limits  $r = 0$  and  $r = 2a(1 - \cos \phi)$ , and is

$$\frac{8a^3}{3}(1 - \cos \phi)^3 \sin \phi d\phi.$$

$$V = \frac{16}{3}a^3 \pi \int_0^\pi (1 - \cos \phi)^3 \sin \phi d\phi,$$


$$V = \frac{64}{3}\pi a^3.$$

EXAMPLE.

A right cone has its vertex on the surface of a sphere, and its axis coincident with the diameter of the sphere passing through that point ; find the volume common to the cone and the sphere.

*Volume of any Solid. Triple Integration.*

153. If we suppose our solid divided into parallelepipeds by planes parallel to the three coördinate planes, the elementary



paralleloiped at any point  $(x,y,z)$  within the solid will have for its volume  $\Delta x \Delta y \Delta z$ , or, if we regard  $x$ ,  $y$ , and  $z$  as independent,  $dx dy dz$  ; and the whole volume

$$V = \iiint dx dy dz, \tag{1}$$

the limits being so chosen as to embrace the whole solid.

The integrations are independent, and may be performed in any order if the limits are suitably chosen.

As it is important to have a perfectly clear conception of the geometrical interpretation of each step in the process of finding

a volume by a triple integration, we will consider one case in detail.

Let the integrations be performed in the order indicated by the formula

$$V = \iiint dzdydx.$$

If the limits are correctly chosen, our first integration gives us the volume of a prism one of whose lateral edges passes through any chosen point  $P, (x, y, z)$  within the solid, is parallel to the axis of  $Z$ , and reaches directly across the solid from surface to surface, while the base of the prism is the rectangle  $dydx$ ; our second integration gives the volume of a right cylinder whose base is a plane section of the solid, passes through the point  $P$ , and is parallel to the plane  $YZ$ , and whose altitude is  $dx$ ; and our third integration gives the volume of the whole solid.

The limits in our first integration are, then, the values of  $z$  belonging to the point in the lower bounding surface and the point in the upper bounding surface which have the coördinates  $x$  and  $y$ ; the limits in the second integration are the values of  $y$  belonging to the two points in the perimeter of the projection of the solid in the plane of  $XY$  which have the coördinate  $x$ ; and the limits in the third integration are the least value and the greatest value of  $x$  belonging to points on the perimeter of the projection of the solid on the plane of  $XY$ .

It is easily seen from what has just been said that the limits in the second and third integrations are precisely those we should use if we were finding the area of the projection of the solid by the formula

$$A = \iint dydx.$$

Of course, it is necessary to have a clear idea of the form of the solid whose volume is required.

For example, let us find the volume of the portion of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

cut off by the coördinate planes.

$$V = \iiint dzdydx,$$

and our limits are, for  $z$ , 0 and  $c\sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}}$ ; for  $y$ , 0 and  $b\sqrt{1 - \frac{x^2}{a^2}}$ ; and for  $x$ , 0 and  $a$ . For, starting at any point  $(x, y, z)$  and integrating on the hypothesis that  $z$  alone varies, we get a column of our elementary parallelepipeds having  $dx dy$  as a base and passing through the point  $(x, y, z)$ . To make this column reach from the plane  $XY$  to the surface,  $z$  must increase from the value zero to the value belonging to the point on the surface of the ellipsoid which has the coördinates  $x$  and  $y$ ; that is, to the value  $c\sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}}$ . Then, integrating on the hypothesis that  $y$  alone varies, we shall sum these columns and shall get a slice of the solid passing through  $(x, y, z)$  and having the thickness  $dx$ . To make this slice reach completely across the solid, we must let  $y$  increase from the value zero to the greatest value it can have in the slice in question; that is, to the value which is the ordinate of that point of the section of the ellipsoid by the plane  $XY$  which has the abscissa  $x$ . The section in question has the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1;$$

therefore the required value of  $y$  is  $b\sqrt{1 - \frac{x^2}{a^2}}$ .

Last, in integrating on the hypothesis that  $x$  alone varies, we must choose our limits so as to include all the slices just described, and must increase  $x$  from zero to  $a$ .

$$\int dz = z = c\sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}}$$

between the limits 0 and  $c\sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}}$



$$\begin{aligned}
c \int \sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}} \cdot dy \\
&= \frac{c}{b} \int \sqrt{b^2 \left(1 - \frac{x^2}{a^2}\right) - y^2} \cdot dy \\
&= \frac{1}{2} \frac{c}{b} \left[ y \sqrt{b^2 \left(1 - \frac{x^2}{a^2}\right) - y^2} + b^2 \left(1 - \frac{x^2}{a^2}\right) \sin^{-1} \frac{y}{b \sqrt{1 - \frac{x^2}{a^2}}} \right] \\
&= \frac{\pi b c}{4} \left(1 - \frac{x^2}{a^2}\right)
\end{aligned}$$

between the limits 0 and  $b \sqrt{1 - \frac{x^2}{a^2}}$ .

$$\frac{\pi b c}{4} \int_0^a \left(1 - \frac{x^2}{a^2}\right) dx = \frac{\pi a b c}{6},$$

the volume required.

#### EXAMPLES.

(1) Find the volume obtained in the present article, performing the integrations in the order indicated by the formula,

$$V = \iiint dx dy dz.$$

(2) Find the volume cut off from the surface

$$\frac{z^2}{c} + \frac{y^2}{b} = 2x$$

by a plane parallel to that of  $YZ$ , at a distance  $a$  from it.

$$\text{Ans. } \pi a^2 \sqrt{bc}.$$

(3) Find the volume enclosed by the surfaces,

$$x^2 + y^2 = cz, \quad x^2 + y^2 = ax, \quad z = 0. \quad \text{Ans. } \frac{3\pi a^4}{32c}$$

(4) Obtain the volume bounded by the surface

$$z = a - \sqrt{x^2 + y^2}$$

and the planes

$$x = z \quad \text{and} \quad x = 0,$$

$$\text{Ans. } \frac{2a^3}{9}.$$

(5) Find the volume of the conoid bounded by the surface  $z^2 + \frac{\alpha^2 y^2}{x^2} = c^2$  and the planes  $x = 0$  and  $x = a$ . *Ans.*  $\frac{\pi c^2 a}{2}$ .

154. If we use polar coördinates we can take as our element of volume

$$r^2 \sin \phi dr d\phi d\theta,$$

an expression easily obtained from the element  $2\pi r^2 \sin \phi dr d\phi$  used in Art. 152.

Then 
$$V = \iiint r^2 \sin \phi dr d\phi d\theta,$$

where the order of the integrations is usually immaterial if the limits are properly chosen.

#### EXAMPLES.

(1) Find the volume of a sphere by polar coördinates.

(2) Find the whole volume of the solid bounded by

$$(x^2 + y^2 + z^2)^3 = 27 \alpha^3 xyz.$$

*Suggestion:* Transform to polar coördinates. *Ans.*  $\frac{9}{2} \alpha^3$ .

## CHAPTER XIII.

## CENTRES OF GRAVITY.


155. The *moment of a force about an axis* perpendicular to its line of direction is the product of the magnitude of the force by the perpendicular distance of its line of direction from the axis, and measures the tendency of the force to produce rotation about the axis.

The *force exerted by gravity* on any material body is proportional to the mass of the body, and may be measured by the mass of the body.

The *Centre of Gravity* of a body is a point so situated that the force of gravity produces no tendency in the body to rotate about any axis passing through this point.

The subject of centres of gravity belongs to Mechanics, and we shall accept the definitions and principles just stated as data for mathematical work, without investigating the mechanical grounds on which they rest.

156. Suppose the points of a body referred to a set of three rectangular axes fixed in the body, and let  $\bar{x}, \bar{y}, \bar{z}$  be the coördinates of the centre of gravity. Place the body with the axes of  $X$  and  $Z$  horizontal, and consider the tendency of the particles of the body to produce rotation about an axis through  $(\bar{x}, \bar{y}, \bar{z})$  parallel to  $OZ$ , under the influence of gravity. Represent the mass of an elementary parallelepiped at any point  $(x, y, z)$  by  $dm$ . The force exerted by gravity on  $dm$  is measured by  $dm$ , and its line of direction is vertical. If the mass of  $dm$  were concentrated at  $P$ , the moment of the force exerted on  $dm$  about the



axis through  $C$  would be  $(x - \bar{x})dm$ , and this moment would represent the tendency of  $dm$  to rotate about the axis in question; the tendency of the whole body to rotate about this axis would be  $\Sigma(x - \bar{x})dm$ . If now we decrease  $dm$  indefinitely, the error committed in assuming that the mass of  $dm$  is concentrated at  $P$  decreases indefinitely, and we shall have as the true expression for the tendency of the whole body to rotate about the axis through  $C$ ,  $\int(x - \bar{x})dm$ ; but this must be zero.

$$\begin{aligned} \text{Hence} \quad \int(x - \bar{x})dm &= 0, \\ \int xdm - \bar{x} \int dm &= 0, \\ \bar{x} &= \frac{\int xdm}{\int dm}. \end{aligned} \quad [1]$$

If we place the body so that the axes of  $Y$  and  $X$  are horizontal, the same reasoning will give us

$$\bar{y} = \frac{\int ydm}{\int dm}; \quad [2]$$

and in like manner we can get

$$\bar{z} = \frac{\int zdm}{\int dm}. \quad [3]$$

Since  $\int dm$  is the mass of the whole body, if we represent it by  $M$  we shall have

$$\begin{aligned} \bar{x} &= \frac{\int xdm}{M}, \\ \bar{y} &= \frac{\int ydm}{M}, \\ \bar{z} &= \frac{\int zdm}{M}. \end{aligned}$$

## EXAMPLE.

Show that the effect of gravity in making a body tend to rotate about any given axis is precisely the same as if the mass of the body were concentrated at its centre of gravity.

157. The mass of any homogeneous body is the product of its volume by its density. If the body is not homogeneous, the density at any point will be a function of the position of that point. Let us represent it by  $\kappa$ . Then we may regard  $dm$  as equal to  $\kappa dv$  if  $dv$  is the element of volume, and we shall have

$$\bar{x} = \frac{\int x\kappa dv}{\int \kappa dv} \quad [1]$$

and corresponding formulas for  $\bar{y}$  and  $\bar{z}$ .

If the body considered is homogeneous,  $\kappa$  is constant, and we shall have

$$\bar{x} = \frac{\int x dv}{\int dv} = \frac{\int x dv}{V}, \quad [2]$$

$$\bar{y} = \frac{\int y dv}{\int dv} = \frac{\int y dv}{V}, \quad [3]$$

$$\bar{z} = \frac{\int z dv}{\int dv} = \frac{\int z dv}{V}. \quad [4]$$

In any particular problem we have only to express  $dv$  in terms of the coördinates.

*Plane Area.*

158. If we use rectangular coördinates, and are dealing with a plane area, where the weight is uniformly distributed, we have

$$dv = dA = dx dy. \quad (\text{Art. 136}).$$

Hence, by 157, [2] and [3],

$$\left. \begin{aligned} \bar{x} &= \frac{\int \int x dx dy}{\int \int dx dy} \\ \bar{y} &= \frac{\int \int y dx dy}{\int \int dx dy} \end{aligned} \right\} \quad [1]$$

If we use polar coördinates,

$$dv = dA = r d\phi dr,$$

and

$$\left. \begin{aligned} \bar{x} &= \frac{\int \int r^2 \cos \phi d\phi dr}{\int \int r d\phi dr} \\ \bar{y} &= \frac{\int \int r^2 \sin \phi d\phi dr}{\int \int r d\phi dr} \end{aligned} \right\} \quad [2]$$

For example; let us find the *centre of gravity* of the area between the *cissoïd* and its asymptote. From the equation of the *cissoïd*

$$y^2 = \frac{x^3}{a-x},$$

we see that the curve is symmetrical with respect to the axis of *X*, passes through the origin, and has the line  $x = a$  as an asymptote. From the symmetry of the area in question,  $\bar{y} = 0$ , and we need only find  $\bar{x}$ .

$$\bar{x} = \frac{\int_0^a \int_{-y}^y x dy dx}{\int_0^a \int_{-y}^y dy dx} = \frac{\int_0^a xy dx}{\int_0^a y dx},$$

$$\bar{x} = \frac{\int_0^a \frac{x^{\frac{3}{2}}}{(a-x)^{\frac{1}{2}}} dx}{\int_0^a \frac{x^{\frac{1}{2}}}{(a-x)^{\frac{1}{2}}} dx} = \frac{\frac{2}{3} a \int_0^a \frac{x^{\frac{1}{2}}}{(a-x)^{\frac{1}{2}}} dx}{\int_0^a \frac{x^{\frac{1}{2}}}{(a-x)^{\frac{1}{2}}} dx}; \text{ by Art. 64 [4].}$$

$$\bar{x} = \frac{2}{3} a.$$

As an example of the use of the polar formulas [2], let us find the *centre of gravity* of the cardioid

$$r = 2a(1 - \cos \phi).$$

Here, from the fact that the axis of  $X$  is an axis of symmetry, we know that  $\bar{y} = 0$ .

$$\bar{x} = \frac{\int_0^{2\pi} \int_0^r r^2 \cos \phi dr d\phi}{\int_0^{2\pi} \int_0^r r \bar{r} dr d\phi}$$

$$= \frac{\frac{1}{3} \int_0^{2\pi} r^3 \cos \phi d\phi}{\frac{1}{2} \int_0^{2\pi} r^2 d\phi} = \frac{\frac{8a^3}{3} \int_0^{2\pi} (1 - \cos \phi)^3 \cos \phi d\phi}{2a^2 \int_0^{2\pi} (1 - \cos \phi)^2 d\phi},$$

$$\int_0^{2\pi} (\cos \phi - 3 \cos^2 \phi + 3 \cos^3 \phi - \cos^4 \phi) d\phi = -\frac{1}{4} \pi;$$

and  $\int_0^{2\pi} (1 - 2 \cos \phi + \cos^2 \phi) d\phi = 3\pi.$

Hence  $\bar{x} = -\frac{1}{4} a.$

#### EXAMPLES.

1. Show that formulas [1] hold even when we use *oblique coördinates*.

2. Find the centre of gravity of a segment of a parabola cut off by any chord.

*Ans.*  $\bar{x} = \frac{2}{3} a, \bar{y} = 0.$  If the axes are the tangent parallel to the chord and the diameter bisecting the chord.

3. Find the centre of gravity of the area bounded by the semi-cubical parabola  $ay^2 = x^3$  and a double ordinate. *Ans.*  $\bar{x} = \frac{4}{3}x$ .

4. Find the centre of gravity of a semi-ellipse, the bisecting line being any diameter.

*Ans.* If the bisecting diameter is taken as the axis of  $Y$ , and the conjugate diameter as the axis of  $X$ ,  $\bar{x} = \frac{4a}{3\pi}$ ,  $\bar{y} = 0$ .

5. Find the centre of gravity of the curve  $y^2 = b^2 \frac{a-x}{x}$ .

*Ans.*  $\bar{x} = \frac{1}{2}a$ .

6. Find the centre of gravity of the cycloid.

*Ans.*  $\bar{x} = a\pi$ ,  $\bar{y} = \frac{3}{8}a$ .

7. Find the centre of gravity of the lemniscate  $r^2 = a^2 \cos 2\phi$ .

*Ans.*  $\bar{x} = \frac{\pi\sqrt{2}}{8}a$ .

8. Find the centre of gravity of a circular sector.

*Ans.* If we take the radius bisecting the sector as the axis of  $X$ , and represent the angle of the sector by  $2\alpha$ ,  $\bar{x} = \frac{3}{8} \frac{a \sin \alpha}{\alpha}$ .

9. Find the centre of gravity of the segment of an ellipse cut off by a quadrantal chord. *Ans.*  $\bar{x} = \frac{3}{8} \frac{a}{\pi - 2}$ ,  $\bar{y} = \frac{3}{8} \frac{b}{\pi - 2}$ .

10. Find the centre of gravity of a quadrant of the area of the curve  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$ . *Ans.*  $\bar{x} = \bar{y} = \frac{25}{48} \frac{a}{\pi}$ .

159. If we are dealing with a homogeneous solid formed by the revolution of a plane curve about the axis of  $X$ , we have

$$dv = 2\pi y dy dx. \quad (\text{Art. 151 [1]})$$

Hence, by Art. 157 [2],

$$\bar{x} = \frac{\int \int xy dx dy}{\int \int y dx dy}. \quad [1]$$



If we use polar coördinates,

$$dv = 2\pi r^2 \sin \phi dr d\phi. \quad (\text{Art. 152 [1].})$$

$$\text{Hence} \quad \bar{x} = \frac{\int \int r^3 \sin \phi \cos \phi dr d\phi}{\int \int r^2 \sin \phi dr d\phi}. \quad [2]$$

For example ; let us find the centre of gravity of a hemisphere. The equation of the revolving curve is  $x^2 + y^2 = a^2$ .

$$\bar{x} = \frac{\int_0^a \int_0^{\sqrt{a^2-x^2}} xy dy dx}{\int_0^a \int_0^{\sqrt{a^2-x^2}} y dy dx} = \frac{\frac{1}{3} a^4}{\frac{1}{3} a^3} = \frac{3}{8} a.$$

If we use polar coördinates the equation of the revolving curve is  $r = a$ .

$$\text{Here} \quad \bar{x} = \frac{\int_0^a \int_0^{\frac{\pi}{2}} r^3 \sin \phi \cos \phi d\phi dr}{\int_0^a \int_0^{\frac{\pi}{2}} r^2 \sin \phi d\phi dr} = \frac{\frac{1}{8} a^4}{\frac{1}{3} a^3} = \frac{3}{8} a.$$

#### EXAMPLES.

1. Find the centre of gravity of the solid formed by the revolution of the sector of a circle about one of its extreme radii.

*Ans.*  $\bar{x} = \frac{3}{4} a \cos^2 \frac{1}{2} \beta$ , where  $\beta$  is the angle of the sector.

2. Find the centre of gravity of the segment of a paraboloid of revolution cut off by a plane perpendicular to the axis.

*Ans.*  $\bar{x} = \frac{3}{8} a$ , where  $x = a$  is the plane.

3. Find the centre of gravity of the solid formed by scooping out a cone from a given paraboloid of revolution, the bases of the two volumes being coincident as well as their vertices.

*Ans.* The centre of gravity bisects the axis.

4. A cardioide is made to revolve about its axis; find the centre of gravity of the solid generated. *Ans.*  $\bar{x} = -\frac{2}{3}a$ .

5. Obtain formulas for the centre of gravity of any homogeneous solid.

6. Find the centre of gravity of the solid bounded by the surface  $z^2 = xy$  and the five planes  $x=0$ ,  $y=0$ ,  $z=0$ ,  $x=a$ ,  $y=b$ .  
*Ans.*  $\bar{x} = \frac{2}{3}a$ ,  $\bar{y} = \frac{2}{3}b$ ,  $\bar{z} = \frac{2}{3}\sqrt{\frac{1}{2}ab}$ .

160. If we are dealing with the arc of a plane curve, the formulas of Art. 157 reduce to

$$\bar{x} = \frac{\int x ds}{\int ds}, \quad [1]$$

$$\bar{y} = \frac{\int y ds}{\int ds}. \quad [2]$$

#### EXAMPLES.

1. Find the centre of gravity of an arc of a circle, taking the diameter bisecting the arc as the axis of  $X$  and the centre as the origin.  
*Ans.*  $\bar{x} = \frac{ac}{s}$ , where  $c$  is the chord of the arc.

2. Find the centre of gravity of the arc of the curve  $x^2 + y^2 = a^2$  between two successive cusps.  
*Ans.*  $\bar{x} = \bar{y} = \frac{2}{3}a$ .

3. Find the centre of gravity of the arc of a semi-cycloid.  
*Ans.*  $\bar{x} = (\pi - \frac{4}{3})a$ ,  $\bar{y} = -\frac{2}{3}a$ .

4. Find the centre of gravity of the arc of a catenary cut off by any horizontal chord.

*Ans.*  $\bar{x} = 0$ ,  $\bar{y} = \frac{ax + sy}{2s}$ , where  $2s$  is the length of the arc.

5. Obtain formulas for the centre of gravity of a surface of revolution, the weight being uniformly distributed over the surface.

6. Find the centre of gravity of any zone of a sphere.

*Ans.* The centre of gravity bisects the line joining the centres of the bases of the zone.

7. A cardioide revolves about its axis; find the centre of gravity of the surface generated. *Ans.*  $\bar{x} = -\frac{1}{8} \frac{a}{3}$ .

8. Find the centre of gravity of the surface of a hemisphere when the density at each point of the surface varies as its perpendicular distance from the base of the hemisphere.

$$\text{Ans. } \bar{x} = \frac{2}{3} a.$$

9. Find the centre of gravity of a quadrant of a circle, the density at any point of which varies as the  $n$ th power of its distance from the centre.

$$\text{Ans. } \bar{x} = \bar{y} = \frac{n+2}{n+3} \cdot \frac{2a}{\pi}.$$

10. Find the centre of gravity of a hemisphere, the density of which varies as the distance from the centre of the sphere.

$$\text{Ans. } \bar{x} = \frac{2}{3} a.$$

*Properties of Guldin.*

161. I. If a plane area revolve about an axis external to itself through any assigned angle, the volume of the solid generated will be equal to a prism whose base is the revolving area and whose altitude is the length of the path described by the centre of gravity of the area.

II. If the arc of a plane curve revolve about an external axis in its own plane through any assigned angle, the area of the surface generated will be equal to that of a rectangle, one side of which is the length of the revolving curve, and the other the length of the path described by its centre of gravity.

First; let the area in question revolve about the axis of  $X$  through an angle  $\Theta$ . The ordinate of the centre of gravity of the area in question is

$$\bar{y} = \frac{\iint y dx dy}{\iint dx dy}, \quad \text{by Art. 158 [1].}$$

The length of the path described by the centre of gravity

$$\bar{y}\Theta = \frac{\Theta \iint y dx dy}{\iint dx dy}. \quad (1)$$

The volume generated is

$$V = \Theta \iint y dx dy, \quad \text{by Art. 151.}$$

Hence 
$$V = \bar{y}\Theta \iint dx dy.$$

But  $\iint dx dy$  is the revolving area, and the first theorem is established.

We leave the proof of the second theorem to the student.

#### EXAMPLES.

1. Find the surface and volume of a sphere, regarding it as generated by the revolution of a semicircle.

2. Find the surface and volume of the solid generated by the revolution of a cycloid about its base.

3. Find the volume and the surface of the ring generated by the revolution of a circle about an external axis.

*Ans.*  $V = 2\pi^2 a^2 b$ ,  $S = 4\pi^2 ab$ , where  $b$  is the distance of the centre of the circle from the axis.

4. Find the volume of the ring generated by the revolution of an ellipse about an external axis.

*Ans.*  $V = 2\pi^2 abc$ , where  $c$  is the distance of the centre of the ellipse from the axis.

## CHAPTER XIV.

## LINE, SURFACE, AND SPACE INTEGRALS.

162. Any variable which depends for its value solely upon the position of a point, as, for example, any function of the rectangular or polar coördinates of the point, may be called a *point-function*.

A point-function is said to be continuous along a given line if its value changes continuously as the point, on whose position the function depends for its value, moves along the line; it is said to be continuous over a given surface if its value changes continuously as the point is made to move at pleasure over the surface; and it is said to be continuous throughout a given space if its value changes continuously as the point is made to move about at pleasure within the space.

163. If a given line is divided in any way into infinitesimal elements, and the length of each element is multiplied by the value a given point-function, which is continuous along the line, has at some point within the element, the limit approached by the sum of these products as each element is indefinitely decreased, is called the *line integral* of the given function along the line in question.

If a given surface is divided in any way into infinitesimal elements such that the distance between the two most widely separated points within each element is infinitesimal, and the area of each element is multiplied by the value a given point-function, which is continuous over the surface, has at some point within the element, the limit approached by the sum of these products as each element is indefinitely decreased, is called the *surface integral* of the given function over the surface in question.

If a given space is divided in any way into infinitesimal elements such that the distance between the two most widely separated points within each element is infinitesimal, and the volume of each element is multiplied by the value a given point-function, which is continuous throughout the space, has at some point within the element, the limit approached by the sum of these products as each element is indefinitely decreased, is called the *space integral* of the given function throughout the space in question.

It is easily seen that the line integral of unity along a given line is the length of the line; that the surface integral of unity over a given surface is the area of the surface; and that the space integral of unity throughout a given space is the volume of the space.

In the chapter on Centres of Gravity we have had numerous simple examples of line, surface, and space integrals.

164. That the value of a line, surface, or space integral is independent of the position in each element of the point at which the value of the given function is taken can be proved as follows: The distance apart of any two points in the same infinitesimal element is infinitesimal (Art. 163), therefore the values of a continuous function taken at any two points in the same element will differ in general by an infinitesimal; the products obtained by multiplying these two values by the magnitude of the element will, then, differ by an infinitesimal of higher order than that of the element; therefore, in forming the integral either of these products may be used in place of the other without changing the result. (I. Art. 161.)

165. The line integral of a function along a given line is absolutely independent of the manner in which the line is broken up into infinitesimal elements, and is equal to the length of the line multiplied by the *mean value* of the function along the line; the *mean value* of the function being defined as follows: Suppose a set of points uniformly distributed along the

line, that is, so distributed that the number of points in any portion of the line is proportional to the length of the portion; take the value of the function at each of these points; divide the sum of these values by the number of the points; and the limit approached by this quotient as the number of the points is indefinitely increased is the *mean value* of the given function along the line; and this mean value is in general finite and determinate.

To prove our proposition, we have only to consider in detail the method of finding the mean value in question. Let the number of points in a unit of length of the line be  $k$ . Then, no matter how the line is broken up into infinitesimal elements, the number of points in each element is  $k$  times the length of the element. Since any two values of the function corresponding to points in the same element differ by an infinitesimal, in finding our limit we may replace all values corresponding to points in the same element by any one; hence the sum of the values corresponding to points in the same element may be replaced by one value multiplied by the number of points taken in that element, that is, this sum may be replaced by  $k$  times the product of one value by the length of the element; and the sum of the values corresponding to all the points taken in the line may be replaced by  $k$  times the sum of the terms obtained by multiplying the length of each element by the value of the function at some point within the element. When we divide this sum by the whole number of points considered, that is, by  $k$  times the length of the line, the  $k$ 's cancel out, and the required mean value reduces to the limit of the numerator divided by the length of the line, and the limit of the numerator is the line integral of the function along the line. Therefore the line integral is the mean value of the function multiplied by the length of the line.

The same proof may be given for a surface integral or for a space integral. The former is the product of the area of the surface by the mean value of the function over the surface; the latter is the volume of the space multiplied by the mean value of the function throughout the space; and both are inde-

pendent of the way in which the surface or space may be divided into infinitesimal elements.

166. If the line along which the integral is taken is a plane curve, it is easy to get a geometrical representation of the integral. For, if at every point of the line a perpendicular to the plane of the line is erected whose length is equal to the value of the function at the point, the line integral required clearly represents the area of the cylindrical surface containing the perpendiculars if the values are all of the same sign, and represents the difference of the areas of the portions of the cylindrical surface which lie on opposite sides of the line if the values of the function are not all of the same sign.

A similar construction shows that a surface integral over a plane surface may be represented by a volume or by the differences of volumes. Consequently, in each case if the function is finite and continuous, the integral is finite and determinate.

167. As examples of line, surface, and space integrals, we will calculate a few *moments of inertia*.

The *moment of inertia* of a body about a given axis may be defined as the space integral of the product of the density at any point of the body by the square of the distance of the point from the axis; the integral being taken throughout the space occupied by the body.

If the body considered is a material surface or a material line, the integral reduces to a surface integral or to a line integral.

In the examples taken below the body is supposed to be homogeneous.

(a) The moment of inertia of a circumference about a given diameter.

Using polar coördinates and taking the diameter as our axis,

$$\begin{aligned}
 I &= \int_0^{2\pi} a^2 \sin^2 \phi \cdot ka d\phi = ka^3 \pi \\
 &= \frac{1}{2} Ma^2,
 \end{aligned}
 \tag{1}$$



if  $I$  is the moment of inertia, and  $a$  the radius,  $k$  the density, and  $M$  the mass of the circumference in question.

(b) The moment of inertia of the perimeter of a square about an axis passing through the centre of the square and parallel to a side.

$$\begin{aligned} I &= 2 \int_{-a}^a y^2 k dy + 2 \int_{-a}^a a^2 k dx \\ &= \frac{4}{3} k a^3 + 4 k a^3 = \frac{16}{3} k a^3 \\ &= \frac{2}{3} M a^2, \end{aligned} \quad [2]$$

if  $2a$  is the length of a side.

(c) The moment of inertia of a circle about a diameter.

$$\begin{aligned} I &= \int_0^a \int_0^{2\pi} r^2 \sin^2 \phi \cdot k r d\phi dr = \frac{1}{4} k \pi a^4 \\ &= \frac{1}{4} M a^2. \end{aligned} \quad [3]$$

(d) The moment of inertia of a square about an axis through the centre of the square and parallel to a side.

$$\begin{aligned} I &= \int_{-a}^a \int_{-a}^a y^2 k dx dy = \frac{4}{3} k a^4 \\ &= \frac{1}{3} M a^2. \end{aligned} \quad [4]$$

(e) The moment of inertia of the surface of a sphere about a diameter.

$$\begin{aligned} I &= \int_0^{2\pi} \int_0^{\pi} a^2 \sin^2 \phi \cdot k a^2 \sin \phi d\phi d\theta = \frac{8}{3} k \pi a^4 \\ &= \frac{2}{3} M a^2. \end{aligned} \quad [5]$$

(f) The moment of inertia of the surface of a cube about an axis parallel to an edge and passing through the centre.

$$\begin{aligned} I &= 4 \int_{-a}^a \int_{-a}^a (a^2 + z^2) k dx dz + 2 \int_{-a}^a \int_{-a}^a (y^2 + z^2) k dy dz \\ &= \frac{6}{3} k a^4 + \frac{1}{3} k a^4 \\ &= \frac{10}{3} M a^2. \end{aligned} \quad [6]$$

(g) The moment of inertia of a sphere about a diameter.

$$I = \int_0^{2\pi} \int_0^{\pi} \int_0^a r^2 \sin^2 \phi \cdot kr^2 \sin \phi \, dr d\phi d\theta = \frac{8}{15} k\pi a^5$$

$$= \frac{2}{5} Ma^2. \quad [7]$$

(h) The moment of inertia of a cube about an axis through the centre and parallel to an edge.

$$I = \int_{-a}^a \int_{-a}^a \int_{-a}^a (y^2 + z^2) k \, dx dy dz = \frac{1}{3} ka^5$$

$$= \frac{2}{5} Ma^2. \quad [8]$$

EXAMPLES.

Find the moments of inertia of the following bodies :

(1) Of a straight line about a perpendicular through an extremity ; about a perpendicular through its middle point.

*Ans.*  $\frac{1}{3} Ml^2$  ;  $\frac{1}{12} Ml^2$ .

(2) Of the circumference of a circle about an axis through its centre perpendicular to its plane.

*Ans.*  $Ma^2$ .

(3) Of a circle about an axis through its centre perpendicular to its plane.

*Ans.*  $\frac{1}{2} Ma^2$ .

(4) Of a rectangle whose sides are  $2a$ ,  $2b$ , about an axis through its centre perpendicular to its plane ; about an axis through its centre parallel to the side  $2b$ .

*Ans.*  $\frac{1}{3} M(a^2 + b^2)$  ;  $\frac{1}{3} Ma^2$ .

(5) Of an ellipse about its major axis ; about its minor axis ; about an axis through the centre perpendicular to the plane of the ellipse.

*Ans.*  $\frac{1}{4} Mb^2$  ;  $\frac{1}{4} Ma^2$  ;  $\frac{1}{4} M(a^2 + b^2)$ .

(6) Of an ellipsoid about the axis  $a$ .

*Ans.*  $\frac{1}{5} M(b^2 + c^2)$ .

(7) Of a rectangular parallelepiped about an axis through the centre parallel to the edge  $2a$ .

*Ans.*  $\frac{1}{5} M(b^2 + c^2)$ .

(8) Of a segment of a parabola about the principal axis.


*Ans.*  $\frac{1}{5} Mb^2$ , where  $2b$  is the breadth of the segment.

168. If  $u$ ,  $D_x u$ , and  $D_y u$  are finite, continuous, and single-valued for all points in a given plane surface bounded by a closed curve  $T$ , the surface integral of  $D_x u$  taken over the surface is equal to the line integral of  $u \cos \alpha$  taken around the whole bounding curve, where  $\alpha$  is the angle made with the axis of  $X$  by the external normal at any point of the boundary.

This may be formulated thus :

$$\iint D_x u dx dy = \int u \cos \alpha . ds. \tag{1}$$

Let the axes be chosen so that the surface in question lies in the first quadrant, and divide the projection of  $T$  on the axis of  $Y$  into infinitesimal elements of which any one is  $dy$ .



On each of these elements as a base erect a rectangle ; and since  $T$  is a closed curve, each of these rectangles will cut it an even number of times.

Let us call the values of  $u$  at the points where the lower side of any one of these rectangles cuts  $T$ ,  $u_1, u_2, u_3, u_4$ , etc., respectively ; the angles which this side makes with the exterior normals at these points,  $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ , etc. ; and the elements which the rectangle cuts from  $T$ ,  $ds_1, ds_2, ds_3, ds_4$ , etc.

It is evident that whenever a line parallel to the axis of  $X$  cuts into the surface bounded by  $T$ , the corresponding value of  $\alpha$  is obtuse and its cosine negative ; that whenever it cuts out,

$\alpha$  is acute and its cosine positive; and that any value of  $\alpha$  is the angle which the contour  $T$  itself makes at the point in question with the axis of  $Y$  if we suppose the contour traced by a point moving so as to keep the bounded surface always on the left hand.

We have then approximately,

$$dy = -ds_1 \cdot \cos \alpha_1 = ds_2 \cdot \cos \alpha_2 = -ds_3 \cdot \cos \alpha_3 = ds_4 \cdot \cos \alpha_4 = \dots \quad [2]$$

If, now, in  $\iint D_x u dx dy$  we perform the integration with respect to  $x$ , and introduce the proper limits, we shall have

$$\iint D_x u dx dy = \int dy (-u_1 + u_2 - u_3 + u_4 \dots); \quad [3]$$

and the second member indicates that we are to form a quantity corresponding to that in parenthesis for every rectangle which cuts  $T$ , to multiply it by the base of the rectangle, and then to take the limit of the sum of the results as all the bases are indefinitely decreased.

By [2],

$$\begin{aligned} & dy (-u_1 + u_2 - u_3 + u_4 \dots) \\ &= u_1 \cos \alpha_1 ds_1 + u_2 \cos \alpha_2 ds_2 + u_3 \cos \alpha_3 ds_3 + u_4 \cos \alpha_4 ds_4 + \dots; \quad [4] \end{aligned}$$

and the limit of the sum of the values any one of which is represented by the second member of [4] is clearly  $\int u \cos \alpha ds$  taken around the whole of  $T$ .

EXAMPLE.

Prove that under the conditions stated in the last article

$$\iint D_x u dx dy = \int u \cos \beta \cdot ds,$$

where  $\beta$  is the angle made with the axis of  $Y$  by the exterior normal.

169. As an illustration of the last proposition, let us find the centre of gravity of a semicircle.

We have 
$$\bar{y} = \frac{k}{M} \int \int y dx dy. \quad (1)$$

But we may write  $y = D_x(xy)$ . Hence, by Art. 168,

$$\begin{aligned} \bar{y} &= \frac{k}{M} \int \int y dx dy = \frac{k}{M} \int xy \cos \alpha ds \\ &= \frac{k}{M} \left( \int_0^\pi a \cos \phi a \sin \phi \cos \phi a d\phi + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x \cdot 0 \cdot \cos \frac{\pi}{2} dx \right) \\ &= \frac{k}{k \frac{\pi a^2}{2}} \cdot \frac{2}{3} a^3 + 0 = \frac{4a}{3\pi}, \end{aligned}$$

which agrees with the result of Ex. 8, Art. 158.

As a second example, we shall find the moment of inertia of a circle about a diameter.

We have

$$\begin{aligned} I &= k \int \int y^2 dx dy = k \int xy^2 \cos \phi \cdot ds \\ &= k \int_0^{2\pi} a \cos \phi a^2 \sin^2 \phi \cos \phi a d\phi \\ &= k a^4 \int_0^{2\pi} \sin^2 \phi \cos^2 \phi d\phi = \frac{k}{4} \pi a^4 = \frac{1}{4} M a^2, \end{aligned}$$

which agrees with the result of (c), Art. 167.

#### EXAMPLES.

(1) Find the centre of gravity of a semicircle, using the theorem  $\int \int D_x u dx dy = \int u \cos \beta \cdot ds$ .

(2) Find the moment of inertia of a circle about an axis through its centre perpendicular to its plane, using the principles  $\int \int D_x u dx dy = \int u \cos \alpha \cdot ds$  and  $\int \int D_y u dx dy = \int u \cos \beta \cdot ds$ .

170. Since, as we have seen in Art. 168,  $\alpha$  is the angle which the curve  $T$  makes with the axis of  $Y$ ; if we trace the curve so as to keep the bounded space on our left, it follows that  $\cos \alpha \cdot ds = dy$ .

Hence 
$$\iint D_x u dx dy = \int u dy; \quad [1]$$

and in like manner,

$$\iint D_y u dx dy = - \int u dx; \quad [2]$$

the first integral in [1] and [2] being taken over the bounded surface, and the second around the bounding curve.

For example, the moment of inertia of a square about an axis through the centre and parallel to a side is

$$I = k \iint y^2 dx dy. \quad ((d) \text{ Art. 167.})$$

By [1], 
$$\iint y^2 dx dy = \int xy^2 dy,$$

and the last integral is to be taken around the perimeter.

Hence

$$\begin{aligned} I &= k \left[ \int_{-a}^a ay^2 dy + \int_a^{-a} (-ay^2 dy) \right] = 2ka \int_{-a}^a y^2 dy = \frac{2}{3} ka^3 \\ &= \frac{1}{3} Ma^2. \end{aligned}$$

**EXAMPLE.**


Work Ex. 8, Art. 167, by the aid of (2).

171. If  $U$ ,  $D_x U$ ,  $D_y U$ , and  $D_z U$  are finite, continuous, single-valued functions throughout the space bounded by a given closed surface  $T$ , the space integral of  $D_x U$  taken throughout the space in question is equal to the surface integral, taken over the bounding surface, of  $U \cos \alpha$ , where  $\alpha$  is the angle made with the axis of  $X$  by the exterior normal at any point of the surface.

This may be formulated thus :

$$\iiint D_x U dx dy dz = \int U \cos \alpha \cdot dS. \quad [1]$$

The proof is almost identical with that given in Art. 168, except that for elementary rectangle we use elementary prism. We shall merely indicate the steps.



$$dydz = -dS_1 \cos \alpha_1 = dS_2 \cos \alpha_2 = -dS_3 \cos \alpha_3 = \dots$$

$$\iiint D_x U dx dy dz = \iint dy dz [-U_1 + U_2 - U_3 \dots]$$

= the limit of the sum of terms of the form

$$U_1 \cos \alpha_1 \cdot dS_1 + U_2 \cos \alpha_2 \cdot dS_2 + U_3 \cos \alpha_3 \cdot dS_3 + \dots$$

$$= \int U \cos \alpha \cdot dS.$$

#### EXAMPLE.

Prove that under the conditions of the last article

$$\iiint D_y U dx dy dz = \int U \cos \beta \cdot dS,$$

and

$$\iiint D_x U dx dy dz = \int U \cos \gamma \cdot dS,$$

where  $\beta$  and  $\gamma$  are the angles made with the axes of  $Y$  and  $Z$  respectively by the exterior normal to the bounding surface.

172. As an illustration, let us find the centre of gravity of a hemisphere.

We have

$$\begin{aligned} \bar{x} &= \frac{k}{M} \iiint x dx dy dz = \frac{k}{M} \int \frac{x^2}{2} \cos \alpha \cdot dS \\ &= \frac{k}{2M} \int_0^{2\pi} \int_0^{\frac{\pi}{2}} a^2 \cos^2 \phi \cos \phi a^2 \sin \phi d\phi d\theta \\ &= \frac{a^4 k}{2M} \int_0^{2\pi} \int_0^{\frac{\pi}{2}} \cos^3 \phi \sin \phi d\phi d\theta \\ &= \frac{a^4 k}{\frac{1}{2} \pi a^3 k} \cdot \frac{\pi}{2} = \frac{3}{8} a; \end{aligned}$$

which agrees with the result of Art. 159.

EXAMPLE.

Find the moment of inertia of a sphere about a diameter ; of a cube about an axis through the centre parallel to an edge. Make your work depend upon finding the value of a surface integral.



## CHAPTER XV.

## MEAN VALUE AND PROBABILITY.

173. The application of the Integral Calculus to questions in Mean Value and Probability is a matter of decided interest; but lack of space will prevent our doing more than solving a few problems in illustration of some of the simplest of the methods and devices ordinarily employed. A full and admirable treatment of the subject is given in "Williamson's Integral Calculus" (London: Longmans, Green, & Co.); and numerous interesting problems are published with their solutions in "The Mathematical Visitor" and "The Annals of Mathematics."

174. The *mean* of  $n$  quantities is their sum divided by their number. If the number of quantities considered is supposed to increase indefinitely according to some given law, the problem of finding the limiting value approached by their mean usually calls for the Integral Calculus. The *mean value* of a continuous function of one, two, or three independent variables has been carefully defined in Art. 165, and has been proved to depend upon a line, surface, or space integral.

(a) Let us find the mean distance of all the points on the circumference of a circle from a given point on the circumference.

If we take the given point as origin, the distances whose mean is required are the radii vectores of points uniformly distributed along the circumference of the circle.

The required mean is, therefore, by Art. 165, equal to

---

the quotient obtained by dividing the line integral of  $r$  taken around the circumference by the length of the circumference; that is,

$$M = \frac{\int r ds}{2\pi a}.$$

The polar equation of the circle is

$$r = 2a \cos \phi;$$

$$ds = 2a d\phi,$$

$$M = \frac{1}{2\pi a} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 4a^2 \cos \phi d\phi = \frac{4a}{\pi},$$

the required *mean value*.

(b) Let us find the mean distance of points on the surface of a circle from a fixed point on the circumference.

Here, by Art. 165, the required mean is the surface integral of  $r$  taken over the circle, divided by the area of the circle; that is,

$$M = \frac{1}{\pi a^2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{2a \cos \phi} r^2 dr d\phi = \frac{32a}{9\pi}.$$

(c) The problem of finding the mean distance of points on the surface of a square from a corner of the square can be simplified slightly by considering merely one of the halves into which the square is divided by a diagonal.

Here

$$\begin{aligned} M &= \frac{2}{a^2} \int_0^{\frac{\pi}{4}} \int_0^{a \sec \phi} r \cdot r dr d\phi \\ &= \frac{a}{3} \left( \sqrt{2} + \log \tan \frac{3\pi}{8} \right). \end{aligned}$$

(d) As an example of a device often employed, we shall now solve the problem, To find the mean distance between two points within a given circle.

If  $M$  be the required mean, the sum of the whole number of cases can be represented by  $(\pi r^2)^2 M$ ,  $r$  being the radius of the circle; since for each position of the first point the number of positions of the second point is proportional to the area of the circle, and may be measured by that area; and as the number of possible positions of the first point may also be measured by the area of the circle, the whole number of cases to be considered is represented by the square of the area; and the sum of all the distances to be considered must be the product of the mean distance by the number.

Let us see what change will be produced in this sum by increasing  $r$  by the infinitesimal  $dr$ ; that is, let us find  $d(\pi^2 r^4 M)$ .

If the first point is anywhere on the annulus  $2\pi r.dr$ , which we have just added, its mean distance from the other points of the circle is  $\frac{32r}{9\pi}$ , by (b).

Therefore, the sum of the new distances to be considered, if the first point is on the annulus, is  $\frac{32r}{9\pi} \cdot \pi r^2 \cdot 2\pi r.dr$ ; but the second point may be on the annulus, instead of the first; so that to get the sum of all the new cases brought in by increasing  $r$  by  $dr$ , we must double the value just obtained.

$$\begin{aligned} \text{Hence} \quad d(\pi^2 r^4 M) &= \frac{128}{9} \pi r^4 dr, \\ \pi^2 r^4 M &= \frac{128}{9} \pi \int_0^a r^4 dr = \frac{128}{45} \pi a^5, \\ M &= \frac{128a}{45\pi}. \end{aligned}$$

175. In solving questions in *Probability*, we shall assume that the student is familiar with the elements of the theory as given in "Todhunter's Algebra."

(a) A man starts from the bank of a straight river, and walks till noon in a random direction; he then turns and walks

in another random direction ; what is the probability that he will reach the river by night?

Let  $\theta$  be the angle his first course makes with the river. If the angle through which he turns at noon is less than  $\pi - 2\theta$ , he will reach the river by night. For any given value of  $\theta$ , then, the required probability is  $\frac{\pi - 2\theta}{2\pi}$ . The probability that  $\theta$  shall lie between any given value  $\theta_0$  and  $\theta_0 + d\theta$  is  $\frac{d\theta}{\frac{1}{2}\pi}$ .

The chance that his first course shall make an angle with the river between  $\theta_0$  and  $\theta_0 + d\theta$ , and that he shall get back, is

$$\frac{\pi - 2\theta}{2\pi} \cdot \frac{d\theta}{\frac{1}{2}\pi} = \frac{(\pi - 2\theta)d\theta}{\pi^2}.$$

As  $\theta$  is equally likely to have any value between 0 and  $\frac{\pi}{2}$ , the required probability,

$$p = \int_0^{\frac{1}{2}\pi} \frac{(\pi - 2\theta)d\theta}{\pi^2} = \frac{1}{4}.$$

(b) A floor is ruled with equidistant straight lines ; a rod, shorter than the distance between the lines, is thrown at random on the floor ; to find the chance of its falling on one of the lines.

Let  $x$  be the distance of the centre of the rod from the nearest line ;  $\theta$  the inclination of the rod to a perpendicular to the parallels passing through the centre of the rod ;  $2a$  the common distance of the parallels ;  $2c$  the length of the rod.

In order that the rod may cross a line, we must have  $c \cos \theta > x$ ; the chance of this for any given value  $x_0$  of  $x$  is  $\frac{1}{\frac{1}{2}\pi} \cos^{-1} \frac{x_0}{c}$ .

The probability that  $x$  will have the value  $x_0$  is  $\frac{dx}{a}$ . The probability required is

$$p = \frac{2}{\pi a} \int_0^c \cos^{-1} \frac{x}{c} dx = \frac{2c}{\pi a}.$$

This problem may be solved by another method which possesses considerable interest.

Since all values of  $x$  from 0 to  $a$ , and all values of  $\theta$  from  $-\frac{\pi}{2}$  to  $\frac{\pi}{2}$  are equally probable, the whole number of cases that can arise may be represented by

$$\int_{-\frac{1}{2}\pi}^{\frac{1}{2}\pi} \int_0^a dx d\theta = \pi a.$$

The number of favorable cases will be represented by

$$\int_{-\frac{1}{2}\pi}^{\frac{1}{2}\pi} \int_0^{c \cos \theta} dx d\theta = 2c.$$

Hence

$$p = \frac{2c}{\pi a}.$$

(c) To find the probability that the distance of two stars, taken at random in the northern hemisphere, shall exceed  $90^\circ$ .

Let  $a$  be the latitude of the first star. With the star as a pole, describe an arc of a great circle, dividing the hemisphere into two lunes; the probability that the distance of the second star from the first will exceed  $90^\circ$  is the ratio of the lune not containing the first star to the hemisphere, and is equal to  $\frac{(\frac{1}{2}\pi - a)}{\pi}$ . The probability that the latitude of the first star will be between  $a$  and  $a + da$  is the ratio of the area of the zone, whose bounding circles have the latitudes  $a$  and  $a + da$  respectively, to the area of the hemisphere, and is

$$\frac{2\pi a^2 \cos a da}{2\pi a^2} = \cos a da.$$

Hence 
$$p = \int_0^{\frac{\pi}{2}} \frac{(\frac{1}{2}\pi - a)}{\pi} \cos a da = \frac{1}{\pi}.$$

(d) A random straight line meets a closed convex curve; what is the probability that it will meet a second closed convex curve within the first?

If an infinite number of random lines be drawn in a plane, all directions are equally probable; and lines having any given

direction will be disposed with equal frequency all over the plane. If we determine a line by its distance  $p$  from the origin, and by the angle  $\alpha$  which  $p$  makes with the axis of  $X$ , we can get all the lines to be considered by making  $p$  and  $\alpha$  vary between suitable limits by equal infinitesimal increments.

In our problem, the whole number of lines meeting the external curve can be represented by  $\int \int dp da$ . If the origin is within the curve, the limits for  $p$  must be zero, and the perpendicular distance from the origin to a tangent to the curve; and for  $\alpha$  must be zero and  $2\pi$ . If we call this number  $N$ , we shall have

$$N = \int_0^{2\pi} p da,$$

$p$  being now the perpendicular from the origin to the tangent.

If we regard the distance from a given point of any closed convex curve along the curve to the point of contact of a tangent, and then along the tangent to the foot of the perpendicular let fall upon it from the origin, as a function of the  $\alpha$  used above, its differential is easily seen to be  $p da$ . If we sum these differentials from  $\alpha = 0$  to  $\alpha = 2\pi$ , we shall get the perimeter of the given curve.

Hence 
$$N = \int_0^{2\pi} p da = L,$$

where  $L$  is the perimeter of the curve in question. By the same reasoning, we can see that  $n$ , the number of the random lines which meet the inner curve, is equal to  $l$ , its perimeter. For  $p$ , the required probability, we shall have

$$p = \frac{l}{L}.$$

#### EXAMPLES.

(1) A number  $n$  is divided at random into two parts; find the mean value of their product.

Ans.  $\frac{n^2}{6}$ .

(2) Find the mean value of the ordinates of a semicircle, supposing the series of ordinates taken equidistant. *Ans.*  $\frac{\pi}{4}a$ .

(3) Find the mean value of the ordinates of a semicircle, supposing the ordinates drawn through equidistant points on the circumference. *Ans.*  $\frac{2a}{\pi}$ .

(4) Find the mean values of the roots of the quadratic  $x^2 - ax + b = 0$ , the roots being known to be real, but  $b$  being unknown but positive. *Ans.*  $\frac{5a}{6}$  and  $\frac{a}{6}$ .

(5) Prove that the mean of the radii vectores of an ellipse, the focus being the origin, is equal to half the minor axis when they are drawn at equal angular intervals, and is equal to half the major axis when they are drawn so that the abscissas of their extremities increase uniformly.

(6) Suppose a straight line divided at random into three parts; find the mean value of their product. *Ans.*  $\frac{a^3}{60}$ .

(7) Find the mean square of the distance of a point within a given square (side =  $2a$ ) from the centre of the square. *Ans.*  $\frac{3}{4}a^2$ .

(8) A chord is drawn joining two points taken at random on a circumference; find the mean area of the less of the two segments into which it divides the circle. *Ans.*  $\frac{\pi a^2}{4} - \frac{a^2}{\pi}$ .

(9) Find the mean latitude of all places north of the equator. *Ans.*  $32^\circ.7$ .

(10) Find the mean distance of points within a sphere from a given point of the surface. *Ans.*  $\frac{3}{4}a$ .

(11) Find the mean distance of two points taken at random within a sphere. *Ans.*  $\frac{3}{8}a$ .

(12) Two points are taken at random in a given line  $a$ ; find the chance that their distance shall exceed a given value  $c$ .

$$\text{Ans. } \left(\frac{a-c}{a}\right)^2.$$

(13) Find the chance that the distance of two points within a square shall not exceed a side of the square. *Ans.*  $\pi - \frac{1}{6}$ .

(14) A line crosses a circle at random; find the chances that a point, taken at random within the circle, shall be distant from the line by less than the radius of the circle. *Ans.*  $1 - \frac{2}{3\pi}$ .

(15) A random straight line crosses a circle; find the chance that two points, taken at random in the circle, shall lie on opposite sides of the line. *Ans.*  $\frac{128}{45\pi^2}$ .

(16) A random straight line is drawn across a square; find the chance that it intersects two opposite sides. *Ans.*  $\frac{1}{2} - \frac{\log 2}{\pi}$ .

(17) Two arrows are sticking in a circular target; find the chance that their distance apart is greater than the radius. *Ans.*  $\frac{3\sqrt{3}}{4\pi}$ .

(18) From a point in the circumference of a circular field a projectile is thrown at random with a given velocity which is such that the diameter of the field is equal to the greatest range of the projectile: find the chance of its falling within the field. *Ans.*  $\frac{1}{2} - \frac{2}{\pi}(\sqrt{2} - 1)$ .

(19) On a table a series of equidistant parallel lines is drawn, and a cube is thrown at random on the table. Supposing that the diagonal of the cube is less than the distance between consecutive straight lines, find the chance that the cube will rest without covering any part of the lines.

*Ans.*  $1 - \frac{4a}{\pi c}$ , where  $a$  is the edge of the cube and  $c$  the distance between consecutive lines.

(20) A plane area is ruled with equidistant parallel straight lines, the distance between consecutive lines being  $c$ . A closed curve, having no singular points, whose greatest diameter is less



than  $c$ , is thrown down on the area. Find the chance that the curve falls on one of the lines.

Ans.  $\frac{l}{\pi c}$ , where  $l$  is the perimeter of the curve.

(21) During a heavy rain-storm, a circular pond is formed in a circular field. If a man undertakes to cross the field in the dark, what is the chance that he will walk into the pond?

## CHAPTER XVI.

## ELLIPTIC INTEGRALS.

176. In attempting to solve completely the problem of the motion of a simple pendulum by the methods of I. Chapter VIII. we encounter an integral of great importance which we have not yet considered. The problem is closely analogous to that of the Cycloidal pendulum (I. Art. 119).

For the sake of simplicity we shall suppose the pendulum bob to start from the lowest point of its circular path with the initial velocity that would be acquired by a particle falling freely in a vacuum through the distance  $y_0$ ; and this by I. Art. 114 [1] is  $\sqrt{2gy_0}$ .

Forming our differential equation of motion as in I. Art. 118, but taking the positive direction of the axis of  $Y$  upward, we have

$$\frac{d^2 s}{dt^2} = -g \frac{dy}{ds}. \quad (1)$$

Multiplying by  $2 \frac{ds}{dt}$  and integrating,

$$v^2 = \left( \frac{ds}{dt} \right)^2 = -2gy + C,$$

or, determining  $C$ ,

$$v^2 = \left( \frac{ds}{dt} \right)^2 = 2g(y_0 - y). \quad (2)$$

If the starting-point is taken as the origin, the equation of the circular path is  $x^2 + y^2 - 2ay = 0$ , whence

$$\left( \frac{ds}{dt} \right)^2 = \frac{a^2}{2ay - y^2} \left( \frac{dy}{dt} \right)^2,$$

and we have

$$\frac{a}{\sqrt{2ay - y^2}} \frac{dy}{dt} = \sqrt{2g(y_0 - y)},$$

or 
$$dt = \frac{ady}{\sqrt{2g} \cdot \sqrt{(y_0 - y)(2ay - y^2)}}.$$

Integrating, and determining the arbitrary constant, we get

$$t = \frac{a}{\sqrt{2g}} \int_0^y \frac{dy}{\sqrt{(y_0 - y)(2ay - y^2)}} \quad (3)$$

as the time required to reach that point of the path which has the ordinate  $y$ .

The substitution of  $x^2 = \frac{y}{y_0}$  reduces (3) to the form

$$t = \sqrt{\frac{a}{g}} \int_0^x \frac{dx}{\sqrt{(1 - x^2) \left(1 - \frac{y_0}{2a} x^2\right)}}, \quad (4)$$

where the integral is of the form

$$\int_0^x \frac{dx}{\sqrt{(1 - x^2)(1 - k^2 x^2)}}, \quad (5)$$

$k^2$  being positive and less than unity if  $y_0$  is less than  $2a$ . An examination of equation (2) will show that if this is true, the pendulum will oscillate between the two points of the arc which have the ordinate  $y_0$ .

If  $y_0$  is greater than  $2a$ , the pendulum will make complete revolutions. For this case the substitution of  $x^2 = \frac{y}{2a}$  in (3) will reduce it to

$$t = a \sqrt{\frac{2}{gy_0}} \int_0^x \frac{dx}{\sqrt{(1 - x^2) \left(1 - \frac{2a}{y_0} x^2\right)}}, \quad (6)$$

where the integral is of the form (5),  $k^2$  being positive and less than unity.

The time required for the pendulum to reach its greatest height — that is, in the first case, the time of a half-vibration, and in the second case, the time of a half-revolution — will depend upon

$$\int_0^1 \frac{dx}{\sqrt{(1 - x^2)(1 - k^2 x^2)}}. \quad (7)$$

177. The length of an arc of an Ellipse, measured from the extremity of the minor axis, has been found to be (Art. 107)

$$s = \int_0^x \sqrt{\frac{a^2 - e^2 x^2}{a^2 - x^2}} \cdot dx. \quad (1)$$

If we replace  $\frac{x}{a}$  by  $x$ , (1) becomes

$$s = a \int_0^x \sqrt{\frac{1 - e^2 x^2}{1 - x^2}} \cdot dx, \quad (2)$$

and the integral is of the form

$$\int_0^x \sqrt{\frac{1 - k^2 x^2}{1 - x^2}} \cdot dx, \quad (3)$$

where  $k^2$  is positive and less than unity.

The length of an Elliptic quadrant depends upon the integral

$$\int_0^1 \sqrt{\frac{1 - k^2 x^2}{1 - x^2}} \cdot dx. \quad (4)$$

178. It can be shown by an elaborate investigation, for which we have not room, that the integral of any algebraic function, which is irrational through containing under the square root sign an algebraic polynomial of the third or fourth degree, can by suitable transformations be made to depend upon one or more of the three integrals

$$F(k, x) = \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \quad [1]$$

$$E(k, x) = \int_0^x \sqrt{\frac{1-k^2x^2}{1-x^2}} \cdot dx, \quad [2]$$

$$\Pi(n, k, x) = \int_0^x \frac{dx}{(1+nx^2)\sqrt{(1-x^2)(1-k^2x^2)}}, \quad [3]$$

which are known as the Elliptic Integrals of the first, second, and third class respectively.

$k$ , which may always be taken positive and less than 1, is called the *modulus*; and  $n$ , which may be taken real, is called the *parameter* of the integral.

$$K = F(k, 1) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}, \quad [4]$$

and  $E = E(k, 1) = \int_0^1 \sqrt{\frac{1-k^2x^2}{1-x^2}} \cdot dx, \quad [5]$

are known as the Complete Elliptic Integrals of the first and second classes.

179. The substitution of  $x = \sin \phi$  in the Elliptic Integrals reduces them to the following simpler forms.

$$F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1-k^2\sin^2\phi}} = \int_0^\phi \frac{d\phi}{\Delta\phi}. \quad [1]$$

$$E(k, \phi) = \int_0^\phi \sqrt{1-k^2\sin^2\phi} \cdot d\phi = \int_0^\phi \Delta\phi \cdot d\phi. \quad [2]$$

$$\Pi(n, k, \phi) = \int_0^\phi \frac{d\phi}{(1+n\sin^2\phi)\sqrt{1-k^2\sin^2\phi}} = \int_0^\phi \frac{d\phi}{(1+n\sin^2\phi)\Delta\phi}. \quad [3]$$

$$K = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-k^2\sin^2\phi}} = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi}. \quad [4]$$

$$E = \int_0^{\frac{\pi}{2}} \sqrt{1-k^2\sin^2\phi} \cdot d\phi = \int_0^{\frac{\pi}{2}} \Delta\phi \cdot d\phi. \quad [5]$$

$\phi$  is called the *amplitude* of the Elliptic Integral, and  $\Delta\phi = \sqrt{1-k^2\sin^2\phi}$  is called the *delta* of  $\phi$ , or more simply, *delta*  $\phi$ , and is regarded as a new trigonometric function: it is always taken with the positive sign, and has an analogy with  $\cos \phi$ .

For a given value of  $k$ ,  $\Delta\phi$  is easily seen to be a periodic function of  $\phi$  having the period  $\pi$ . It has its maximum value 1

when  $\phi = 0$  and when  $\phi = \pi$ , and its minimum value  $\sqrt{1-k^2}$ , which is usually represented by  $k'$  and called the *complementary modulus*, when  $\phi = \frac{\pi}{2}$ ; and  $\Delta\left(\frac{\pi}{2} + a\right) = \Delta\left(\frac{\pi}{2} - a\right)$ .

*Landen's Transformation.*

180. The approximate numerical value of an Elliptic Integral of the first class, when  $k$  and  $\phi$  are given, is easily computed by the aid of two valuable reduction formulas due to *Landen*.

If in 
$$F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}}$$

we replace  $\phi$  by  $\phi_1$ ,  $\phi_1$  and  $\phi$  being connected by the relation

$$\tan \phi = \frac{\sin 2\phi_1}{k + \cos 2\phi_1}, \quad (1)$$

which is easily transformable into either of the following :

$$k \sin \phi = \sin (2\phi_1 - \phi), \quad (2)$$

$$\tan (\phi - \phi_1) = \frac{1-k}{1+k} \tan \phi_1, \quad (3)$$

$$\int_0^\phi \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} \text{ reduces to } \frac{2}{1+k} \int_0^{\phi_1} \frac{d\phi_1}{\sqrt{1 - \frac{4k}{(1+k)^2} \sin^2 \phi_1}},$$

which is also an Elliptic Integral of the first class, but has a different modulus and a different amplitude from those of the given integral.

The steps of the process are as follows :

From (1) we easily find

$$\sin^2 \phi = \frac{\sin^2 2\phi_1}{1+k^2+2k \cos 2\phi_1},$$

whence 
$$\sqrt{1-k^2 \sin^2 \phi} = \frac{1+k \cos 2\phi_1}{\sqrt{1+k^2+2k \cos 2\phi_1}}.$$

Differentiating (1), we get

$$\sec^2 \phi d\phi = \frac{2(1+k \cos 2\phi_1)}{(k + \cos 2\phi_1)^2} d\phi_1;$$

but from (1),  $\sec^2 \phi = \frac{1+k^2+2k \cos 2\phi_1}{(k + \cos 2\phi_1)^2};$

hence  $d\phi = \frac{2(1+k \cos 2\phi_1)}{1+k^2+2k \cos 2\phi_1} d\phi_1,$

$$\begin{aligned} \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} &= \frac{2 d\phi_1}{\sqrt{1+k^2+2k \cos 2\phi_1}} = \frac{2 d\phi_1}{\sqrt{1+k^2+2k-4k \sin^2 \phi_1}} \\ &= \frac{2}{1+k} \frac{d\phi_1}{\sqrt{1-\frac{4k}{(1+k)^2} \sin^2 \phi_1}}. \end{aligned}$$

$$\phi_1 = 0 \quad \text{when} \quad \phi = 0,$$

hence  $\int_0^\phi \frac{d\phi}{\sqrt{1-k^2 \sin^2 \phi}} = \frac{2}{1+k} \int_0^{\phi_1} \frac{d\phi_1}{\sqrt{1-\frac{4k}{(1+k)^2} \sin^2 \phi_1}},$

and  $F(k, \phi) = \frac{2}{1+k} F(k_1, \phi_1),$

where  $k_1 = \frac{2\sqrt{k}}{1+k},$

and  $\sin(2\phi_1 - \phi) = k \sin \phi.$

} [4]

$k_1$  is less than 1 and greater than  $k$ ; for  $\frac{2\sqrt{k}}{1+k} < 1$  reduces to  $0 < (1-\sqrt{k})^2$ , which is obviously true, and  $\frac{2\sqrt{k}}{1+k} > k$  reduces to  $4 > k(1+k)^2$ , which is true, since  $k$  is less than 1. If  $\phi$  is not greater than  $\frac{\pi}{2}$ , and the smallest value of  $\phi_1$  consistent with the relation  $\sin(2\phi_1 - \phi) = k \sin \phi$  is taken,  $0 < \phi_1 < \phi$ . Hence (4) is a reduction formula by which we can raise the modulus and lower the amplitude of our given function.

By applying the formula (4)  $n$  times, we get

$$F(k, \phi) = \frac{2}{1+k} \cdot \frac{2}{1+k_1} \cdot \frac{2}{1+k_2} \cdots \frac{2}{1+k_{n-1}} F(k_n, \phi_n);$$

or, since  $\frac{2}{1+k} = \frac{k_1}{\sqrt{k}}$ , etc.,

$$\left. \begin{aligned} F(k, \phi) &= k_n \sqrt{\frac{k_1 \cdot k_2 \cdot k_3 \cdots k_{n-1}}{k}} F(k_n, \phi_n), \\ \text{where } k_p &= \frac{2\sqrt{k_{p-1}}}{1+k_{p-1}}, \text{ and } \sin(2\phi_p - \phi_{p-1}) = k_{p-1} \sin \phi_{p-1}. \end{aligned} \right\} [5]$$

If we suppose  $n$  in (5) to be indefinitely increased, we shall have  $\lim_{n \rightarrow \infty} [k_n] = 1$ ; for if we form the series

$$(1-k) + (1-k_1) + (1-k_2) + \cdots + (1-k_p) + \cdots,$$

we shall have

$$\frac{1-k_{p+1}}{1-k_p} = \frac{1 - \frac{2\sqrt{k_p}}{1+k_p}}{1-k_p} = \frac{(1-\sqrt{k_p})^2}{1-k_p^2} = \frac{1-\sqrt{k_p}}{1+\sqrt{k_p}} \frac{1}{1+k_p},$$

which is always less than unity; hence the terms in the series must decrease indefinitely as  $p$  is increased and  $\lim_{n \rightarrow \infty} [1-k_n] = 0$ . Since, as we have seen above,  $\phi_n$  continually diminishes as  $n$  increases, but does not reach the value zero, it must have some limiting value  $\Phi$ . Hence

$$\begin{aligned} \lim_{n \rightarrow \infty} [F(k_n, \phi_n)] &= F(1, \Phi) = \int_0^\Phi \frac{d\phi}{\sqrt{1-\sin^2\phi}} \\ &= \int_0^\Phi \sec \phi d\phi = \log \tan \left[ \frac{\pi}{4} + \frac{\Phi}{2} \right]; \end{aligned}$$

$$\text{and } F(k, \phi) = \log \tan \left[ \frac{\pi}{4} + \frac{\Phi}{2} \right] \sqrt{\frac{k_1 \cdot k_2 \cdot k_3 \cdots}{k}}. \quad [6]$$

Formulas [5] and [6] lend themselves very readily to numerical computation.



181. Formula [4], Art. 180, may be used to decrease the modulus and increase the amplitude of a given Elliptic Integral. Interchanging the subscripts, and using (3) Art. 180 instead of (2) Art. 180, we have

$$\left. \begin{aligned} F(k, \phi) &= \frac{1+k_1}{2} F(k_1, \phi_1), \\ \text{where} \quad k_1 &= \frac{1-\sqrt{1-k^2}}{1+\sqrt{1-k^2}}, \\ \text{and} \quad \tan(\phi_1 - \phi) &= \sqrt{1-k^2} \tan \phi. \end{aligned} \right\} [1]$$

By repeated application of [1] we get

$$\left. \begin{aligned} F(k, \phi) &= (1+k_1)(1+k_2)\dots(1+k_n) \frac{F(k_n, \phi_n)}{2^n}, \\ \text{where} \quad k_p &= \frac{1-\sqrt{1-k_{p-1}^2}}{1+\sqrt{1-k_{p-1}^2}}, \\ \text{and} \quad \tan(\phi_p - \phi_{p-1}) &= \sqrt{1-k_{p-1}^2} \tan \phi_{p-1}. \end{aligned} \right\} [2]$$

It is easily shown, as in Art. 180, that  $\lim_{n \rightarrow \infty} [k_n] = 0$ , and consequently that  $\lim_{n \rightarrow \infty} F(k_n, \phi_n) = \int_0^\Phi d\phi = \Phi$ , where  $\Phi$  is the limiting value approached by  $\phi$  as  $n$  is increased.

If  $\phi = \frac{\pi}{2}$ , we get from [2],  $\phi_1 = \pi$ ,  $\phi_2 = 2\pi$ , ...  $\phi_n = 2^{n-1}\pi$ ;  
hence  $K = F\left(k, \frac{\pi}{2}\right) = \frac{\pi}{2} (1+k_1)(1+k_2)(1+k_3)\dots$  [3]

Formulas [2] and [3], like formulas [5] and [6] of Art. 180, lend themselves readily to computation.

With a large modulus, it is generally best to use [5] and [6] of Art. 180; with a small modulus, [2] or [3] of the present article will generally work more rapidly.

We give in the next article the whole work of computing the Elliptic Integral  $F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$  by each of the two methods, and

of computing  $K\left(\frac{\sqrt{2}}{2}\right) = F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right)$  by the second method, employing five-place logarithms.

182.  $F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$ . METHOD OF ART. 180.

$$k = 0.70712 \qquad \log k = 9.84949$$

$$1 + k = 1.70712 \qquad \log(1 + k) = 0.23226$$

$$\log \sqrt{k} = 9.92474$$

$$\log 2 = 0.30103$$

$$\text{colog}(1 + k) = \underline{9.76774}$$

$$\log k_1 = 9.99351$$

$$k_1 = 0.98518 \qquad \log k_1 = 9.99351$$

$$1 + k_1 = 1.98518 \qquad \log(1 + k_1) = 0.29780$$

$$\log \sqrt{k_1} = 9.99676$$

$$\log 2 = 0.30103$$

$$\text{colog}(1 + k_1) = \underline{9.70220}$$

$$\log k_2 = 9.99999$$

$$\underline{k_2 = 1}$$

$$\log k = 9.84949$$

$$\log \sin \frac{\pi}{4} = \underline{9.84949}$$

$$\log \sin(2\phi_1 - \phi) = \underline{9.69898}$$

$$2\phi_1 - \phi = 30^\circ \ 0' \ 3''$$

$$2\phi_1 = 75^\circ \ 0' \ 3''$$

$$\phi_1 = 37^\circ \ 30' \ 2''$$

$$\log k_1 = 9.99351$$

$$\log \sin \phi_1 = \underline{9.78445}$$

$$\log \sin(2\phi_2 - \phi_1) = \underline{9.77796}$$

$$2\phi_2 - \phi_1 = 36^\circ 51' 3''$$

$$2\phi_2 = 74^\circ 21' 5''$$

$$\Phi = \phi_2 = 37^\circ 10' 32''$$

$$\frac{1}{2}\Phi + \frac{\pi}{4} = 63^\circ 35' 16''$$

$$\log \tan\left(\frac{\pi}{4} + \frac{1}{2}\Phi\right) = 0.30393$$


---

$$\log \sqrt{k_1} = 9.99676$$

$$\operatorname{colog} \sqrt{k} = 0.07526$$

$$\log \log \tan\left(\frac{\pi}{4} + \frac{1}{2}\Phi\right) = 9.48277$$

$$\operatorname{colog} \mu = 0.36222$$


---

$$\log F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right) = 9.91701$$

$$F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right) = 0.82605$$


---

$\mu = 0.43429$  is the modulus of the common system of logarithms.

$$F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right). \text{ METHOD OF ART. 181.}$$

$$\sqrt{1 - k^2} = k' = 0.70712$$

$$1 - k' = 0.29288$$

$$1 + k' = 1.70712$$

$$k_1 = 0.17157$$

$$1 - k_1 = 0.82843$$

$$1 + k_1 = 1.17157$$

$$k_1' = 0.98520$$

$$1 - k_1' = 0.01480$$

$$1 + k_1' = 1.98520$$

$$k_2 = 0.00746$$

$$\log(1 - k') = 9.46669$$

$$\operatorname{colog}(1 + k') = 9.76774$$

$$\log k_1 = 9.23443$$

$$\log(1 - k_1) = 9.91826$$

$$\log(1 + k_1) = 0.06878$$

$$\log k_1'^2 = 9.98704$$

$$\log k_1' = 9.99352$$

$$\log(1 - k_1') = 8.17026$$

$$\operatorname{colog}(1 + k_1') = 9.70220$$

$$\log k_2 = 7.87246$$

$$1 - k_2 = 0.99254 \qquad \log(1 - k_2) = 9.99675$$

$$1 + k_2 = 1.00746 \qquad \log(1 + k_2) = 0.00323$$

$$\log k_2'^2 = 9.99998$$

$$k_2' = 1$$

$$\log k_2' = 9.99999$$

$$\underline{k_2 = 0}$$

$$\log k' = 9.84949$$

$$\log \tan \phi = 0.00000$$

$$\log \tan(\phi_1 - \phi) = 9.84949$$

$$\phi_1 - \phi = 35^\circ 15' 53''$$

$$\phi_1 = 80^\circ 15' 53''$$

$$\log k_1' = 9.99352$$

$$\log \tan \phi_1 = 0.76557$$

$$\log \tan(\phi_2 - \phi_1) = 0.75909$$

$$\phi_2 - \phi_1 = 80^\circ 7' 17''$$

$$\phi_2 = 160^\circ 23' 10''$$

$$\tan(\phi_3 - \phi_2) = \tan \phi_2$$

$$\Phi = \phi_3 = 2\phi_2 = 320^\circ 46' 20''$$

$$\frac{1}{2^3}\Phi = 40^\circ 5' 48''$$

$$= 144348''$$

$$\pi = 648000''$$

$$\log\left(\frac{1}{2^3}\Phi\right)'' = 5.15942$$

$$\text{colog } \pi'' = 4.18842$$

$$\log \pi = 0.49715$$

$$\underline{\log\left(\frac{1}{2^3}\Phi\right) = 9.84499}$$

$$\begin{aligned}\log(1+k_1) &= 0.06878 \\ \log(1+k_2) &= 0.00323 \\ \log\left(\frac{1}{2^3}\Phi\right) &= 9.84499 \\ \log F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right) &= 9.91700 \\ F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right) &= 0.82605\end{aligned}$$

For  $F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right)$  we have by (3), Art. 181,

$$\begin{aligned}\log(1+k_1) &= 0.06878 \\ \log(1+k_2) &= 0.00323 \\ \log \pi &= 0.49715 \\ \operatorname{colog} 2 &= 9.69897 \\ \log F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right) &= 0.26813 \\ F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right) &= 1.8541\end{aligned}$$

183. Landen's Transformation can also be applied to the computation of Elliptic Integrals of the second class, but the task is a more difficult one; we shall, however, give a brief sketch of the method; and in so doing we shall apply it to a more general form

$$G(k, \phi) = \int_0^\phi \frac{\alpha + \beta \sin^2 \phi}{\sqrt{1 - k^2 \sin^2 \phi}} d\phi, \quad [1]$$

of which  $E(k, \phi)$  is a special case.

From Art. 180 we have

$$\begin{aligned}\sqrt{1 - k^2 \sin^2 \phi} &= \frac{1 + k \cos 2\phi_1}{\sqrt{1 + k^2 + 2k \cos 2\phi_1}}, \\ \cos \phi &= \frac{k + \cos 2\phi_1}{\sqrt{1 + k^2 + 2k \cos 2\phi_1}}, \\ d\phi &= \frac{2(1 + k \cos 2\phi_1)}{1 + k^2 + 2k \cos 2\phi_1} d\phi_1.\end{aligned}$$

Hence  $\sqrt{1 - k^2 \sin^2 \phi} + k \cos \phi = \sqrt{1 + k^2 + 2k \cos 2\phi_1}$ .

$$\begin{aligned} G(k, \phi) &= \int_0^\phi \frac{\alpha + \beta \sin^2 \phi}{\sqrt{1 - k^2 \sin^2 \phi}} d\phi \\ &= \int_0^\phi \left[ \frac{\alpha}{\sqrt{1 - k^2 \sin^2 \phi}} + \frac{\beta}{k^2} \cdot \frac{1 - (1 - k^2 \sin^2 \phi)}{\sqrt{1 - k^2 \sin^2 \phi}} \right] d\phi \\ &= \int_0^\phi \left[ \frac{\alpha + \frac{\beta}{k^2}}{\sqrt{1 - k^2 \sin^2 \phi}} - \frac{\beta}{k^2} \sqrt{1 - k^2 \sin^2 \phi} \right] d\phi, \end{aligned}$$

and

$$\begin{aligned} G(k, \phi) - \frac{\beta}{k} \sin \phi &= \int_0^\phi \left[ \frac{\alpha + \frac{\beta}{k^2}}{\sqrt{1 - k^2 \sin^2 \phi}} - \frac{\beta}{k^2} (\sqrt{1 - k^2 \sin^2 \phi} + k \cos \phi) \right] d\phi. \end{aligned}$$

Substituting  $\phi_1$  for  $\phi$ , this becomes

$$\begin{aligned} G(k, \phi) - \frac{\beta}{k} \sin \phi &= 2 \int_0^{\phi_1} \frac{\alpha - \frac{\beta}{k} \cos 2\phi_1}{\sqrt{1 + k^2 + 2k \cos 2\phi_1}} d\phi_1 \\ &= \frac{2}{1+k} \int_0^{\phi_1} \frac{\alpha - \frac{\beta}{k} + \frac{2\beta}{k} \sin^2 \phi_1}{\sqrt{1 - \frac{4k}{(1+k)^2} \sin^2 \phi_1}} d\phi_1. \end{aligned}$$

Hence  $G(k, \phi) = \frac{\beta}{k} \sin \phi + \frac{2}{1+k} G_1(k_1, \phi_1)$ , [2]

where

$$k_1 = \frac{2\sqrt{k}}{1+k}, \quad \sin(2\phi_1 - \phi) = k \sin \phi, \quad \alpha_1 = \alpha - \frac{\beta}{k}, \quad \beta_1 = \frac{2\beta}{k}. \quad [3]$$

Formulas [2] and [3] enable us to make our given function depend upon one of the same form, but having a greater modulus and a less amplitude. A repeated use of [2], together with the reductions employed in Art. 180, gives us

$$\begin{aligned} \therefore z \cdot \phi &= \frac{\beta}{k} \sin \phi + \frac{\beta}{k_1 k_2} \sin \phi_1 + \sqrt{\frac{k_1}{k}} \cdot \beta_2 \sin \phi_2 \\ &+ \sqrt{\frac{k_1 k_2}{k}} \cdot \beta_3 \sin \phi_3 + \dots + \sqrt{\frac{k_1 k_2 \dots k_{n-1}}{k}} \beta_{n-1} \sin \phi_{n-1} \\ &+ k_n \sqrt{\frac{k_1 k_2 \dots k_n}{k}} \cdot G_n(k_n, \phi_n), \end{aligned} \quad [4]$$

where  $\beta_2 = \frac{\beta_3}{k_1 k_2 \dots k_{n-1}}$

and  $\phi_n = a - \frac{\beta}{k} \left( 1 + \frac{\beta}{k_1} + \frac{\beta^2}{k_1 k_2} + \dots + \frac{\beta^{n-1}}{k_1 k_2 k_3 \dots k_{n-1}} \right)$

Just as in Art. 180  $k_n$  rapidly approaches 1 as  $n$  is increased; the limiting value of  $G_n(k_n, \phi_n)$  is then

$$\begin{aligned} \text{limit } G_n(k_n, \phi_n) &= \int_0^{\phi_n} \frac{\alpha + \beta \sin^2 \phi}{\cos \phi} d\phi \\ &= (\alpha + \beta) \log \tan \left( \frac{\pi}{4} + \frac{\phi_n}{2} \right) - \beta \sin \phi_n. \end{aligned} \quad [6]$$

By Art. 180, [5] and [6],

$$\text{limit } k_n \sqrt{\frac{k_1 k_2 \dots k_n}{k}} \log \tan \left[ \frac{\pi}{4} + \frac{\phi_n}{2} \right] = F(k, \phi).$$

[4] can thus be written

$$\begin{aligned} G(k, \phi) &= F(k, \phi) \left[ \alpha - \frac{\beta}{k} \left( 1 + \frac{\beta}{k_1} + \frac{\beta^2}{k_1 k_2} + \dots \right. \right. \\ &\left. \left. + \frac{\beta^{n-1}}{k_1 k_2 \dots k_{n-1}} - \frac{\beta^n}{k_1 k_2 k_3 \dots k_{n-1}} \right) \right] \\ &+ \frac{\beta}{k} \left[ \sin \phi + \frac{\beta}{\sqrt{k}} \sin \phi_1 + \frac{\beta^2}{\sqrt{k} k_1} \sin \phi_2 + \frac{\beta^3}{\sqrt{k} k_1 k_2} \sin \phi_3 + \dots \right. \\ &\left. + \frac{\beta^{n-1}}{\sqrt{k} k_1 \dots k_{n-1}} \sin \phi_{n-1} - \frac{\beta^n}{\sqrt{k} k_1 \dots k_{n-1}} \sin \phi_n \right]. \end{aligned} \quad [7]$$

If  $\alpha = 1$ , and  $\beta = -k^2$ , [7] reduces to

$$\begin{aligned}
 E(k, \phi) = F(k, \phi) & \left[ 1 + k \left( 1 + \frac{2}{k_1} + \frac{2^2}{k_1 k_2} + \dots \right. \right. \\
 & \left. \left. + \frac{2^{n-1}}{k_1 k_2 \dots k_{n-1}} - \frac{2^n}{k_1 k_2 \dots k_{n-1}} \right) \right] \\
 & - k \left[ \sin \phi + \frac{2}{\sqrt{k}} \sin \phi_1 + \frac{2^2}{\sqrt{k} k_1} \sin \phi_2 + \dots \right. \\
 & \left. + \frac{2^{n-1}}{\sqrt{k} k_1 \dots k_{n-2}} \sin \phi_{n-1} - \frac{2^n}{\sqrt{k} k_1 \dots k_{n-1}} \sin \phi_n \right], \quad [8]
 \end{aligned}$$

where  $k_p = \frac{2\sqrt{k_{p-1}}}{1+k_{p-1}}$ , and  $\sin(2\phi_p - \phi_{p-1}) = k_{p-1} \sin \phi_{p-1}$ . [9]

By Formulas [8] and [9] an Elliptic Integral of the Second Class may be computed without difficulty.

184. Formula [2], Art. 183, may be used to decrease the modulus and increase the amplitude of an Elliptic Integral. Interchanging the subscripts, we have

$$G(k, \phi) = \frac{1+k_1}{2} \left[ G_1(k_1, \phi_1) - \frac{\beta_1}{k_1} \sin \phi_1 \right];$$

or, since  $\frac{\beta_1}{k_1} = \frac{\beta}{2}$ , (Art. 183 [3]),

$$G(k, \phi) = \frac{1+k_1}{2} \left[ G_1(k_1, \phi_1) - \frac{\beta}{2} \sin \phi_1 \right], \quad [1]$$

where

$$k_1 = \frac{1-\sqrt{1-k^2}}{1+\sqrt{1-k^2}}, \quad \tan(\phi_1 - \phi) = \sqrt{1-k^2} \tan \phi, \quad \alpha_1 = \alpha + \frac{\beta}{2}, \quad \beta_1 = \frac{k_1 \beta}{2}. \quad [2]$$



A repeated use of [2] gives

$$G(k, \phi) = \frac{1-k}{2} \cdot \frac{1-k_1}{2} \cdot \frac{1-k_2}{2} \cdots \frac{1-k_n}{2} G_n(k, \phi) \\ - \frac{1-k_1}{2} \frac{1-k_2}{2} \sin \phi_1 - \frac{1-k_2}{2} \frac{1-k_3}{2} \sin \phi_2 + \cdots \\ + \frac{1-k_1}{2} \cdot \frac{1-k_2}{2} \cdots \frac{1-k_n}{2} \sin \phi_n \quad [3]$$

where  $\sin \phi_n = \frac{1-k_1 k_2 \cdots k_n}{2^n}$

and  $\sin \phi_{n-1} = \frac{1}{2} \left( 1 - \frac{1-k_1}{2} - \frac{1-k_2}{2} - \frac{k_1 k_2}{2^2} - \cdots + \frac{k_1 k_2 \cdots k_{n-1}}{2^{n-1}} \right)$

Just as in Art. 131, limit  $k_n = 0$ , therefore limit  $G_n = 0$  and  
 limit  $G(k, \phi) = \int_0^{\sin \phi} 1 \cdot d\phi = \sin \phi$ .

By Art. 131, [2],  $\frac{1-k_1}{2} \cdot \frac{1-k_2}{2} \cdots \frac{1-k_n}{2} \sin \phi_n = F(k, \phi)$ .

By Art. 130, [1],  $\frac{1-k_1}{2} = \frac{1-\sqrt{k_1}}{k_1}$ ,  $\frac{1-k_2}{2} = \frac{1-\sqrt{k_2}}{k_2}$ , etc.

Hence [3] becomes

$$G(k, \phi) = F(k, \phi) \left[ 1 + \frac{3}{2} \left( 1 - \frac{k_1}{2} - \frac{k_1 k_2}{2^2} - \frac{k_1 k_2 k_3}{2^3} - \cdots \right) \right] \\ - \frac{\beta \sqrt{k_1 k_2}}{k_1 - 2} \sin \phi_1 + \frac{\beta \sqrt{k_2 k_3}}{2^2} \sin \phi_2 + \frac{\beta \sqrt{k_3 k_4}}{2^3} \sin \phi_3 + \cdots \quad [4]$$

If  $\alpha = 1$  and  $\beta = -k^2$ , [4] reduces to

$$E(k, \phi) = F(k, \phi) \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} + \frac{k_1 k_2 k_3}{2^3} + \dots \right) \right] \\ + k \left[ \frac{\sqrt{k_1}}{2} \sin \phi_1 + \frac{\sqrt{k_1 k_2}}{2^2} \sin \phi_2 + \frac{\sqrt{k_1 k_2 k_3}}{2^3} \sin \phi_3 + \dots \right], \quad [5]$$

where

$$k_p = \frac{1 - \sqrt{1 - k_{p-1}^2}}{1 + \sqrt{1 - k_{p-1}^2}}, \text{ and } \tan(\phi_p - \phi_{p-1}) = \sqrt{1 - k_{p-1}^2} \cdot \tan \phi_{p-1}. \quad [6]$$

We have seen in Art. 181 that if  $\phi = \frac{\pi}{2}$ ,  $\phi_p = 2^{p-1}\pi$ .

Therefore, for a complete Elliptic Integral of the second class we have

$$E\left(k, \frac{\pi}{2}\right) = F\left(k, \frac{\pi}{2}\right) \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} + \frac{k_1 k_2 k_3}{2^3} + \dots \right) \right]. \quad [7]$$

Formulas [5] and [7] are admirably adapted to computation.

We give in the next article the work of computing  $E\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$  by each of the methods just given, and of computing  $E\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right)$  by the second method; using, as far as possible, the values already employed or obtained in Art. 182.

185.  $E\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right)$ . METHOD OF ART. 183.

Here, as we have seen in Art. 182, if we carry the work only to five decimal places,  $k_2 = 1$ , and our working formula will be

$$E(k, \phi) = F(k, \phi) \left[ 1 + k \left( 1 - \frac{2}{k_1} \right) \right] \\ - k \left[ \sin \phi + \frac{2}{\sqrt{k}} \sin \phi_1 - \frac{2^2}{\sqrt{k} k_1} \sin \phi_2 \right].$$

$$\begin{aligned} \log 2 &= 0.30103 \\ \log k &= 9.84949 \\ \text{colog } k_1 &= 0.00649 \\ \hline &0.15701 \end{aligned}$$

$$\frac{2k}{k_1} = 1.43553$$

$$1 + k = 1.70712$$

$$\log \left( 1 + k - \frac{2k}{k_1} \right) = 9.43391$$

$$1 + k - \frac{2k}{k_1} = 0.27159$$

$$\log F \left( \frac{\sqrt{2}}{2}, \frac{\pi}{4} \right) = \frac{9.91701}{9.35092}$$

$$F \left( \frac{\sqrt{2}}{2}, \frac{\pi}{4} \right) \left( 1 + k - \frac{2k}{k_1} \right) = 0.22435$$

$$\begin{aligned} \log k &= 9.84949 \\ \log \sin \phi &= 9.84949 \\ \hline &9.69898 \end{aligned}$$

$$k \sin \phi = 0.5$$

$$\begin{aligned} \log 2 &= 0.30103 \\ \log \sqrt{k} &= 9.92474 \\ \log \sin \phi_1 &= 9.78445 \\ \hline &0.01022 \end{aligned}$$

$$\frac{2k}{\sqrt{k}} \sin \phi_1 = 1.0238$$

$$\begin{aligned} \log 2^2 &= 0.60206 \\ \log \sqrt{k} &= 9.92474 \\ \text{colog } \sqrt{k_1} &= 0.00324 \\ \log \sin \phi_2 &= 9.78122 \\ \hline &0.31126 \end{aligned}$$

$$\frac{2^2 k}{\sqrt{k} k_1} \sin \phi_2 = 2.0477$$

$$-k \left( \sin \phi + \frac{2}{\sqrt{k}} \sin \phi_1 - \frac{2^2}{\sqrt{k} k_1} \sin \phi_2 \right) = 0.5239$$

$$F \left( \frac{\sqrt{2}}{2}, \frac{\pi}{4} \right) \left( 1 + k - \frac{2k}{k_1} \right) = 0.22435$$

$$E \left( \frac{\sqrt{2}}{2}, \frac{\pi}{4} \right) = 0.74825$$

$$E\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right). \text{ METHOD OF ART. 184.}$$

$k_3 = 0$ . Therefore our formula is

$$E(k, \phi) = F(k, \phi) \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} \right) \right] + k \left( \frac{\sqrt{k_1}}{2} \sin \phi_1 + \frac{\sqrt{k_1 k_2}}{2^2} \sin \phi_2 \right).$$

$$\log k_1 = 9.23443$$

$$\log k_2 = 7.87246$$

$$\text{colog } 4 = 9.39794$$

$$\hline 6.50483$$

$$\frac{k_1 k_2}{2^2} = 0.00032$$

$$\frac{k_1}{2} = 0.08578$$

$$\hline 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} = 1.08610$$

$$\frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} \right) = 0.271525$$

$$\log 0.728475 = 9.862415 \quad 1 - \frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} \right) = 0.728475$$

$$\log F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right) = 9.91700$$

$$\hline 9.779415$$

$$F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{4}\right) (0.728475) = 0.60178$$

$$\log k = 9.84949$$

$$\log \sqrt{k_1} = 9.61722$$

$$\text{colog } 2 = 9.69897$$

$$\log \sin \phi_1 = 9.99370$$

$$\hline 9.15938$$

$$\frac{k\sqrt{k_1}}{2} \sin \phi_1 = 0.14434$$

$$\begin{aligned}
 \log k &= 9.84949 \\
 \log \sqrt{k_1} &= 9.61722 \\
 \log \sqrt{k_2} &= 8.93623 \\
 \text{colog } 4 &= 9.39794 \\
 \log \sin \phi_2 &= 9.52592 \\
 \hline
 &= 7.32680
 \end{aligned}$$

$$\frac{k \sqrt{k_1 k_2}}{2^2} \sin \phi_2 = 0.00212$$

$$k \left( \frac{\sqrt{k_1}}{2^2} \sin \phi_1 + \frac{\sqrt{k_1 k_2}}{2^2} \sin \phi_2 \right) = 0.14646$$

$$F \left( \frac{\sqrt{2}}{2}, \frac{\pi}{4} \right) (0.728475) = 0.60178$$

$$E \left( \frac{\sqrt{2}}{2}, \frac{\pi}{4} \right) = 0.74824$$

$$E \left( \frac{\sqrt{2}}{2}, \frac{\pi}{2} \right). \quad \text{METHOD OF ART. 184.}$$

$$E \left( k, \frac{\pi}{2} \right) = F \left( k, \frac{\pi}{2} \right) \left[ 1 - \frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} \right) \right].$$

$$1 - \frac{k^2}{2} \left( 1 + \frac{k_1}{2} + \frac{k_1 k_2}{2^2} \right) = 0.728475 \quad \log 0.728475 = 9.862415$$

$$\log F \left( \frac{\sqrt{2}}{2}, \frac{\pi}{2} \right) = 0.26813$$

$$E \left( \frac{\sqrt{2}}{2}, \frac{\pi}{2} \right) = 1.3507 \quad \log E \left( \frac{\sqrt{2}}{2}, \frac{\pi}{2} \right) = 0.13054$$

186. An Elliptic Integral of the first or second class, whose amplitude is greater than  $\frac{\pi}{2}$ , can be made to depend upon one whose amplitude is less than  $\frac{\pi}{2}$ , and upon the corresponding Complete Elliptic Integral.

We have

$$F(k, \pi) = \int_0^{\pi} \frac{d\phi}{\Delta\phi} = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} + \int_{\frac{\pi}{2}}^{\pi} \frac{d\phi}{\Delta\phi} = K + \int_{\frac{\pi}{2}}^{\pi} \frac{d\phi}{\Delta\phi}, \text{ by [4], Art. 179.}$$

In  $\int_{\frac{\pi}{2}}^{\pi} \frac{d\phi}{\Delta\phi}$  let  $\phi = \pi - \psi$ ;

then  $d\phi = -d\psi$  and  $\Delta\phi = \sqrt{1 - k^2 \sin^2 \phi} = \sqrt{1 - k^2 \sin^2 \psi} = \Delta\psi$ ,

and we have  $\int_{\frac{\pi}{2}}^{\pi} \frac{d\phi}{\Delta\phi} = -\int_{\frac{\pi}{2}}^0 \frac{d\psi}{\Delta\psi} = -\int_{\frac{\pi}{2}}^0 \frac{d\phi}{\Delta\phi} = \int_0^{\frac{\pi}{2}} \frac{d\phi}{\Delta\phi} = K$ .

Hence  $F(k, \pi) = \int_0^{\pi} \frac{d\phi}{\Delta\phi} = 2K$ . [1]

$$F(k, n\pi + \rho) = \int_0^{n\pi + \rho} \frac{d\phi}{\Delta\phi} \\ = \int_0^{\pi} \frac{d\phi}{\Delta\phi} + \int_{\pi}^{2\pi} \frac{d\phi}{\Delta\phi} + \int_{2\pi}^{3\pi} \frac{d\phi}{\Delta\phi} + \dots + \int_{(p-1)\pi}^{p\pi} \frac{d\phi}{\Delta\phi} + \dots + \int_{n\pi}^{n\pi + \rho} \frac{d\phi}{\Delta\phi}.$$

In  $\int_{p\pi}^{(p+1)\pi} \frac{d\phi}{\Delta\phi}$  let  $\phi = p\pi + \psi$ ; then  $d\phi = d\psi$ , and  $\Delta\phi = \Delta\psi$ ,

and we have  $\int_{p\pi}^{(p+1)\pi} \frac{d\phi}{\Delta\phi} = \int_0^{\pi} \frac{d\psi}{\Delta\psi} = \int_0^{\pi} \frac{d\phi}{\Delta\phi} = 2K$ .

The substitution of  $\psi$  for  $\phi - n\pi$  in  $\int_{n\pi}^{n\pi+\rho} \frac{d\phi}{\Delta\phi}$  gives us

$$\int_{n\pi}^{n\pi+\rho} \frac{d\phi}{\Delta\phi} = \int_0^{\rho} \frac{d\psi}{\Delta\psi} = \int_0^{\rho} \frac{d\phi}{\Delta\phi} = F(k, \rho).$$

$$\text{Therefore} \quad F(k, n\pi + \rho) = 2nK + F(k, \rho). \quad [2]$$

In like manner it can be proved that

$$F(k, n\pi - \rho) = 2nK - F(k, \rho), \quad [3]$$

$$E(k, n\pi + \rho) = 2nE + E(k, \rho), \quad [4]$$

$$E(k, n\pi - \rho) = 2nE - E(k, \rho), \quad [5]$$

where  $E = E\left(k, \frac{\pi}{2}\right)$  is the complete Elliptic Integral of the second class.

A table giving the values of the Elliptic Integrals of the first and second classes for values of the amplitude between 0 and  $\frac{\pi}{2}$  is, then, a complete table.

Such a table, carried out to ten decimal places, is given by Legendre in his "Traité des Fonctions Elliptiques." We give in the next article a small three-place table.

It must be noted that the first column gives  $F(0, \phi)$  and  $E(0, \phi)$ , that is,  $\int_0^{\phi} d\phi = \phi$ ; and that the last column gives  $F(1, \phi)$  and  $E(1, \phi)$ , that is,  $\log \tan\left(\frac{\pi}{4} + \frac{\phi}{2}\right)$  and  $\sin \phi$ .

The complete Elliptic Integrals,

$$K = F\left(k, \frac{\pi}{2}\right) \text{ and } E = E\left(k, \frac{\pi}{2}\right),$$

are given in the last line of each table.

$F(k, \phi)$ .

$\phi$	$k=0$ $\sin 0^\circ$	$k=0.1$ $\sin 6^\circ$	$k=0.2$ $\sin 12^\circ$	$k=0.3$ $\sin 18^\circ$	$k=0.4$ $\sin 24^\circ$	$k=0.5$ $\sin 30^\circ$	$k=0.6$ $\sin 37^\circ$	$k=0.7$ $\sin 45^\circ$	$k=0.8$ $\sin 53^\circ$	$k=0.9$ $\sin 64^\circ$	$k=1.$ $\sin 90^\circ$
$0^\circ$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
5	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087
10	0.175	0.175	0.175	0.175	0.175	0.175	0.175	0.175	0.175	0.175	0.175
$15^\circ$	0.262	0.262	0.262	0.262	0.262	0.263	0.263	0.263	0.264	0.264	0.265
20	0.349	0.349	0.349	0.350	0.350	0.351	0.352	0.353	0.354	0.355	0.356
25	0.436	0.436	0.437	0.438	0.439	0.440	0.441	0.443	0.445	0.448	0.451
$30^\circ$	0.524	0.524	0.525	0.526	0.527	0.529	0.532	0.536	0.539	0.544	0.549
35	0.611	0.611	0.612	0.614	0.617	0.620	0.624	0.630	0.636	0.644	0.653
40	0.698	0.699	0.700	0.703	0.707	0.712	0.718	0.727	0.736	0.748	0.763
$45^\circ$	0.785	0.786	0.789	0.792	0.798	0.804	0.814	0.826	0.839	0.858	0.881
50	0.873	0.874	0.877	0.882	0.889	0.898	0.911	0.928	0.947	0.974	1.011
55	0.960	0.961	0.965	0.972	0.981	0.993	1.010	1.034	1.060	1.099	1.154
$60^\circ$	1.047	1.049	1.054	1.062	1.074	1.090	1.112	1.142	1.178	1.233	1.317
65	1.134	1.137	1.143	1.153	1.168	1.187	1.215	1.254	1.302	1.377	1.506
70	1.222	1.224	1.232	1.244	1.262	1.285	1.320	1.370	1.431	1.534	1.735
$75^\circ$	1.309	1.312	1.321	1.336	1.357	1.385	1.426	1.488	1.566	1.703	2.028
80	1.396	1.400	1.410	1.427	1.452	1.485	1.534	1.608	1.705	1.885	2.436
85	1.484	1.487	1.499	1.519	1.547	1.585	1.643	1.731	1.848	2.077	3.131
$k. 90^\circ$	1.571	1.575	1.588	1.610	1.643	1.686	1.752	1.854	1.993	2.275	$\infty$



E(k, φ).

φ	k = 0		k = 0.1		k = 0.2		k = 0.3		k = 0.4		k = 0.5		k = 0.6		k = 0.7		k = 0.8		k = 0.9		k = 1.		
	sin 0°	sin 0°	sin 0°	sin 0°	sin 12°	sin 12°	sin 18°	sin 18°	sin 24°	sin 24°	sin 30°	sin 30°	sin 37°	sin 37°	sin 46°	sin 46°	sin 53°	sin 53°	sin 64°	sin 64°	sin 90°	sin 90°	
0°	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
5	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087	0.087
10	0.175	0.175	0.175	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174	0.174
15°	0.262	0.262	0.262	0.262	0.262	0.262	0.262	0.262	0.261	0.261	0.261	0.261	0.261	0.261	0.260	0.260	0.260	0.260	0.259	0.259	0.259	0.259	0.259
20	0.349	0.349	0.349	0.349	0.349	0.349	0.349	0.349	0.348	0.348	0.347	0.347	0.347	0.347	0.346	0.346	0.345	0.345	0.343	0.343	0.342	0.342	0.342
25	0.436	0.436	0.436	0.436	0.436	0.436	0.435	0.435	0.434	0.434	0.433	0.433	0.433	0.433	0.430	0.430	0.428	0.428	0.425	0.425	0.423	0.423	0.423
30°	0.524	0.523	0.523	0.523	0.523	0.521	0.521	0.521	0.520	0.518	0.518	0.515	0.515	0.515	0.512	0.512	0.509	0.509	0.505	0.505	0.500	0.500	0.500
35	0.611	0.610	0.610	0.609	0.609	0.607	0.607	0.607	0.605	0.602	0.602	0.598	0.598	0.598	0.593	0.593	0.588	0.588	0.581	0.581	0.574	0.574	0.574
40	0.698	0.698	0.698	0.696	0.696	0.693	0.693	0.693	0.690	0.685	0.685	0.679	0.679	0.679	0.672	0.672	0.664	0.664	0.654	0.654	0.643	0.643	0.643
45°	0.785	0.785	0.785	0.782	0.779	0.779	0.779	0.779	0.773	0.767	0.767	0.759	0.759	0.759	0.748	0.748	0.737	0.737	0.723	0.723	0.707	0.707	0.707
50	0.873	0.872	0.869	0.864	0.864	0.864	0.864	0.864	0.857	0.848	0.848	0.837	0.837	0.837	0.823	0.823	0.808	0.808	0.789	0.789	0.766	0.766	0.766
55	0.960	0.959	0.955	0.948	0.948	0.948	0.948	0.948	0.939	0.928	0.928	0.914	0.914	0.914	0.895	0.895	0.875	0.875	0.850	0.850	0.819	0.819	0.819
60°	1.047	1.046	1.041	1.032	1.032	1.032	1.032	1.032	1.021	1.008	1.008	0.989	0.989	0.989	0.965	0.965	0.940	0.940	0.907	0.907	0.866	0.866	0.866
65	1.134	1.132	1.126	1.116	1.116	1.116	1.116	1.116	1.103	1.086	1.086	1.063	1.063	1.063	1.033	1.033	1.001	1.001	0.960	0.960	0.906	0.906	0.906
70	1.222	1.219	1.212	1.200	1.200	1.200	1.200	1.200	1.184	1.163	1.163	1.135	1.135	1.135	1.099	1.099	1.060	1.060	1.008	1.008	0.940	0.940	0.940
75°	1.309	1.306	1.297	1.283	1.283	1.283	1.283	1.283	1.264	1.240	1.240	1.207	1.207	1.207	1.163	1.163	1.117	1.117	1.053	1.053	0.966	0.966	0.966
80	1.396	1.393	1.383	1.367	1.367	1.367	1.367	1.367	1.344	1.316	1.316	1.277	1.277	1.277	1.227	1.227	1.172	1.172	1.095	1.095	0.985	0.985	0.985
85	1.484	1.480	1.468	1.450	1.450	1.450	1.450	1.450	1.424	1.392	1.392	1.347	1.347	1.347	1.289	1.289	1.225	1.225	1.135	1.135	0.996	0.996	0.996
E. 90°	1.571	1.566	1.554	1.533	1.533	1.533	1.533	1.533	1.504	1.467	1.467	1.417	1.417	1.417	1.351	1.351	1.278	1.278	1.173	1.173	1.000	1.000	1.000

*Addition Formulas.*

188. The Elliptic Integrals,  $F(k, x)$  and  $E(k, x)$ , may be regarded as new functions of  $x$ , defined by the aid of definite integrals; namely,

$$F(k, x) = \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}},$$

$$E(k, x) = \int_0^x \sqrt{\frac{1-k^2x^2}{1-x^2}} \cdot dx;$$

see Art. 178, [1] and [2].

We have seen how we may compute their values to any required degree of approximation when  $k$  and  $x$  are given. It remains to study their properties.

We are familiar with other and much simpler functions which may be defined as definite integrals, and whose most important properties can be deduced from these definitions.

For example, we may define  $\log x$  as  $\int_1^x \frac{dx}{x}$ ,  $\sin^{-1}x$  as  $\int_0^x \frac{dx}{\sqrt{1-x^2}}$ ,  $\tan^{-1}x$  as  $\int_0^x \frac{dx}{1+x^2}$ , and the theory of these functions may be based upon these definitions. For instance, the fundamental property of the logarithm is expressed by what is called the *addition formula*,

$$\log x + \log y = \log (xy),$$

and the whole theory of logarithms may be based on this property; and there are addition formulas for the other functions defined above; namely,

$$\sin^{-1}x + \sin^{-1}y = \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2}),$$

$$\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right).$$

These three important formulas are usually obtained by more or less elaborate methods involving the theory of the functions which are the inverse or anti-functions of the  $\log x$ , the  $\sin^{-1} x$ , and the  $\tan^{-1} x$ , that is, of  $e^x$ ,  $\sin x$ , and  $\tan x$ ; but they may be obtained without difficulty from the definitions of  $\log x$ ,  $\sin^{-1} x$ , and  $\tan^{-1} x$ , as definite integrals.

Take first 
$$\log x = \int_1^x \frac{dx}{x}.$$

Let us determine  $y$  in terms of  $x$ , so that

$$\log x + \log y = \log c, \quad (1)$$

where  $c$  is a given constant.

Since 
$$\log y = \int_1^y \frac{dy}{y},$$

if we differentiate (1), we have

$$\frac{dx}{x} + \frac{dy}{y} = 0,$$

or 
$$ydx + xdy = 0. \quad (2)$$

Integrate (2), and we get

$$\int ydx + \int xdy = C. \quad (3)$$

Simplify the first member of (3) by *integration by parts*;

$$xy - \int xdy + xy - \int ydx = C,$$

or 
$$2xy - \int (xdy + ydx) = C.$$

Reducing by the aid of (2),  $2xy = C,$

or 
$$xy = C_1, \quad (4)$$

where  $C_1$  is an undetermined constant. To determine  $C_1$ , let  $x = 1$  in (4), and we have  $y = C_1$  when  $x = 1$ ; let  $x = 1$  in (1), then  $\log x = \int_1^1 \frac{dx}{x} = 0$ ,  $\log y = \log c$ , and  $y = c$ , when  $x = 1$ . Therefore  $C_1 = c$ , and  $xy = c$ . Consequently  $y = \frac{c}{x}$  is the required value of  $y$ , and we have (1)

$$\log x + \log \frac{c}{x} = \log c.$$

We can express this relation more neatly by replacing  $c$  by its value  $xy$ , and thus we reach our required addition formula

$$\log x + \log y = \log(xy). \quad [5]$$

189. The addition formula for the  $\sin^{-1}$  can be deduced in exactly the same way. We wish to determine  $y$  so that

$$\sin^{-1}x + \sin^{-1}y = \sin^{-1}c. \quad (1)$$

We have  $\sin^{-1}x = \int_0^x \frac{dx}{\sqrt{1-x^2}}$ ,  $\sin^{-1}y = \int_0^y \frac{dy}{\sqrt{1-y^2}}$ .

Differentiate (1).

$$\frac{dx}{\sqrt{1-x^2}} + \frac{dy}{\sqrt{1-y^2}} = 0, \quad (2)$$

or  $\sqrt{1-y^2} \cdot dx + \sqrt{1-x^2} \cdot dy = 0$ ,

$$\int \sqrt{1-y^2} \cdot dx + \int \sqrt{1-x^2} \cdot dy = C.$$

Integrate by parts, and

$$x\sqrt{1-y^2} + y\sqrt{1-x^2} + \int xy \left( \frac{dx}{\sqrt{1-x^2}} + \frac{dy}{\sqrt{1-y^2}} \right) = C;$$

or, reducing by (2),

$$x\sqrt{1-y^2} + y\sqrt{1-x^2} = C. \quad (3)$$

To determine  $C$ , we have from (3)  $y = C$  when  $x = 0$ , and from (1)  $y = c$  when  $x = 0$ , since  $\sin^{-1}x = \int_0^x \frac{dx}{\sqrt{1-x^2}} = 0$ , when  $x = 0$ . Hence  $C = c$ , and  $x\sqrt{1-y^2} + y\sqrt{1-x^2} = c$ , and, finally,

$$\sin^{-1}x + \sin^{-1}y = \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2}). \quad [4]$$

To get an addition formula for the  $\tan^{-1}$ , a slight device is required, that of dividing the differential equation corresponding to (2) by  $1 - x^2y^2$ .

As before, let

$$\tan^{-1}x + \tan^{-1}y = \tan^{-1}c, \quad (5)$$

where  $\tan^{-1}x = \int_0^x \frac{dx}{1+x^2},$

and  $\tan^{-1}y = \int_0^y \frac{dy}{1+y^2}.$

$$\frac{dx}{1+x^2} + \frac{dy}{1+y^2} = 0,$$

or  $(1+y^2)dx + (1+x^2)dy = 0. \quad (6)$

Divide by  $1 - x^2y^2$  and integrate.

$$\int \frac{1+y^2}{1-x^2y^2} \cdot dx + \int \frac{1+x^2}{1-x^2y^2} \cdot dy = C.$$

Integrate by parts. We have

$$d \cdot \frac{1+y^2}{1-x^2y^2} = \frac{2y}{(1-x^2y^2)^2} [(1+x^2)dy + xy(1+y^2)dx],$$

$$d \cdot \frac{1+x^2}{1-x^2y^2} = \frac{2x}{(1-x^2y^2)^2} [(1+y^2)dx + xy(1+x^2)dy],$$

and

$$x \cdot \frac{1+y^2}{1-x^2y^2} + y \cdot \frac{1+x^2}{1-x^2y^2} = \frac{x+y}{1-xy}.$$

Hence

$$\frac{x+y}{1-xy} - \int \frac{2xy(1+xy)}{(1-x^2y^2)^2} [(1+y^2)dx + (1+x^2)dy] = C.$$

Therefore, by (6),  $\frac{x+y}{1-xy} = C.$  (7)

To determine  $C$ , we have from (7)  $y=C$  when  $x=0$ , and from (5)  $y=c$  when  $x=0$ , since  $\tan^{-1}x = \int_0^x \frac{dx}{1-x^2} = 0$  when  $x=0$ .

Hence  $C=c$ , and  $\frac{x+y}{1-xy} = c$ ,

and, finally,  $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right).$  [8]

190. To get an addition formula for  $F(k, x)$ , as before let  $F(k, x) + F(k, y) = F(k, c)$ , (1)

where  $F(k, x) = \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$ ,

and  $F(k, y) = \int_0^y \frac{dy}{\sqrt{(1-y^2)(1-k^2y^2)}}$

$$\frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \frac{dy}{\sqrt{(1-y^2)(1-k^2y^2)}} = 0, \quad (2)$$

or

$$\sqrt{(1-y^2)(1-k^2y^2)} \cdot dx + \sqrt{(1-x^2)(1-k^2x^2)} \cdot dy = 0. \quad (3)$$

Divide by  $1 - k^2x^2y^2$  and integrate.

$$\int \frac{\sqrt{(1-y^2)(1-k^2y^2)}}{1-k^2x^2y^2} \cdot dx + \int \frac{\sqrt{(1-x^2)(1-k^2x^2)}}{1-k^2x^2y^2} \cdot dy = C.$$

Integrate by parts. We have

$$d \cdot \frac{\sqrt{(1-y^2)(1-k^2y^2)}}{1-k^2x^2y^2} = \frac{y}{(1-k^2x^2y^2)^2} \{ [2k^2(x^2+y^2) - (1+k^2)(1+k^2x^2y^2)] \frac{dy}{\sqrt{(1-y^2)(1-k^2y^2)}} + 2k^2xy\sqrt{(1-y^2)(1-k^2y^2)} \cdot dx \};$$

$$d \cdot \frac{\sqrt{(1-x^2)(1-k^2x^2)}}{1-k^2x^2y^2} = \frac{x}{(1-k^2x^2y^2)^2} \{ [2k^2(x^2+y^2) - (1+k^2)(1+k^2x^2y^2)] \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + 2k^2xy\sqrt{(1-x^2)(1-k^2x^2)} \cdot dy \}.$$

Hence

$$\begin{aligned} & \frac{x\sqrt{(1-y^2)(1-k^2y^2)} + y\sqrt{(1-x^2)(1-k^2x^2)}}{1-k^2x^2y^2} \\ & - \int \frac{xy}{(1-k^2x^2y^2)^2} \{ [2k^2(x^2+y^2) - (1+k^2)(1+k^2x^2y^2)] \\ & \left[ \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} + \frac{dy}{\sqrt{(1-y^2)(1-k^2y^2)}} \right] \\ & + 2k^2xy[\sqrt{(1-y^2)(1-k^2y^2)} \cdot dx \\ & + \sqrt{(1-x^2)(1-k^2x^2)} \cdot dy] \} = C. \end{aligned}$$

Reducing, by the aid of (2) and (3), we have

$$\frac{x\sqrt{(1-y^2)(1-k^2y^2)} + y\sqrt{(1-x^2)(1-k^2x^2)}}{1-k^2x^2y^2} = C. \quad (4)$$

To determine  $C$ , from (4)  $y = C$  when  $x = 0$ , and from (1)  $y = c$  when  $x = 0$ . Therefore  $C = c$ , and we get

$$\begin{aligned}
 &F(k, x) + F(k, y) \\
 &= F\left(k, \frac{x\sqrt{(1-y^2)(1-k^2y^2)} + y\sqrt{(1-x^2)(1-k^2x^2)}}{1-k^2x^2y^2}\right), \quad [5]
 \end{aligned}$$

our required addition formula.

An addition formula for  $E(k, x)$  can be obtained in very much the same way, but the work is rather complicated, and it is better to use a method which will be explained later.

THE ELLIPTIC FUNCTIONS.

191. We have just seen that there is an analogy between the Elliptic Integral  $F(k, x)$ , and the familiar functions  $\log x$ ,  $\sin^{-1}x$ , and  $\tan^{-1}x$ ; and we know that the theory of these functions is ultimately connected with that of their inverse functions,  $\log^{-1}u$  or  $e^u$ ,  $\sin u$ , and  $\tan u$ ; and, indeed, that the latter are so much simpler than the former that it is customary to regard them as the direct functions, and the logarithm, the anti-sine, and the anti-tangent as the inverse functions.

For example: the first three addition formulas just obtained are much simpler when we express them in terms of the direct functions, and they become

$$\begin{aligned}
 &\log^{-1}(u + v) = \log^{-1}u + \log^{-1}v, \\
 \text{or} \quad &e^{(u+v)} = e^u \cdot e^v, \quad [1]
 \end{aligned}$$

$$\begin{aligned}
 &\sin(u + v) = \sin u \sqrt{1 - \sin^2 v} + \sin v \sqrt{1 - \sin^2 u}; \\
 \text{or} \quad &\sin(u + v) = \sin u \cos v + \cos u \sin v, \quad [2]
 \end{aligned}$$

$$\tan(u + v) = \frac{\tan u + \tan v}{1 - \tan u \cdot \tan v}; \quad [3]$$

and in this form they seem to better deserve the name of addition formulas.



In the same way the addition formula for  $F(k, x)$  can be more simply written in terms of the function which we might naturally represent by  $F^{-1}u \pmod{k}$ ; and, as we might expect, this function has many interesting and important properties which well deserve investigation.

Since in most of the work which follows we shall generally employ the same modulus throughout, we shall not take the trouble to write it except in the few cases where its omission might give rise to confusion, and then we shall put  $\pmod{k}$  after the function, as above with  $F^{-1}u \pmod{k}$ .

192. In Arts. 178 and 179 we have adopted two forms of notation for an Elliptic Integral of the first class,  $F(k, x)$  and  $F(k, \phi)$ ;

$$F(k, x) = \int_0^x \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}},$$

$$F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1-k^2\sin^2\phi}} = \int_0^\phi \frac{d\phi}{\Delta\phi},$$

where

$$x = \sin \phi, \quad \sqrt{1-x^2} = \cos \phi,$$

and

$$\sqrt{1-k^2x^2} = \sqrt{1-k^2\sin^2\phi} = \Delta\phi.$$

If we let

$$u = F(k, x) = F(k, \phi),$$

we have in Art. 179 called  $\phi$  the amplitude of  $u$ , and  $\sin \phi$ ,  $\cos \phi$ , and  $\Delta\phi$  may be called the sine, the cosine, and the delta of the amplitude of  $u$ ; and  $\phi$ ,  $\sin \phi$ ,  $\cos \phi$ , and  $\Delta\phi$  may be written  $\text{am } u$ ,  $\text{sinam } u$ ,  $\text{cosam } u$ , and  $\Delta \text{am } u$ , or, more briefly,  $\text{am } u$ ,  $\text{sn } u$ ,  $\text{cn } u$ , and  $\text{dn } u$ ; and may be read amplitude  $u$ , sine amplitude  $u$ , cosine amplitude  $u$ , and delta amplitude  $u$ . Formulating, we have

$$\left. \begin{aligned} u &= F(k, x) = F(k, \phi), \\ \phi &= \text{am } u, \\ x &= \sin \phi = \text{sn } u, \\ \sqrt{1-x^2} &= \cos \phi = \text{cn } u, \\ \sqrt{1-k^2x^2} &= \Delta\phi = \text{dn } u, \end{aligned} \right\} \quad [1]$$

$\operatorname{sn} u$ ,  $\operatorname{cn} u$ ,  $\operatorname{dn} u$ , are trigonometric functions of  $\phi$ , the amplitude of  $u$ , but they may be regarded as new and somewhat complicated functions of  $u$  itself, and from this point of view they are called *Elliptic Functions* of  $u$ .

$\operatorname{am} u$  also is sometimes called an Elliptic Function; and there are various allied functions that are sometimes included under the general title of Elliptic Functions. We shall, however, restrict the name to  $\operatorname{sn} u$ ,  $\operatorname{cn} u$ , and  $\operatorname{dn} u$ . They have an analogy with trigonometric functions, and have a theory which closely resembles that of trigonometric functions, and which we shall proceed to develop. It must, however, be kept in mind that the independent variable  $u$  is not an angle, as in the case of the trigonometric functions.

Of course, with our notation,  $u = F(k, x) = \operatorname{sn}^{-1} x \pmod{k}$ , or  $u = F(k, \phi) = \operatorname{am}^{-1} \phi \pmod{k}$ .

The fundamental formulas connecting the Elliptic Functions of a single quantity follow immediately from the definitions [1], and are

$$\operatorname{sn}^2 u + \operatorname{cn}^2 u = 1, \quad [2]$$

$$\operatorname{dn}^2 u + k^2 \operatorname{sn}^2 u = 1, \quad [3]$$

$$\frac{d \operatorname{am} u}{du} = \operatorname{dn} u, \quad [4]$$

$$\frac{d \operatorname{sn} u}{du} = \operatorname{cn} u \cdot \operatorname{dn} u, \quad [5]$$

$$\frac{d \operatorname{cn} u}{du} = -\operatorname{sn} u \cdot \operatorname{dn} u, \quad [6]$$

$$\frac{d \operatorname{dn} u}{du} = -k^2 \operatorname{sn} u \cdot \operatorname{cn} u, \quad [7]$$

The only one of this set which needs any explanation is [4].

We have 
$$u = \int_0^\phi \frac{d\phi}{\Delta\phi};$$

hence 
$$du = \frac{d\phi}{\Delta\phi}, \quad d\phi = \frac{d \operatorname{am} u}{\operatorname{dn} u};$$

and, finally, 
$$\frac{d \operatorname{am} u}{du} = \operatorname{dn} u.$$

Since 
$$\int_0^{-\phi} \frac{d\phi}{\Delta\phi} = \int_0^{\phi} \frac{d(-\phi)}{\Delta(-\phi)} = -\int_0^{\phi} \frac{d\phi}{\Delta\phi},$$

we see that

$$\left. \begin{aligned} \operatorname{am}(-u) &= -\operatorname{am} u, \\ \operatorname{sn}(-u) &= -\operatorname{sn} u, \\ \operatorname{cn}(-u) &= \operatorname{cn} u, \\ \operatorname{dn}(-u) &= \operatorname{dn} u, \end{aligned} \right\} \quad [8]$$

That is,  $\operatorname{sn} u$  is an odd function of  $u$ , and  $\operatorname{cn} u$  and  $\operatorname{dn} u$  are even functions of  $u$ .

Since 
$$\int_0^0 \frac{d\phi}{\Delta\phi} = 0,$$

we have

$$\left. \begin{aligned} \operatorname{am}(0) &= 0, \\ \operatorname{sn}(0) &= 0, \\ \operatorname{cn}(0) &= 1, \\ \operatorname{dn}(0) &= 1, \end{aligned} \right\} \quad [9]$$

193. Our addition formula for the sine amplitude flows immediately from [5], Art. 190. Let  $u = F(k, x)$  and  $v = F(k, y)$ , and take the sine amplitude of each member of [5], Art. 190; we get

$$\operatorname{sn}(u+v) = \frac{\operatorname{sn} u \cdot \operatorname{cn} v \cdot \operatorname{dn} v + \operatorname{cn} u \cdot \operatorname{sn} v \cdot \operatorname{dn} u}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}.$$

If now we replace  $v$  by  $-v$ , and simplify by [8], Art. 192, we have

$$\operatorname{sn}(u-v) = \frac{\operatorname{sn} u \cdot \operatorname{cn} v \cdot \operatorname{dn} v - \operatorname{cn} u \cdot \operatorname{sn} v \cdot \operatorname{dn} u}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v},$$

and the two formulas can be combined if we use the sign  $\pm$ ;

$$\operatorname{sn}(u \pm v) = \frac{\operatorname{sn} u \cdot \operatorname{cn} v \cdot \operatorname{dn} v \pm \operatorname{cn} u \cdot \operatorname{sn} v \cdot \operatorname{dn} u}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [1]$$

From [1], with the aid of [2] and [3], Art. 192, we can get, after a rather elaborate reduction, the addition formulas for  $\operatorname{cn}$  and  $\operatorname{dn}$ .

$$\operatorname{cn}(u \pm v) = \frac{\operatorname{cn} u \cdot \operatorname{cn} v \mp \operatorname{sn} u \cdot \operatorname{sn} v \cdot \operatorname{dn} u \cdot \operatorname{dn} v}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [2]$$

$$\operatorname{dn}(u \pm v) = \frac{\operatorname{dn} u \cdot \operatorname{dn} v \mp k^2 \cdot \operatorname{sn} u \cdot \operatorname{sn} v \cdot \operatorname{cn} u \cdot \operatorname{cn} v}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [3]$$

From [1], [2], and [3] a large number of formulas can be readily obtained. We give only those for  $\operatorname{sn}$ ; there are similar ones for  $\operatorname{cn}$  and  $\operatorname{dn}$ .

$$\operatorname{sn}(u + v) + \operatorname{sn}(u - v) = \frac{2 \operatorname{sn} u \cdot \operatorname{cn} v \cdot \operatorname{dn} v}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [4]$$

$$\operatorname{sn}(u + v) - \operatorname{sn}(u - v) = \frac{2 \operatorname{cn} u \cdot \operatorname{sn} v \cdot \operatorname{dn} u}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [5]$$

$$\operatorname{sn}(u + v) \cdot \operatorname{sn}(u - v) = \frac{\operatorname{sn}^2 u - \operatorname{sn}^2 v}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [6]$$

$$1 + \operatorname{sn}(u + v) \cdot \operatorname{sn}(u - v) = \frac{\operatorname{cn}^2 v + \operatorname{sn}^2 u \cdot \operatorname{dn}^2 v}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [7]$$

$$1 + k^2 \operatorname{sn}(u + v) \cdot \operatorname{sn}(u - v) = \frac{\operatorname{dn}^2 v + k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{cn}^2 v}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [8]$$

$$[1 + \operatorname{sn}(u + v)][1 + \operatorname{sn}(u - v)] = \frac{(\operatorname{cn} v + \operatorname{sn} u \cdot \operatorname{dn} v)^2}{1 - k^2 \cdot \operatorname{sn}^2 u \cdot \operatorname{sn}^2 v}. \quad [9]$$

From [2] and [3] comes the useful formula

$$\operatorname{cn}(u + v) = \operatorname{cn} u \cdot \operatorname{cn} v - \operatorname{sn} u \cdot \operatorname{sn} v \cdot \operatorname{dn}(u + v). \quad [10]$$

194. If in formulas [1], [2], and [3] of Art. 193 we let  $v = u$ , we get the following formulas for  $\text{sn } 2u$ ,  $\text{cn } 2u$ , and  $\text{dn } 2u$ :

$$\text{sn } 2u = \frac{2 \text{sn } u \cdot \text{cn } u \cdot \text{dn } u}{1 - k^2 \text{sn}^4 u}, \quad [1]$$

$$\text{cn } 2u = \frac{\text{cn}^2 u - \text{sn}^2 u \cdot \text{dn}^2 u}{1 - k^2 \text{sn}^4 u} = \frac{1 - 2 \text{sn}^2 u + k^2 \text{sn}^4 u}{1 - k^2 \text{sn}^4 u}, \quad [2]$$

$$\text{dn } 2u = \frac{\text{dn}^2 u - k^2 \cdot \text{sn}^2 u \cdot \text{cn}^2 u}{1 - k^2 \text{sn}^4 u} = \frac{1 - 2k^2 \text{sn}^2 u + k^2 \text{sn}^4 u}{1 - k^2 \text{sn}^4 u}. \quad [3]$$

From these come readily

$$1 - \text{cn } 2u = \frac{2 \text{sn}^2 u \cdot \text{dn}^2 u}{1 - k^2 \text{sn}^4 u}, \quad [4]$$

$$1 + \text{cn } 2u = \frac{2 \text{cn}^2 u}{1 - k^2 \text{sn}^4 u}, \quad [5]$$

$$1 - \text{dn } 2u = \frac{2k^2 \text{sn}^2 u \cdot \text{cn}^2 u}{1 - k^2 \text{sn}^4 u}, \quad [6]$$

$$1 + \text{dn } 2u = \frac{2 \text{dn}^2 u}{1 - k^2 \text{sn}^4 u}. \quad [7]$$

195. Replacing  $u$  by  $\frac{u}{2}$ , and dividing [4] by [7] and [6] by [5], Art. 194, we have

$$\text{sn}^2 \frac{u}{2} = \frac{1 - \text{cn } u}{1 + \text{dn } u} = \frac{1 - \text{dn } u}{k^2(1 + \text{cn } u)}, \quad [1]$$

$$\text{cn}^2 \frac{u}{2} = \frac{\text{dn } u + \text{cn } u}{1 + \text{dn } u} = \frac{-k'^2 + k^2 \text{cn } u + \text{dn } u}{k^2(1 + \text{cn } u)}, \quad [2]$$

$$\text{dn}^2 \frac{u}{2} = \frac{k'^2 + \text{dn } u + k^2 \text{cn } u}{1 + \text{dn } u} = \frac{(\text{cn } u + \text{dn } u)}{(1 + \text{cn } u)}, \quad [3]$$

where  $k'^2 = 1 - k^2$ , and is the square of the complementary modulus.

From [1], [2], and [3], we can get without difficulty the set

$$\operatorname{sn}^2 \frac{u}{2} = \frac{\operatorname{dn} u - \operatorname{cn} u}{k'^2 + \operatorname{dn} u - k^2 \operatorname{cn} u}, \quad [4]$$

$$\operatorname{cn}^2 \frac{u}{2} = \frac{k'^2(1 + \operatorname{cn} u)}{k'^2 + \operatorname{dn} u - k^2 \operatorname{cn} u}, \quad [5]$$

$$\operatorname{dn}^2 \frac{u}{2} = \frac{k'^2(1 + \operatorname{dn} u)}{k'^2 + \operatorname{dn} u - k^2 \operatorname{cn} u}. \quad [6]$$

Numerous additional formulas can be obtained by the exercise of a little ingenuity, but we have given the most useful and important ones, and they form a set as complete as the usual collections of trigonometric formulas.

*Periodicity of the Elliptic Functions.*

196. We have seen (Art. 186, [2]) that

$$F(k, n\pi + \rho) = 2nK + F(k, \rho), \quad [1]$$

where  $K$  is the complete Elliptic Integral of the first class.

Let  $u = F(k, \rho)$ , and take the amplitude of each member of [1]; we get

$$\operatorname{am}(u + 2nK) = n\pi + \operatorname{am} u; \quad [2]$$

or, replacing  $n$  by  $2n$ ,

$$\operatorname{am}(u + 4nK) = 2n\pi + \operatorname{am} u; \quad [3]$$

whence

$$\left. \begin{aligned} \operatorname{sn}(u + 4nK) &= \operatorname{sn} u, \\ \operatorname{cn}(u + 4nK) &= \operatorname{cn} u, \\ \operatorname{dn}(u + 4nK) &= \operatorname{dn} u, \end{aligned} \right\}; \quad [4]$$

and  $\operatorname{sn} u$ ,  $\operatorname{cn} u$ ,  $\operatorname{dn} u$  are periodic functions, and have the real period  $4K$ .  $\operatorname{dn} u$  actually has the smaller period  $2K$ , as may be seen by taking the delta of both members of [2].

Since the amplitude of  $K$  is  $\frac{\pi}{2}$ , we have

$$\left. \begin{aligned} \operatorname{sn} K &= 1, \\ \operatorname{cn} K &= 0, \\ \operatorname{dn} K &= k', \end{aligned} \right\}, \quad [5]$$

and our addition formulas [1], [2], [3], Art. 193, give us readily

$$\left. \begin{aligned} \operatorname{sn}(u + K) &= \frac{\operatorname{cn} u}{\operatorname{dn} u}, \\ \operatorname{cn}(u + K) &= -\frac{k' \operatorname{sn} u}{\operatorname{dn} u}, \\ \operatorname{dn}(u + K) &= \frac{k'}{\operatorname{dn} u}, \end{aligned} \right\}, \quad [6]$$

$$\left. \begin{aligned} \operatorname{sn}(u + 2K) &= -\operatorname{sn} u, \\ \operatorname{cn}(u + 2K) &= -\operatorname{cn} u, \\ \operatorname{dn}(u + 2K) &= \operatorname{dn} u, \end{aligned} \right\}, \quad [7]$$

$$\left. \begin{aligned} \operatorname{sn}(u + 3K) &= -\frac{\operatorname{cn} u}{\operatorname{dn} u}, \\ \operatorname{cn}(u + 3K) &= \frac{k' \operatorname{sn} u}{\operatorname{dn} u}, \\ \operatorname{dn}(u + 3K) &= \frac{k'}{\operatorname{dn} u}, \end{aligned} \right\}, \quad [8]$$

$$\left. \begin{aligned} \operatorname{sn}(u + 4K) &= \operatorname{sn} u, \\ \operatorname{cn}(u + 4K) &= \operatorname{cn} u, \\ \operatorname{dn}(u + 4K) &= \operatorname{dn} u, \end{aligned} \right\}, \quad [9]$$

a confirmation of [4].

197. It is easy to get formulas for the  $\text{sn}$ ,  $\text{cn}$ , and  $\text{dn}$  of an imaginary variable,  $u\sqrt{-1}$ , by the aid of a transformation due to Jacobi.

Let 
$$v = F(k, \phi) = \int_0^\phi \frac{d\phi}{\Delta\phi}, \tag{1}$$

so that  $\phi = \text{am } v$ ,  $\sin \phi = \text{sn } v$ , and  $\cos \phi = \text{cn } v$ . In (1), replace  $\phi$  by  $\psi$ ,  $\phi$  and  $\psi$  being connected by the relation

$$\sin \phi = \sqrt{-1} \cdot \tan \psi, \tag{2}$$

whence 
$$\cos \phi = \sec \psi, \tag{3}$$

$$\Delta\phi = \sqrt{1 - k^2 \sin^2 \phi} = \sqrt{1 + k^2 \tan^2 \psi}, \tag{4}$$

and 
$$d\phi = \sqrt{-1} \cdot \sec \psi \cdot d\psi.$$

Since  $\psi$  and  $\phi$  equal zero together,

$$\begin{aligned} v &= \int_0^\phi \frac{d\phi}{\Delta\phi} = \sqrt{-1} \int_0^\psi \frac{\sec \psi d\psi}{\sqrt{1 + k^2 \tan^2 \psi}} \\ &= \sqrt{-1} \int_0^\psi \frac{d\psi}{\sqrt{1 - k'^2 \sin^2 \psi}} = \sqrt{-1} \cdot F(k', \psi). \end{aligned}$$

If now we let  $u = F(k', \psi)$ ,

we have 
$$v = u\sqrt{-1}. \tag{5}$$

Hence, since  $\psi = \text{am } u \pmod{k'}$ , we have from (2), (3), and (4),

$$\text{sn } v \pmod{k} = \sqrt{-1} \frac{\text{sn } u \pmod{k'}}{\text{cn } u \pmod{k'}},$$

$$\text{cn } v \pmod{k} = \frac{1}{\text{cn } u \pmod{k'}},$$

$$\text{dn } v \pmod{k} = \frac{\text{dn } u \pmod{k'}}{\text{cn } u \pmod{k'}};$$

or, as  $v = u\sqrt{-1}$ ,





Since the amplitude of  $K$  is  $\frac{\pi}{2}$ , we have

$$\left. \begin{aligned} \operatorname{sn} K &= 1, \\ \operatorname{cn} K &= 0, \\ \operatorname{dn} K &= k', \end{aligned} \right\}, \quad [5]$$

and our addition formulas [1], [2], [3], Art. 193, give us readily

$$\left. \begin{aligned} \operatorname{sn}(u + K) &= \frac{\operatorname{cn} u}{\operatorname{dn} u}, \\ \operatorname{cn}(u + K) &= -\frac{k' \operatorname{sn} u}{\operatorname{dn} u}, \\ \operatorname{dn}(u + K) &= \frac{k'}{\operatorname{dn} u}, \end{aligned} \right\}, \quad [6]$$

$$\left. \begin{aligned} \operatorname{sn}(u + 2K) &= -\operatorname{sn} u, \\ \operatorname{cn}(u + 2K) &= -\operatorname{cn} u, \\ \operatorname{dn}(u + 2K) &= \operatorname{dn} u, \end{aligned} \right\}, \quad [7]$$

$$\left. \begin{aligned} \operatorname{sn}(u + 3K) &= -\frac{\operatorname{cn} u}{\operatorname{dn} u}, \\ \operatorname{cn}(u + 3K) &= \frac{k' \operatorname{sn} u}{\operatorname{dn} u}, \\ \operatorname{dn}(u + 3K) &= \frac{k'}{\operatorname{dn} u}, \end{aligned} \right\}, \quad [8]$$

$$\left. \begin{aligned} \operatorname{sn}(u + 4K) &= \operatorname{sn} u, \\ \operatorname{cn}(u + 4K) &= \operatorname{cn} u, \\ \operatorname{dn}(u + 4K) &= \operatorname{dn} u, \end{aligned} \right\}, \quad [9]$$

• confirmation of [4].

197. It is easy to get formulas for the  $\text{sn}$ ,  $\text{cn}$ , and  $\text{dn}$  of an imaginary variable,  $u\sqrt{-1}$ , by the aid of a transformation due to Jacobi.

$$\text{Let } v = F(k, \phi) = \int_0^\phi \frac{d\phi}{\Delta\phi}, \quad (1)$$

so that  $\phi = \text{am } v$ ,  $\sin \phi = \text{sn } v$ , and  $\cos \phi = \text{cn } v$ . In (1), replace  $\phi$  by  $\psi$ ,  $\phi$  and  $\psi$  being connected by the relation

$$\sin \phi = \sqrt{-1} \cdot \tan \psi, \quad (2)$$

$$\text{whence } \cos \phi = \sec \psi, \quad (3)$$

$$\Delta\phi = \sqrt{1 - k^2 \sin^2 \phi} = \sqrt{1 + k^2 \tan^2 \psi}, \quad (4)$$

$$\text{and } d\phi = \sqrt{-1} \cdot \sec \psi \cdot d\psi.$$

Since  $\psi$  and  $\phi$  equal zero together,

$$\begin{aligned} v &= \int_0^\phi \frac{d\phi}{\Delta\phi} = \sqrt{-1} \int_0^\psi \frac{\sec \psi d\psi}{\sqrt{1 + k^2 \tan^2 \psi}} \\ &= \sqrt{-1} \int_0^\psi \frac{d\psi}{\sqrt{1 - k'^2 \sin^2 \psi}} = \sqrt{-1} \cdot F(k', \psi). \end{aligned}$$

If now we let  $u = F(k', \psi)$ ,

$$\text{we have } v = u\sqrt{-1}. \quad (5)$$

Hence, since  $\psi = \text{am } u \pmod{k'}$ , we have from (2), (3), and (4),

$$\text{sn } v \pmod{k} = \sqrt{-1} \frac{\text{sn } u \pmod{k'}}{\text{cn } u \pmod{k'}},$$

$$\text{cn } v \pmod{k} = \frac{1}{\text{cn } u \pmod{k'}},$$

$$\text{dn } v \pmod{k} = \frac{\text{dn } u \pmod{k'}}{\text{cn } u \pmod{k'}};$$

or, as  $v = u\sqrt{-1}$ ,

$$\left. \begin{aligned} \operatorname{sn} u \sqrt{-1} \pmod{k} &= \sqrt{-1} \frac{\operatorname{sn} u \pmod{k'}}{\operatorname{cn} u \pmod{k'}}, \\ \operatorname{cn} u \sqrt{-1} \pmod{k} &= \frac{1}{\operatorname{cn} u \pmod{k'}}, \\ \operatorname{dn} u \sqrt{-1} \pmod{k} &= \frac{\operatorname{dn} u \pmod{k'}}{\operatorname{cn} u \pmod{k'}} \end{aligned} \right\} \quad [6]$$

It is interesting to note that if  $u$  is replaced in (6) by  $u\sqrt{-1}$ , the formulas reduce to

$$\begin{aligned} \operatorname{sn}(-u) &= -\operatorname{sn} u, \\ \operatorname{cn}(-u) &= \operatorname{cn} u, \\ \operatorname{dn}(-u) &= \operatorname{dn} u, \end{aligned}$$

and are still true. Consequently, in (6),  $u$  may be either a real or a pure imaginary.

$$\text{Let} \quad \int_0^{\frac{\pi}{2}} \frac{d\psi}{\Delta\psi \pmod{k'}} = \int_0^{\frac{\pi}{2}} \frac{d\psi}{\sqrt{1-k'^2 \sin^2 \psi}} = K'.$$

Then, by Art. 196,  $4K'$  is a period for  $\operatorname{sn} u \pmod{k'}$ ,  $\operatorname{cn} u \pmod{k'}$ , and  $\operatorname{dn} u \pmod{k'}$ .

Hence

$$\begin{aligned} \operatorname{sn}(u\sqrt{-1} + 4nK'\sqrt{-1}) &= \operatorname{sn} u\sqrt{-1}, \\ \operatorname{cn}(u\sqrt{-1} + 4nK'\sqrt{-1}) &= \operatorname{cn} u\sqrt{-1}, \\ \operatorname{dn}(u\sqrt{-1} + 4nK'\sqrt{-1}) &= \operatorname{dn} u\sqrt{-1}; \end{aligned}$$

or, replacing  $u\sqrt{-1}$  by  $v$ ,

$$\left. \begin{aligned} \operatorname{sn}(v + 4nK'\sqrt{-1}) &= \operatorname{sn} v, \\ \operatorname{cn}(v + 4nK'\sqrt{-1}) &= \operatorname{cn} v, \\ \operatorname{dn}(v + 4nK'\sqrt{-1}) &= \operatorname{dn} v, \end{aligned} \right\} \quad [7]$$

and  $4K'\sqrt{-1}$  is a period for  $\operatorname{sn}$ ,  $\operatorname{cn}$ , and  $\operatorname{dn}$ .

We see, then, that our Elliptic Functions, like Trigonometric Functions, have a real period, and, like Exponential Functions, have a pure imaginary period. They are, then, what may be called

To make a table of Elliptic Functions:

Take from Hill's See. Var. or Legendre's Traité the tabular values of  $u_1 = F(K, \phi)$ ,  $u_2 = E(K, \phi)$ ,  $u_3 = \Pi(u, k, \phi)$ . Tabulate the values of  $u_1, u_2, u_3$ , and the values corresponding of  $\phi_1, \phi_2, \phi_3$ .

Then form new tables of  $u_1, u_2, u_3$ , using the  $\phi_1, \phi_2, \phi_3$  for taking out <sup>corresponding</sup>  $\sin u, \cos u$  from <sup>table of</sup> trigonometric natural sines and cosines.

Thus we get for any  $u$  the corresponding  $\sin u, \cos u$

$u$	$\sin u$	$\cos u$	$dn u$
.....	.....	.....	.....

The last column  $dn u$  must be calculated by the formula:

$$\sqrt{1-k^2x^2} = \Delta \phi = \sqrt{1-k^2 \sin^2 \phi} = dn u,$$

The values of  $k$ , and  $\phi$  being known.

As a matter of fact no complete set of tables for the Elliptic Functions has been published, and their values are usually obtained indirectly from Legendre's Tables of Elliptic Integrals (v. Arts. 186, 187), unless especial accuracy is required, in which case they must be computed by methods which we have not space to give.

198. The Elliptic Integral of the second class  $E(k, \phi)$  can be expressed in terms of Elliptic Functions, and for some purposes there is a decided advantage in the new form.

We have 
$$E(k, \phi) = \int_0^\phi \Delta \phi \cdot d\phi.$$

Let  $u = F(k, \phi)$ , then  $\phi = \text{am } u$ , and  $E(k, \phi)$  may be written  $E(k, \text{am } u)$ , or, more simply,  $E(\text{am } u)$ , if the modulus can be omitted without danger of confusion.

Then 
$$E(\text{am } u) = \int_0^{\text{am } u} \text{dn } u \cdot d\text{am } u;$$

or, since by (4), Art. 192,

$$d\text{am } u = \text{dn } u \cdot du,$$

$$E(\text{am } u) = \int_0^u \text{dn}^2 u \cdot du. \quad [1]$$

As an example of the usefulness of the form just given in [1], we will employ it in getting an *addition formula* for Elliptic Integrals of the second class.

$$\begin{aligned} E(\text{am } u) + E(\text{am } v) &= \int_0^u \text{dn}^2 u \cdot du + \int_0^v \text{dn}^2 v \cdot dv \\ &= \int_0^u \text{dn}^2 z \cdot dz + \int_0^v \text{dn}^2 z \cdot dz \\ &= \int_0^{u+v} \text{dn}^2 z \cdot dz + \int_0^v \text{dn}^2 z \cdot dz - \int_u^{u+v} \text{dn}^2 z \cdot dz \\ &= E[\text{am}(u+v)] + \int_0^v \text{dn}^2 z \cdot dz - \int_u^{u+v} \text{dn}^2 z \cdot dz. \end{aligned}$$

Replacing  $z$  by  $u + z$ , and remembering that  $u$  and  $v$  are given constants,

$$\int_u^{u+v} \text{dn}^2 z \cdot dz = \int_0^v \text{dn}^2 (u + z) dz,$$

and

$$E(\text{am } u) + E(\text{am } v) = E[\text{am } (u + v)] - \int_0^v [\text{dn}^2 (u + z) - \text{dn}^2 z] dz. \quad (2)$$

$$\text{dn}^2 (u + z) - \text{dn}^2 z = [\text{dn } (u + z) + \text{dn } z][\text{dn } (u + z) - \text{dn } z]. \quad (3)$$

We can obtain from [3], Art. 193, the following formulas analogous to [4] and [5], Art. 193,

$$\text{dn } (u + v) + \text{dn } (u - v) = \frac{2 \text{dn } u \cdot \text{dn } v}{1 - k^2 \text{sn}^2 u \cdot \text{sn}^2 v}, \quad (4)$$

$$\text{dn } (u + v) - \text{dn } (u - v) = -\frac{2 k^2 \text{sn } u \cdot \text{sn } v \cdot \text{cn } u \cdot \text{cn } v}{1 - k^2 \text{sn}^2 u \cdot \text{sn}^2 v}. \quad (5)$$

If in (4) and (5) we let  $u + v = u + z$ , and  $u - v = z$ , and substitute the results in (3), we get

$$\begin{aligned} & \text{dn}^2 (u + z) - \text{dn}^2 z \\ &= \frac{4 k^2 \text{sn} \left(\frac{u}{2} + z\right) \text{cn} \left(\frac{u}{2} + z\right) \text{dn} \left(\frac{u}{2} + z\right) \text{sn} \frac{u}{2} \text{cn} \frac{u}{2} \text{dn} \frac{u}{2}}{\left[1 - k^2 \text{sn}^2 \frac{u}{2} \text{sn}^2 \left(\frac{u}{2} + z\right)\right]^2}; \end{aligned}$$

and

$$\begin{aligned} & \int [\text{dn}^2 (u + z) - \text{dn}^2 z] dz \\ &= -2 \text{sn} \frac{u}{2} \text{cn} \frac{u}{2} \text{dn} \frac{u}{2} \int \frac{2 k^2 \text{sn} \left(\frac{u}{2} + z\right) \text{cn} \left(\frac{u}{2} + z\right) \text{dn} \left(\frac{u}{2} + z\right) dz}{\left[1 - k^2 \text{sn}^2 \frac{u}{2} \text{sn}^2 \left(\frac{u}{2} + z\right)\right]^2} \\ &= \frac{2 \text{sn} \frac{u}{2} \text{cn} \frac{u}{2} \text{dn} \frac{u}{2}}{\text{sn}^2 \frac{u}{2}} \cdot \frac{1}{1 - k^2 \text{sn}^2 \frac{u}{2} \text{sn}^2 \left(\frac{u}{2} + z\right)} \end{aligned}$$

since  $-2k^2 \operatorname{sn}^2 \frac{u}{2} \operatorname{sn} \left( \frac{u}{2} + z \right) \operatorname{cn} \left( \frac{u}{2} + z \right) \operatorname{dn} \left( \frac{u}{2} + z \right) dz$  is the differential of  $1 - k^2 \operatorname{sn}^2 \frac{u}{2} \operatorname{sn}^2 \left( \frac{u}{2} + z \right)$ .

$$\begin{aligned} & \int_0^v [\operatorname{dn}^2(u+z) - \operatorname{dn}^2 z] dz \\ &= -\frac{2 \operatorname{sn} \frac{u}{2} \operatorname{cn} \frac{u}{2} \operatorname{dn} \frac{u}{2}}{\operatorname{sn}^2 \frac{u}{2}} \left[ \frac{1}{1 - k^2 \operatorname{sn}^2 \frac{u}{2} \operatorname{sn}^2 \left( \frac{u}{2} + v \right)} - \frac{1}{1 - k^2 \operatorname{sn}^2 \frac{u}{2}} \right] \\ &= -\frac{k^2 2 \operatorname{sn} \frac{u}{2} \operatorname{cn} \frac{u}{2} \operatorname{dn} \frac{u}{2}}{1 - k^2 \operatorname{sn}^2 \frac{u}{2}} \cdot \frac{\operatorname{sn}^2 \left( \frac{u}{2} + v \right) - \operatorname{sn}^2 \frac{u}{2}}{1 - k^2 \operatorname{sn}^2 \frac{u}{2} \operatorname{sn}^2 \left( \frac{u}{2} + v \right)} \\ &= -k^2 \cdot \operatorname{sn} u \cdot \operatorname{sn} v \cdot \operatorname{sn}(u+v), \end{aligned}$$

by (1), Art. 194, and [6], Art. 193.

Hence by (2),

$$E(\operatorname{am} u) + E(\operatorname{am} v) = E[\operatorname{am}(u+v)] + k^2 \operatorname{sn} u \cdot \operatorname{sn} v \cdot \operatorname{sn}(u+v), \quad [6]$$

our required addition formula.

#### APPLICATIONS.

##### *Rectification of the Lemniscate.*

199. From the polar equation of the Lemniscate,  $r^2 = a^2 \cos 2\theta$ , referred to its centre as origin and its axis as axis, we get as the length of the arc, measured from the vertex to any point,  $P$ , whose coördinates are  $r$  and  $\theta$ .

$$s = a \int_0^\theta \frac{d\theta}{\sqrt{\cos 2\theta}} = a \int_0^\theta \frac{d\theta}{\sqrt{1 - 2 \sin^2 \theta}}, \quad [1]$$

and for the arc of the quadrant of the Lemniscate, that is, the arc from vertex to centre,

$$s_v = \int_0^{\frac{\pi}{4}} \frac{d\theta}{\sqrt{1-2\sin^2\theta}}. \tag{2}$$

These differ from Elliptic Integrals of the first class only in that the coefficient of  $\sin^2\theta$  is greater than unity, and they may be reduced to the standard form by a simple device.

Introduce in [1]  $\phi$  in place of  $\theta$ ,  $\phi$  and  $\theta$  being connected by the relation  $\sin^2\phi = 2\sin^2\theta$ .


Then we have  $\sqrt{1-2\sin^2\theta} = \cos\phi$ ,

and  $d\theta = \frac{\sqrt{2}}{2} \frac{\cos\phi d\phi}{\sqrt{1-\frac{1}{2}\sin^2\phi}}$ .

Hence  $s = \frac{a\sqrt{2}}{2} \int_0^\phi \frac{d\phi}{\sqrt{1-\frac{1}{2}\sin^2\phi}} = \frac{a\sqrt{2}}{2} F\left(\frac{\sqrt{2}}{2}, \phi\right)$ , [3]

and  $s_v = \frac{a\sqrt{2}}{2} \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-\frac{1}{2}\sin^2\phi}} = \frac{a\sqrt{2}}{2} F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right)$ . [4]

The auxiliary angle  $\phi$  is very easily constructed when the point  $P$  of the Lemniscate is given. We have  $r = a\sqrt{\cos 2\theta}$ , and we have seen that  $\sqrt{\cos 2\theta} = \cos\phi$ ; hence  $r = a\cos\phi$ . If, then, on  $a$  as a diameter we describe a semi-circumference, and with the centre  $O$  of the Lemniscate as a centre, and with a radius equal to  $r$ , we describe an arc, and join with  $O$  the point  $Q$  where this arc intersects the semi-circumference, the angle made by  $OQ$  with  $a$  is equal to  $\phi$ . For  $OQ = a\cos\angle AOQ$  and  $OP = a\sqrt{\cos 2\theta}$ .





## EXAMPLES.

(1) Find the numerical value of  $\int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-4\sin^2\phi}}$ .

*Ans.* 0.843.

(2) Reduce  $\int_0^{\phi} \frac{d\phi}{\sqrt{1-n\sin^2\phi}}$  to an Elliptic Integral of the first class, when  $n > 1$ .

*Ans.*  $\frac{1}{\sqrt{n}} \int_0^{\psi} \frac{d\psi}{\sqrt{1-\frac{1}{n}\sin^2\psi}}$  where  $\sin^2\psi = n\sin^2\phi$ .

(3) The half-axis of a Lemniscate is 2. What is the length of the arc of a quadrant? of the arc from the vertex to the point whose polar angle is  $30^\circ$ ? *Ans.* 2.622; 1.168.

In the inverse problem of cutting off an arc of given length the Elliptic Functions are of service. As an interesting example, let us find the point which bisects the quadrantal arc of the Lemniscate.

Here  $s = \frac{a\sqrt{2}}{2} \frac{1}{2} F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right),$

and we wish to find  $\phi$  and then  $\theta$ .

Let  $u = F\left(\frac{\sqrt{2}}{2}, \frac{\pi}{2}\right);$  we need  $\operatorname{am} \frac{u}{2}.$

$$\operatorname{am} u = \frac{\pi}{2}, \operatorname{sn} u = 1, \operatorname{cn} u = 0, \text{ and } \operatorname{dn} u = \frac{\sqrt{2}}{2}.$$

By [1] and [2], Art. 195,

$$\operatorname{sn}^2 \frac{u}{2} = \frac{1 - \operatorname{cn} u}{1 + \operatorname{dn} u}, \quad \operatorname{cn}^2 \frac{u}{2} = \frac{\operatorname{dn} u + \operatorname{cn} u}{1 + \operatorname{dn} u}.$$

Therefore,

$$\frac{\operatorname{sn}^2 \frac{u}{2}}{\operatorname{cn}^2 \frac{u}{2}} = \operatorname{tn}^2 \frac{u}{2} = \frac{1 - \operatorname{cn} u}{\operatorname{dn} u + \operatorname{cn} u} = \frac{1}{\frac{\sqrt{2}}{2}} = \sqrt{2}.$$

If, then, the required amplitude is  $\phi$ ,


$$\tan^2 \phi = \sqrt{2},$$

and

$$\tan \phi = \sqrt[4]{2}.$$

Since  $\sin^2 \phi = 2 \sin^2 \theta$ , we can compute  $\theta$  without difficulty, and so get our required point. If, however, a construction will suffice, a very simple one gives the point.

Erect at  $A$  a perpendicular whose length is a mean proportional between  $a$  and  $a\sqrt{2}$ . The angle subtended at  $O$  by this perpendicular is  $\phi$ , and the corresponding point,  $P$ , is found by the method described on page 255.



*Rectification of the Ellipse.*

200. We have seen in Art. 177 that the length of an arc of an Ellipse measured from the end of the minor axis is

$$s = \int_0^x \sqrt{\frac{a^2 - e^2 x^2}{a^2 - x^2}} \cdot dx. \tag{1}$$

If we let  $x = a \sin \phi$ , [1] becomes


$$s = a \int_0^\phi \sqrt{1 - e^2 \sin^2 \phi} \cdot d\phi = a E(e, \phi), \tag{2}$$

$e$ , the modulus of the Elliptic Integral, being the eccentricity of the Ellipse. If  $x = a$ ,  $\phi = \frac{\pi}{2}$ , and the length of the Elliptic quadrant is

$$s_q = a \int_0^{\frac{\pi}{2}} \sqrt{1 - e^2 \sin^2 \phi} \cdot d\phi = a E\left(e, \frac{\pi}{2}\right). \tag{3}$$

The length of an arc of the Elliptic quadrant, not measured from the extremity of the minor axis, can of course be expressed as the difference between two Elliptic Integrals of the second class.

The amplitude  $\phi$ , corresponding to a given point  $P$ , of the Ellipse, is easily constructed as follows: On the major axis as diameter describe a circumference; extend the ordinate of  $P$  until it meets the circumference, and join the point of intersection with the centre of the ellipse. The angle the joining line makes with the minor axis is seen to be the required amplitude  $\phi$ . If  $\phi$  is given,  $P$  may be found by reversing the order of the steps of the construction.



#### EXAMPLES.

The equation of an ellipse is  $\frac{x^2}{16} + \frac{y^2}{8} = 1$ , required the length of the quadrantal arc: of the arc whose extremities have the abscissas 2 and  $2\sqrt{2}$ . *Ans.* 5.4; 0.944.

(2) Find the abscissa of the end of the unit arc measured from the extremity of the minor axis in the ellipse  $\frac{x^2}{16} + \frac{y^2}{8} = 1$ ; of the point which bisects the arc of the quadrant. *Ans.* 0.996; 2.57.

201. By the aid of the addition formula

$$E(\text{am } u) + E(\text{am } v) = E[\text{am } (u + v)] + k^2 \text{sn } u \text{sn } v \text{sn } (u + v) \quad ([6], \text{ Art. 198})$$

it is always possible to find an arc of an ellipse differing from the sum of two given arcs by an expression which is algebraic in terms of the abscissas of the extremities of the three arcs. This will be clearer if we modify slightly the form of our addition formula.

Let  $\phi = \text{am } u$ ,  $\psi = \text{am } v$ , and  $\sigma = \text{am}(u + v)$ .

Then the formula given above becomes

$$E(k, \phi) + E(k, \psi) = E(k, \sigma) + k^2 \sin \phi \sin \psi \sin \sigma, \quad [1]$$

where  $\phi$ ,  $\psi$ , and  $\sigma$  are three angles connected by the relation

$$\cos \sigma = \cos \phi \cos \psi - \sin \phi \sin \psi \Delta \sigma, \quad [2]$$

by [10], Art. 193.

If we multiply [1] by  $a$  and take  $k$  equal to  $e$ , we get

$$aE(e, \phi) + aE(e, \psi) = aE(e, \sigma) + \frac{e^2}{a^2} x_1 \cdot x_2 \cdot x_3,$$

if  $x_1$ ,  $x_2$ , and  $x_3$  are the abscissas of the points whose amplitudes are  $\phi$ ,  $\psi$ , and  $\sigma$ .

The most interesting case is when  $\sigma = \frac{\pi}{2}$ , in which case  $aE(e, \sigma)$  is the arc of a quadrant. [2] then reduces to

$$0 = \cos \phi \cos \psi - \sin \phi \sin \psi \sqrt{1 - e^2},$$


or 
$$\frac{b}{a} \sin \phi \sin \psi = \cos \phi \cos \psi,$$

or 
$$\tan \phi \tan \psi = \frac{a}{b}, \quad [3]$$

and we get from [1]

$$aE(e, \phi) - \left[ aE\left(e, \frac{\pi}{2}\right) - aE(e, \psi) \right] = ae^2 \sin \phi \sin \psi. \quad [4]$$

If, then, any point,  $P$ , is given, [3] will enable us to get the amplitude of a second point,  $Q$ , and thus to find  $Q$ ,  $Q$  and  $P$  being so related that the arc  $BP$ , minus the arc  $AQ$ , shall be equal to a quantity which is proportional to the product of the abscissas of  $P$  and  $Q$ .



For the special case where  $\phi$  and  $\psi$  are equal we have from

$$[3], \quad \tan \phi = \sqrt{\frac{a}{b}},$$

and from [4],

$$BP - AP = ae^2 \sin^2 \phi = \frac{a^2 e^2}{a + b} = a - b.$$

This point, which divides the quadrant into two arcs whose difference is equal to the difference between the semi-axes, has a number of curious properties, and is known as *Fagnani's point*.

#### EXAMPLES.

(1) Show that the distance of the normal at Fagnani's point, from the centre of the ellipse, is equal to  $a - b$ .

(2) Show that the angle between the normals at  $P$  and  $Q$  in the figure is equal to  $\psi - \phi$ ; that the normals are equidistant from  $O$ ; that this distance is  $BP - AQ$ .

#### *Rectification of the Hyperbola.*

202. If the arc of the Hyperbola is measured from the vertex to any given point,  $P$ , whose coördinates are  $x$  and  $y$ , its length is easily found to be

$$s = \int_0^y \left( \frac{1 + \frac{a^2 + b^2}{b^4} y^2}{1 + \frac{y^2}{b^2}} \right)^{\frac{1}{2}} dy, \quad [1]$$

$$\text{or} \quad s = \int_0^y \left( \frac{1 + \frac{a^2 e^2}{b^4} y^2}{1 + \frac{y^2}{b^2}} \right)^{\frac{1}{2}} dy, \quad [2]$$

if  $e$  is the eccentricity of the Hyperbola. Let

$$\frac{ae}{b^2} y = \tan \phi,$$

and [2] becomes

$$s = \frac{b^2}{ae} \int_0^\phi \frac{\sec^2 \phi \, d\phi}{\sqrt{1 - \frac{1}{e^2} \sin^2 \phi}};$$

hence 
$$s = \frac{b^2}{ae} \int_0^\phi \frac{\sec^2 \phi \, d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = \frac{b^2}{ae} \int_0^\phi \frac{\sec^2 \phi \, d\phi}{\Delta \phi} \quad [3]$$

if 
$$k = \frac{1}{e}.$$

Now 
$$\frac{1}{\Delta \phi} = \frac{1 - k^2}{1 - k^2} \cdot \frac{1}{\Delta \phi} = \frac{1}{1 - k^2} \cdot \frac{1 - k^2 \sin^2 \phi - k^2 \cos^2 \phi}{\Delta \phi}$$

$$= \frac{1}{1 - k^2} \left[ \Delta \phi - \frac{k^2 \cos^2 \phi}{\Delta \phi} \right];$$

and 
$$s = \frac{b^2}{ae} \cdot \frac{1}{1 - k^2} \left[ \int_0^\phi \sec^2 \phi \, \Delta \phi \, d\phi - k^2 \int_0^\phi \frac{d\phi}{\Delta \phi} \right]$$

$$= \frac{b^2}{ae} \cdot \frac{1}{1 - k^2} \left[ \int_0^\phi \sec^2 \phi \, \Delta \phi \, d\phi - k^2 F(k, \phi) \right].$$

If we integrate by parts,

$$\int_0^\phi \sec^2 \phi \, \Delta \phi \, d\phi = \tan \phi \, \Delta \phi + k^2 \int_0^\phi \frac{\sin^2 \phi}{\Delta \phi} \, d\phi;$$

but 
$$k^2 \frac{\sin^2 \phi}{\Delta \phi} = \frac{1}{\Delta \phi} - \Delta \phi,$$

and 
$$k^2 \int_0^\phi \frac{\sin^2 \phi}{\Delta \phi} \cdot d\phi = F(k, \phi) - E(k, \phi).$$

Hence


$$s = \frac{b^2}{ae} F(k, \phi) - \frac{b^2}{ae(1 - k^2)} [E(k, \phi) - \tan \phi \, \Delta \phi].$$

But 
$$1 - k^2 = \frac{b^2}{a^2 e^2},$$

therefore  $s = \frac{b^2}{ae} F(k, \phi) - aeE(k, \phi) + ae \tan \phi \Delta \phi,$

or  $s = \frac{b^2}{ae} F\left(\frac{1}{e}, \phi\right) - aeE\left(\frac{1}{e}, \phi\right) + ae \tan \phi \Delta \phi. \quad [4]$

The angle  $\phi$  corresponding to a given point  $P$  is easily con-



structed. We have only to erect a perpendicular to the transverse axis at a distance  $\frac{b^2}{ae} = \frac{b^2}{\sqrt{a^2 + b^2}}$  from the origin; that is, at a distance from the centre equal to the projection of  $b$  on the asymptote, and to join the projection of  $P$  on this line with the centre. The angle made by the joining line with the transverse axis is  $\phi$ , for its tangent is clearly  $\frac{y}{\frac{b^2}{\sqrt{a^2 + b^2}}}$ .

#### EXAMPLES.

(1) Find the length of the arc of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ , measured from the vertex to the point whose ordinate is 2.

*Ans.* 2.194.

(2) Show that  $ae \tan \phi \Delta \phi$  is the distance from the centre to the normal at  $P$ .

(3) Show that the limiting value approached by the difference between the arc and the portion of the asymptote cut off by a perpendicular upon it from  $P$ , as  $P$  recedes indefinitely from

the origin, is  $aeE\left(\frac{1}{e}, \frac{\pi}{2}\right) - \frac{b^2}{ae}F\left(\frac{1}{e}, \frac{\pi}{2}\right)$ . This is generally referred to as the difference between the length of the infinite arc of the hyperbola and the length of the asymptote.

Show that in example (1) this difference is equal to 2.803.


*The Pendulum.*

203. We have seen in Art. 176 that if a pendulum starts from rest at a point of its arc whose distance above the lowest point is  $y_0$ , the time required in rising from the lowest point to a point whose distance above the lowest point is  $y$ , is

$$t = \sqrt{\frac{a}{g}} \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = \sqrt{\frac{a}{g}} F(k, \phi), \quad [1]$$

where  $k = \sqrt{\frac{y_0}{2a}}$ , and  $\sin \phi = \sqrt{\frac{y}{y_0}}$ .

In the figure let  $A$  be the lowest point of the arc,  $B$  the



highest point reached by the pendulum, and  $P$  the point reached at the expiration of the time  $t$ . Call  $AOB$   $\alpha$ , and  $AOP$   $\theta$ .

Then  $\frac{y_0}{a} = 1 - \cos \alpha$ , and  $\sqrt{\frac{y_0}{2a}} = \sqrt{\frac{1}{2}(1 - \cos \alpha)} = \sin \frac{\alpha}{2} = k$ .

Consequently the modulus of the Elliptic Integral in [1] is




the sine of one-fourth the angle through which the pendulum swings.

$$\frac{y}{a} = 1 - \cos \theta.$$

$$\text{and } \sqrt{\frac{y}{2a}} = \sqrt{\frac{1}{2}(1 - \cos \theta)} = \sin \frac{\theta}{2},$$

$$\text{and } \sin \phi = \sqrt{\frac{y}{y_0}} = \frac{\sqrt{\frac{y}{2a}}}{\sqrt{\frac{y_0}{2a}}} = \frac{\sin \frac{\theta}{2}}{\sin \frac{\alpha}{2}},$$

and therefore the sine of the amplitude of the Elliptic Integral in [1] is easily computed when the angle through which the pendulum has risen is given. When  $\theta = \alpha$ ,  $\sin \phi = 1$ , and  $\phi = \frac{\pi}{2}$ ; so that the time of a half-oscillation is  $\sqrt{\frac{a}{g}} F\left(\sin \frac{\alpha}{2}, \frac{\pi}{2}\right)$ , a confirmation of [7]. Art. 176. The construction indicated



in the figure gives the angle  $\phi$ , corresponding to any given arc  $AP$ . For

$$\frac{y}{\frac{1}{2}y_0} = 1 - \cos \angle O'Q,$$

$$\text{and } \sqrt{\frac{y}{y_0}} = \sqrt{\frac{1}{2}(1 - \cos \angle O'Q)} = \sin \frac{\angle O'Q}{2} = \sin \angle ACQ.$$

Therefore  $\angle ACQ = \phi$ .

It is very easy to express the angle  $\theta$  in terms of  $t$ .

We have 
$$t = \sqrt{\frac{a}{g}} F\left(\sin \frac{\alpha}{2}, \phi\right);$$

hence 
$$t \sqrt{\frac{g}{a}} = F\left(\sin \frac{\alpha}{2}, \phi\right),$$

$$\phi = \text{am}\left(t \sqrt{\frac{g}{a}}\right),$$

$$\sin \phi = \text{sn}\left(t \sqrt{\frac{g}{a}}\right),$$

and 
$$\sin \frac{\theta}{2} = \sin \frac{\alpha}{2} \text{sn}\left(t \sqrt{\frac{g}{a}}\right) \pmod{\sin \frac{\alpha}{2}};$$

then 
$$\cos \frac{\theta}{2} = \text{dn}\left(t \sqrt{\frac{g}{a}}\right) \pmod{\sin \frac{\alpha}{2}},$$

and 
$$\sin \theta = 2 \sin \frac{\alpha}{2} \text{sn}\left(t \sqrt{\frac{g}{a}}\right) \text{dn}\left(t \sqrt{\frac{g}{a}}\right) \pmod{\sin \frac{\alpha}{2}}.$$

EXAMPLES.

(1) A pendulum swings through an angle of  $180^\circ$ ; required the time of oscillation.

*Ans.*  $3.708 \sqrt{\frac{a}{g}}$ .

(2) Compare the times required by the pendulum in Ex. (1) to descend through the first  $30^\circ$ , the second  $30^\circ$ , and the third  $30^\circ$  of its arc respectively.

*Ans.*  $1.028 \sqrt{\frac{a}{g}}; 0.446 \sqrt{\frac{a}{g}}; 0.380 \sqrt{\frac{a}{g}}$ .

(3) The time of vibration of a pendulum swinging in an arc of  $72^\circ$  is observed to be 2 seconds; how long does it take it to fall through an arc of  $5^\circ$ , beginning at a point  $20^\circ$  from the highest point of the arc of swing? *Ans.* 0.095 seconds.

(4) A pendulum for which  $\sqrt{\frac{a}{g}}$  has been determined, and is equal to  $\frac{1}{2}$ , vibrates through an arc of  $180^\circ$ ; through what arc does it rise in the first half-second after it has passed its lowest point? in the first  $\frac{1}{8}$  of a second? *Ans.*  $69^\circ$ ;  $20^\circ 6'$ .

(5) It has been shown in Art. 176 that if  $y_0 > 2a$  the pendulum will make complete revolutions, and that the time required to pass from the lowest point to any point whose distance above the lowest point is  $y$ , is

$$t = a \sqrt{\frac{2}{gy_0}} \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = a \sqrt{\frac{2}{gy_0}} F(k, \phi),$$

where  $k = \sqrt{\frac{2a}{y_0}}$  and  $\sin \phi = \sqrt{\frac{y}{2a}}$ .

Show that in this case  $\phi = \frac{\theta}{2}$ , and that  $\sin \frac{\theta}{2} = \operatorname{sn} \left( \frac{t}{a} \sqrt{\frac{gy_0}{2}} \right)$ .

NOTE.—In working with a pendulum it is often about as easy to compute  $F(k, \phi)$  by developing by the binomial theorem and integrating two or three terms, as to use a table of Elliptic Integrals.

We have 
$$F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}},$$

$$(1 - k^2 \sin^2 \phi)^{-\frac{1}{2}} = 1 + \frac{1}{2} k^2 \sin^2 \phi + \frac{1 \cdot 3}{2 \cdot 4} k^4 \sin^4 \phi + \dots,$$

and 
$$F(k, \phi) = \int_0^\phi \frac{d\phi}{\sqrt{1 - k^2 \sin^2 \phi}} = \phi + \frac{k^2}{4} (\phi - \sin \phi \cos \phi) - \frac{3}{32} k^4 \sin^3 \phi \cos \phi + \frac{9}{64} k^4 (\phi - \sin \phi \cos \phi) \dots$$

## CHAPTER XVII.

## INTRODUCTION TO THE THEORY OF FUNCTIONS.

204. A function having but a single value for any given value, real or imaginary, of the variable is called a *single-valued* function. Rational Algebraic Functions, Exponential Functions, the direct Trigonometric Functions, and the Elliptic Functions are single-valued.

A function which has in general two or more values for any given value of the variable is called a *multiple-valued* function. Irrational Algebraic Functions, Logarithmic Functions, the inverse or anti-Trigonometric Functions, and the Elliptic Integrals, are multiple-valued.


205. In Chapter II. we have explained the customary graphical method of representing an imaginary by the position of a point in a plane, the rectangular coördinates of the point being the real term and the real coefficient of the pure imaginary term of the imaginary in question.

In the ordinary treatment of the Theory of Functions this method of representation is of the greatest service, and enables us to bring the study of functions of imaginary variables within the province of Pure Geometry, and to give it great definiteness and precision.

For the sake of brevity we shall in future use the symbol  $i$  for  $\sqrt{-1}$  and  $\text{cis } \phi$  for  $\cos \phi + \sqrt{-1} \sin \phi$ , so that we shall write our typical imaginary as  $x + yi$  or as  $r \text{cis } \phi$ , instead of using the longer forms  $x + y\sqrt{-1}$ , and  $r(\cos \phi + \sqrt{-1} \sin \phi)$ .

We shall also use the name *complex quantity* for an imaginary of the typical form when it is necessary to distinguish it from a pure imaginary.

206. A complex variable  $z = x + yi$  is said to vary *continuously* when it varies in such a manner that the path traced by the point  $(x, y)$  representing it is a continuous line.



Thus if  $z$  changes from the value  $\alpha$  to the value  $\beta$ , so that the point representing it traces any of the four lines in the figure,  $z$  varies continuously.

It will be seen that a variable can pass from the first to the second of two given values, real or imaginary, by any one of an infinite number of different paths without discontinuity if the variable in question is not restricted to real values; while a real variable can change continuously from one given value to another in but one way, since the point representing it is confined in its motion to the *axis of reals*.

207. A single-valued function  $w$  of a complex variable  $z$  is called a *continuous function* if the point representing it traces a continuous path whenever the point representing  $z$  traces a continuous path.

A multiple-valued function of  $z$  is continuous if each of the  $n$  points representing values corresponding to a value of  $z$  traces a continuous path whenever  $z$  traces a continuous path. These  $n$  paths are in general distinct, but two or more of them may intersect, a point of intersection corresponding to a value of  $z$  for which two or more of the  $n$  values of  $w$ , usually distinct, happen to coincide. Such a value of  $z$  is sometimes called a

*critical value*, and the consideration of critical values plays an important part in the Theory of Functions.

In studying a multiple-valued function we may confine our attention to any one of its  $n$  values, and except for the possible presence of critical points this value may be treated just as we treat a single-valued function.

In representing graphically the changes produced in a function  $w$  by changing the variable  $z$  on which it depends, it is customary to avoid confusion by using separate sets of axes for  $w$  and  $z$ .

208. If we use the word *function* in its widest sense,  $w = u + vi$  will be a function of a complex variable  $z = x + yi$ , if  $u$  and  $v$  are any given functions of  $x$  and  $y$ . For example,


$$xi, 6y, x^2 + y^2, x - yi, x^2 - y^2 + 2xyi, \frac{x - y + xi}{\sqrt{x^2 + y^2 + 4}}$$

may all be regarded as functions of  $z$ .

We have seen in Chapter II., Arts. 36-42, that with this definition of function the derivative with respect to  $z$  of a function of  $w$  is in general indeterminate; but that there are various functions of  $z$ , for instance,  $z^m, \log z, e^z, \sin z$ , where the derivative is not indeterminate. We are now ready to investigate more in detail the general question of the existence of a determinate derivative of a function of a complex variable.

Let  $w = u + vi$  be a function of  $z$ ;  $u$  and  $v$ , which are *real*, being functions of  $x$  and  $y$ .

Starting with the value  $z_0 = x_0 + y_0i$  of  $z$  and the corresponding value  $w_0 = u_0 + v_0i$  of  $w$ , let us change  $z$  by giving to  $x$  increment  $\Delta x$  without changing  $y$ .




Let  $\Delta_x u$  and  $\Delta_x v$  be the corresponding increments of  $u$  and  $v$ ; and  $z_1$  and  $w_1$  the new values of  $z$  and  $w$ .

We have  $z_1 = z_0 + \Delta x$ ,  $w_1 = w_0 + \Delta_x u + i \Delta_x v$ .

Then  $\frac{w_1 - w_0}{z_1 - z_0} = \frac{\Delta_x u}{\Delta x} + \frac{i \Delta_x v}{\Delta x}$ ,

and the derivative of  $w$  with respect to  $z$  under the given circumstances is

$$\lim_{\Delta z \rightarrow 0} \left[ \frac{w_1 - w_0}{z_1 - z_0} \right] = D_x u + i D_x v. \quad [1]$$



If, however, starting with the same value  $z_0$  of  $z$ , we change  $z$  by giving  $y$  the increment  $\Delta y$  without changing  $x$ , we have

$$z_1 = x_0 + (y_0 + \Delta y)i = z_0 + i \Delta y,$$

$$w_1 = u_0 + \Delta_y u + (v_0 + \Delta_y v)i = w_0 + \Delta_y v + i \Delta_y v,$$

$$\frac{w_1 - w_0}{z_1 - z_0} = \frac{\Delta_y v}{i \Delta y} + \frac{i \Delta_y v}{i \Delta y},$$

and

$$\lim_{\Delta z \rightarrow 0} \left[ \frac{w_1 - w_0}{z_1 - z_0} \right] = D_y v - i D_y u, \quad [2]$$

and this is the derivative of  $w$  with respect to  $z$  when we change  $y$  and do not change  $x$ .

Comparing [1] with [2], we see that if we start with a given value of  $z$ , and change  $z$  in the two different ways just considered, the limits of the ratios of the corresponding changes in  $w$  to the changes in  $z$  need not be the same. Indeed, the two values for  $\frac{dw}{dz}$  given in [1] and [2] will not be the same unless  $w = u + vi$  is such a function of  $z = x + yi$  that

$$D_x u = D_y v \quad \text{and} \quad D_x v = -D_y u. \quad [3]$$

We shall now show that if  $w$  is such a function of  $z$  that equations [3] are satisfied,  $\lim_{\Delta z \rightarrow 0} \left[ \frac{\Delta w}{\Delta z} \right]$  will be the same if we start with a given value  $z_0$  of  $z$ , no matter in what manner  $z$  may change; that is, no matter in what direction the point representing  $z$  may be supposed to move; or, in other words, no matter what may be the value of  $\lim_{\Delta x \rightarrow 0} \left[ \frac{\Delta y}{\Delta x} \right]$ .

We have in general, since  $w$  is a function of the two variables  $x$  and  $y$ ,

$$\Delta w = (D_x u + iD_x v) \Delta x + (D_y u + iD_y v) \Delta y + \epsilon,$$

where  $\epsilon$  is an infinitesimal of higher order than  $\Delta x$  or  $\Delta y$ .

(I., Art. 198.)

$$\Delta z = \Delta x + i\Delta y.$$

Hence 
$$\frac{\Delta w}{\Delta z} = \frac{D_x u \cdot \Delta x + iD_y v \cdot \Delta y + iD_x v \cdot \Delta x + D_y u \cdot \Delta y + \epsilon}{\Delta x + i\Delta y}$$

$$= \frac{D_x u + iD_y v \cdot \frac{\Delta y}{\Delta x} + iD_x v + D_y u \cdot \frac{\Delta y}{\Delta x} + \frac{\epsilon}{\Delta x}}{1 + i \frac{\Delta y}{\Delta x}},$$

and

$$\lim_{\Delta z \rightarrow 0} \left[ \frac{\Delta w}{\Delta z} \right] = \frac{dw}{dz}$$

$$= \frac{D_x u + iD_y v \cdot \lim_{\Delta x \rightarrow 0} \left[ \frac{\Delta y}{\Delta x} \right] + i \left( D_x v - iD_y u \cdot \lim_{\Delta x \rightarrow 0} \left[ \frac{\Delta y}{\Delta x} \right] \right)}{1 + i \lim_{\Delta x \rightarrow 0} \left[ \frac{\Delta y}{\Delta x} \right]} \quad [4]$$

a value involving  $\lim_{\Delta x \rightarrow 0} \left[ \frac{\Delta y}{\Delta x} \right]$ , and therefore dependent upon the direction in which  $z$  is made to move.

If, however, [3] is satisfied, [4] reduces to

$$\frac{dw}{dz} = D_x u + iD_x v, \quad [5]$$

and the derivative of  $w$  is independent of  $\lim_{\Delta x \rightarrow 0} \left[ \frac{\Delta y}{\Delta x} \right]$ .



A function which satisfies equations [3], and which, therefore, has a derivative whose value depends only upon the value of the independent variable, and not upon the direction in which the point representing the variable is supposed to move, is called by some writers a *monogenic* function, by others a *function which has a derivative*; still others refuse to dignify with the name of function any other than *monogenic* functions.

209. Any function involving  $z$  as a whole, that is, any function which can be formed by performing an operation or a series of operations upon  $z$  as a whole, without introducing  $x$  and  $y$  except as they occur in  $z$ , is a monogenic function of  $z$ .

For if  $w = fz = f(x + yi)$ ,

where  $fz$  can be formed by operating upon  $z$  as a whole,

$$D_x w = f'z, \quad \text{and} \quad D_y w = i f'z;$$

therefore  $i D_x w = D_y w$ , or  $i D_x (u + vi) = D_y (u + vi)$ ;

whence  $D_x u = D_y v$ , and  $D_y u = -D_x v$ ;

and [3], Art. 208, is satisfied. Consequently  $w$  is monogenic. This accounts for the results of Arts. 38–42.

If  $w$  is a multiple-valued function of  $z$ , there may be several different values of  $\frac{dw}{dz}$ , corresponding to the same value of  $z$ ; but if  $w$  is monogenic, each of these values depends only upon  $z$ , and not upon the way in which  $z$  is supposed to change.

In future, unless something is said to the contrary, we shall give the name function only to monogenic functions. Thus we shall not call such expressions as  $x - yi$ , or  $x^2 + y^2 + 2xyi$ , functions of  $z$ .

#### *Conjugate Functions.*

210. If  $u$  and  $v$  are functions of  $x$  and  $y$ , satisfying equations [3], Art. 208, it is easy to prove that

$$D_x^2 u + D_y^2 u = 0 \quad \text{and} \quad D_x^2 v + D_y^2 v = 0.$$

For since  $D_x u = D_y v$  and  $D_x v = -D_y u$ ,  
 we have  $D_x^2 u = D_x D_y v$  and  $D_y^2 u = -D_x D_y v$ ,  
 $D_x^2 v = -D_x D_y u$  and  $D_y^2 v = D_x D_y u$ ;

$u$  and  $v$  are then solutions of Laplace's equation,

$$D_x^2 V + D_y^2 V = 0. \tag{1}$$

Any two functions  $\phi$  and  $\psi$  of  $x$  and  $y$ , such that  $\phi(x, y) + i\psi(x, y)$  is a monogenic function of  $x + yi$ , are called *conjugate functions*; and, by what has just been proved, each of a pair of conjugate functions is always a solution of Laplace's Equation [1].

Thus  $x^2 - y^2$ ,  $2xy$ ;  $e^x \cos y$ ,  $e^x \sin y$ ;  $\frac{1}{2} \log(x^2 + y^2)$ ,  $\tan^{-1} \frac{y}{x}$ ; are three pairs of conjugate functions, since  $x^2 - y^2 + 2xyi = (x + yi)^2$ ,  $e^x \cos y + ie^x \sin y = e^{x+yi}$ ,  $\frac{1}{2} \log(x^2 + y^2) + i \tan^{-1} \frac{y}{x} = \log(x + yi)$ , and consequently, by Art. 209, are all monogenic. Therefore each of the six functions at the beginning of this paragraph is a solution of Laplace's Equation [1].

It is clear that we can form pairs of conjugate functions at pleasure by merely forming functions of  $x + yi$  and breaking them up into their real parts, and their pure imaginary parts; that is, throwing them into the typical form  $u + vi$ .

If each of a pair of conjugate functions,  $\phi$  and  $\psi$ , is written equal to a constant, the equations thus formed will represent a pair of curves which intersect at right angles. For let  $(x, y)$  be a point of intersection of the curves  $\phi = a$ ,  $\psi = b$ ; the slopes of the two curves at  $(x, y)$  are respectively  $-\frac{D_x \phi}{D_y \phi}$ ,  $-\frac{D_x \psi}{D_y \psi}$  by I., Art. 202; and since  $D_x \phi = D_y \psi$  and  $D_x \psi = -D_y \phi$ , the second slope is minus the reciprocal of the first, and the curves are perpendicular to each other at the point in question.

Thus  $x^2 - y^2 = a$ ,  $2xy = b$ , cut each other orthogonally; as do

also  $\frac{1}{2} \log (x^2 + y^2) = a$ ,  $\tan^{-1} \frac{y}{x} = b$ ; or, what amounts to the same thing,  $x^2 + y^2 = a_1 \frac{y}{x} = b_1$ . It must be observed, however, that  $x^2 + y^2$  and  $\frac{y}{x}$  are not conjugate functions, and that in general the converse of our proposition does not hold.

It may be easily proved that if  $\phi$  and  $\psi$  are conjugate functions of  $x$  and  $y$ , and  $f$  and  $F$  are any second pair of conjugate functions of  $x$  and  $y$ , the new pair of functions formed by replacing  $x$  and  $y$  in  $\phi$  and  $\psi$  by  $f$  and  $F$  respectively will be conjugate.

$$\text{Thus} \quad (e^x \cos y)^2 - (e^x \sin y)^2, \quad 2 e^x \cos y \cdot e^x \sin y,$$

$$\text{or, reducing,} \quad e^{2x} \cos 2y, \quad e^{2x} \sin 2y,$$

are conjugate functions ;

$$\frac{1}{2} \log [(x^2 - y^2)^2 + (2xy)^2], \quad \tan^{-1} \left( \frac{2xy}{x^2 - y^2} \right),$$

$$\text{or, reducing,} \quad \log (x^2 + y^2), \quad \tan^{-1} \left( \frac{2xy}{x^2 - y^2} \right),$$

are conjugate.

The properties of conjugate functions given in this article are of great importance in many branches of Mathematical Physics.


#### EXAMPLE.

Show that if  $x'$  and  $y'$  are conjugate functions of  $x$  and  $y$ ,  $x$  and  $y$  are conjugate functions of  $x'$  and  $y'$ .

#### *Preservation of Angles.*

211. If  $w$  is a single-valued monogenic function of  $z$ , and the point representing  $z$  traces two arcs intersecting at a given angle, the corresponding arcs traced by the point representing  $w$  will in general intersect at the same angle.

For let  $z_0$  be the point of intersection of the curves in the  $z$  plane, and  $w_0$  the corresponding point in the  $w$  plane. Let  $z_1$  be a point on the first curve, and  $z_2$  a point on the second; and let



$w_1$  and  $w_2$  be the corresponding points in the  $w$  figure.

Let  $r_1, r_2, s_1,$  and  $s_2$  be the moduli of  $z_1 - z_0, z_2 - z_0, w_1 - w_0,$  and  $w_2 - w_0$  respectively,  $\phi_1, \phi_2, \psi_1,$  and  $\psi_2$  their arguments; then, since  $w$  is a monogenic function of  $z$ , we must have

$$\lim \left[ \frac{w_1 - w_0}{z_1 - z_0} \right] = \lim \left[ \frac{w_2 - w_0}{z_2 - z_0} \right],$$

or 
$$\lim \left[ \frac{s_1 \text{ cis } \psi_1}{r_1 \text{ cis } \phi_1} \right] = \lim \left[ \frac{s_2 \text{ cis } \psi_2}{r_2 \text{ cis } \phi_2} \right];$$

whence, by Art. 23,

$$\lim \left[ \frac{s_1}{r_1} \text{ cis } (\psi_1 - \phi_1) \right] = \lim \left[ \frac{s_2}{r_2} \text{ cis } (\psi_2 - \phi_2) \right];$$

and since, when two imaginaries are equal, their moduli must be equal, and their arguments must be equal, unless the moduli are both zero or both infinite,

$$\lim (\psi_2 - \psi_1) = \lim (\phi_2 - \phi_1);$$

that is, the angle between the arcs in the  $w$  figure is equal to the angle between the corresponding arcs in the  $z$  figure; unless

$$\left[ \frac{dw}{dz} \right]_{z=z_0} = 0, \text{ or } \left[ \frac{dw}{dz} \right]_{z=z_0} = \infty.$$

If  $w$  is a multiple-valued monogenic function of  $z$ . and if starting from any point  $z_0$ , the point which represents  $z$  traces

out two curves intersecting at an angle  $\alpha$ , each of the  $n$  points representing the corresponding values of  $w$  will trace out a pair of curves intersecting at the angle  $\alpha$ ; unless  $z_0$  is a point at which  $\frac{dw}{dz}$  is zero or infinite.

If, then,  $w$  is any monogenic function of  $z$ , and the point representing  $z$  is made to trace out any figure however complex, the point representing  $w$  will trace out a figure in which all the angles occurring in the  $z$  figure are preserved unchanged, except those having their vertices at points representing values of  $z$  which make  $\frac{dw}{dz}$  zero or infinite.

This principle leads to the following working rule for transforming any given figure into another, in which the angles are preserved unchanged.

Substitute  $x'$  and  $y'$  for  $x$  and  $y$  in the equations of the curves which compose the given figure,  $x'$  and  $y'$  being any pair of conjugate functions (Art. 210) of  $x$  and  $y$ , and the new equations thus obtained will represent a set of curves forming a second figure in which all the angles of the given figure are preserved unchanged, except those having their vertices at points at which  $D_x x'$  and  $D_y y'$  are both zero, or at which one of them is infinite.

For example,  $x - y = a,$  (1)

$$x + y = b, \quad (2)$$

are a pair of perpendicular right lines. Replace  $x$  by  $x^2 - y^2$  and  $y$  by  $2xy$ , and we get

$$x^2 - 2xy - y^2 = a, \quad (3)$$

$$x^2 + 2xy - y^2 = b, \quad (4)$$

a pair of hyperbolas that cut orthogonally.

212. If  $w$  is a *single-valued* continuous function of  $z$ , it is clear that if  $w_0$  and  $w_1$  are the values corresponding to  $z_0$  and  $z_1$ ,



and the point  $z$  moves from  $z_0$  to  $z_1$  by two different paths, the corresponding paths traced by  $w$  will begin at  $w_0$  and end at  $w_1$ , and consequently that if  $z$  describes any closed contour,  $w$  also will describe a closed contour.

If  $w$  is a double-valued function of  $z$ , since to each value of  $z$  there will correspond two values of  $w$ , it is conceivable that if  $w_1$  and  $w_1'$  are the values of  $w$  corresponding to  $z_1$ , and  $z$  moves from  $z_0$  to  $z_1$  by two different paths,  $w$  may in one case move from  $w_0$  to  $w_1$ , and in the other case from  $w_0$  to  $w_1'$ .

We can prove, however, that if the two paths traced by  $z$  do not enclose a *critical point* (Art. 207), and  $w$  is finite and continuous for the portion of the plane considered, this will not take place, and that the two paths starting from  $w_0$  will terminate at the same point  $w_1$ .

For as  $z$  traces the first path, each of the two points representing the two values of  $w$  will trace a path, one starting at  $w_0$ , and the other at  $w_0'$ , and unless the  $z$  path passes through a critical point, the two  $w$  paths will not intersect, but will be entirely separate and distinct, and will lead, one from  $w_0$  to  $w_1$ , the other from  $w_0'$  to  $w_1'$ .

If, now, the  $z$  path be gradually swung into a second position without changing its beginning or its end, since  $w$  is a continuous function, the two  $w$  paths will be gradually swung into new positions; but, provided that the  $z$  path in its changing does not at any time pass through a critical point, the two  $w$  paths will at no time intersect, and consequently it will be impossible for the  $w$  points to pass over from one path to the other, and therefore the point which starts at  $w_0$  must always come out at  $w_1$ , and not at  $w_1'$ .

It follows readily from this reasoning that if  $z$  describes a closed contour not embracing a critical point, each of the  $w$  points will describe a closed contour, and these contours will not intersect.

Of course, the proof given above holds for any multiple-valued function.

In any portion of the plane, then, not containing critical

points the separate values of a multiple-valued function may be separately considered, and may be regarded and treated as single-valued functions.

213. That in the case of a double-valued function two paths in the  $z$  plane, including between them a critical point, but having the same beginning and the same end, may lead to different values of the function, is easily shown by an example.

Let  $w = \sqrt{z}$ , and let  $z$ , starting with the value 1, move to the value  $-1$  by the semi-circular path in the figure. That one of




FIG. 1.

the corresponding values of  $w$  which starts with  $+1$  will describe the quadrant shown in the figure, and will reach the point  $1 \cdot \text{cis} \frac{\pi}{2}$ , or  $i$ . If, however,  $z$  moves from  $+1$  to  $-1$  by




FIG. 2.

the semi-circular path in the second figure, the value of  $w$  which starts with  $+1$  will describe the quadrant shown in the second figure, and will reach the value  $1 \cdot \text{cis} \left(-\frac{\pi}{2}\right)$ , or  $-i$ . These

two paths described by  $z$ , then, although beginning at the same point  $+1$  and ending at the same point  $-1$ , cause that value of the function which begins with  $+1$  to reach two different values; and the two paths in question embrace the point  $z = 0$ , which is clearly a point at which the two values of  $w$ , ordinarily different, coincide; that is, a critical point.

It is easily seen that if  $z$ , starting with the value  $+1$ , describes a complete circumference about the origin, the value of  $w$  which starts from the point  $+1$  will not describe a closed contour, but will move through a semi-circumference and end with the point  $1 \cdot \text{cis } \pi$  or  $-1$ . Now, by Art. 212 any path





FIG. 3.


described by  $z$  beginning with  $+1$  and ending with  $-1$  and passing above the origin, since it can be deformed into the semi-circumference of Fig. 1 without passing through a critical point, will cause the value of  $w$  beginning with  $+1$  to end with  $+i$ ; and any path described by  $z$  beginning with  $+1$  and ending with  $-1$  and passing below the origin, since it can be deformed into the semi-circumference of Fig. 2 without passing through a critical point, will cause the value of  $w$  beginning with  $+1$  to end with  $-i$ . Therefore any two paths described by  $z$  beginning with  $+1$  and ending with  $-1$  will, if they include the critical point  $z = 0$  between them, lead to different values of  $w$ , provided that the same value of  $w$  is taken at the start.

214. If  $w$  is a double-valued function of  $z$ , and  $z$  describes a closed contour about a single critical point, this contour may be deformed into a circle about the critical point, and a line leading from the starting point to the circumference of the circle, without affecting the final value of  $w$  (Art. 212). Thus, in the figure, the two paths  $ABCD A$ ,  $AB'C'D'B'A$  lead from the same initial to the same final value of  $w$ ; and this is true no matter how small the radius of the circle  $B'C'D'$ .






Let  $z_0$  be the critical point, and let  $w_0$  be the corresponding point in the  $w$  figure. As  $z$  moves from  $z_1$  towards  $z_0$ , the points



representing the corresponding values of  $w$  will start at  $w_1$  and  $w_1'$  and move towards  $w_0$ , tracing distinct paths.

If, now,  $z$  describes a circumference about  $z_0$ , and then returns along its original path to  $z_1$ , the first value of  $w$  will either make a complete revolution about  $w_0$  and return along the branch (1) to its initial value  $w_1$ , or it will describe about



$w_0$  a path ending with the branch (2) of the  $w$  curve, and move along that branch to the value  $w_1'$ .

In the first case, and in that case only, the value of  $w$  describes a closed contour when  $z$  describes a closed contour, and is practically a single-valued function.

If  $z_0$  is a point at which  $\frac{dw}{dz}$  is neither zero nor infinite (v. Art. 211), when  $z$  describes about  $z_0$  a circle of infinitesimal radius,  $w$  will make about  $w_0$  a complete revolution; for since if two radii are drawn from  $z_0$ , the curves corresponding to them will form at  $w_0$  an angle equal to the angle between the radii, when a radius drawn to the moving point which is describing the circle about  $z_0$  revolves through an angle of  $360^\circ$ , the cor-

responding line joining  $w_0$  with the moving point representing  $w$  will revolve through  $360^\circ$ , and we shall have what we have called Case I.

If, then, we avoid the points at which  $\frac{dw}{dz}$  is zero or infinite, we shall avoid all critical points that can vitiate the results obtained by treating our double-valued or multiple-valued functions as we treat single-valued functions.

A *critical point* of such a character that when  $z$  describes a closed contour about it the corresponding path traced by any one of the values of  $w$  is not closed, we shall call a *branch point*.

When a function is finite, continuous, and single-valued for all values of  $z$  lying in a given portion of the  $z$  plane, or when if multiple-valued it is finite and continuous, and has no *branch points* in the portion of the plane in question, it is said to be *holomorphic* in that portion of the plane.

*Definite Integrals.*

215. In the case of real variables,  $\int_{z_0}^z fz . dz$  was defined in Art. 80 in effect as follows :

$$\int_{z_0}^z fz . dz = \lim_{n = \infty} [fz_0(z_1 - z_0) + fz_1(z_2 - z_1) + fz_2(z_3 - z_2) + \dots + fz_{n-1}(Z - z_{n-1})], \quad [1]$$

where  $z_1, z_2, z_3, \dots z_{n-1}$  are values of  $z$  dividing the interval between  $z_0$  and  $Z$  into  $n$  parts, each of which is made to approach zero as its limit as  $n$  is indefinitely increased.

In other words,  $\int_{z_0}^z$  is the *line integral* of  $fz$  (Art. 163) taken along the straight line, joining  $z_0$  and  $Z$  if  $z_0$  and  $Z$  are represented as in the Calculus of Imaginaries.

It has been proved that if  $fz$  is finite and continuous between  $z_0$  and  $Z$ , this integral depends merely upon the initial and final values of  $z$ , and is equal to  $FZ - Fz_0$  where  $Fz$  is the indefinite integral  $\int fz . dz$ .

If  $z$  is a *complex* variable, and passes from  $z_0$  to  $Z$  along any given path, we shall still define the definite integral  $\int_{z_0}^Z fz . dz$  by [1] where now  $z_1, z_2, z_3, \dots, z_{n-1}$  are points in the given path.

Two important results follow immediately from this definition :

$$1st. \text{ That } \int_r^{z_0} fz . dz = - \int_{z_0}^r fz . dz, \quad [2]$$

if  $z$  traverses in each integral the same path connecting  $z_0$  and  $Z$ .

2d. That the modulus of  $\int_{z_0}^Z fz . dz$  is not greater than the line-integral of the modulus of  $fz$  taken along the given path joining  $z_0$  and  $Z$ .

If we let

$$fz = w = u + vi, \quad z = x + yi, \quad u = \phi(x, y), \quad \text{and} \quad v = \psi(x, y),$$

$$\begin{aligned} \text{then} \quad \int_{z_0}^Z fz . dz &= \int (u + vi)(dx + idy) \\ &= \int \phi(x, y) dx + i \int \psi(x, y) dx - \int \psi(x, y) dy + i \int \phi(x, y) dy, \end{aligned} \quad [3]$$

each of the integrals in the last member being the line-integral of a real function of real variables, taken along the given path connecting  $z_0$  and  $Z$ .

If the given path is changed, each of the integrals in the last member of [3] will in general change, and the value of  $\int_{z_0}^Z fz . dz$  will change ; and, since  $z$  may pass from  $z_0$  to  $Z$  by an infinite number of different paths, we have no reason to expect that  $\int_{z_0}^Z fz . dz$  will in general be determinate.

We shall, however, prove that in a large and important class of cases  $\int_{z_0}^Z fz . dz$  is determinate, and depends for its value upon  $z_0$  and  $Z$ , and not at all upon the nature of the path traversed by  $z$  in going from  $z_0$  to  $Z$ .

216. If  $fz$  is holomorphic in a given portion of the plane,

$$\int_{z_0}^{z_0} fz \cdot dz = 0 \tag{1}$$

if  $z$  describes any closed contour lying wholly within that portion of the plane.

From [3], Art. 215, we have

$$\int_{z_0}^{z_0} fz \cdot dz = \int w \cdot dz = \int u dx + i \int v dx - \int v dy + i \int u dy, \tag{2}$$

the integral in each case being the line-integral around the closed contour in question.

Since  $w = fz$  is holomorphic,  $u = \phi(x, y)$ , and  $v = \psi(x, y)$ , and  $D_x u$ ,  $D_y u$ ,  $D_x v$ , and  $D_y v$  are easily seen to be finite, continuous, and single-valued in the portion of the plane considered. Therefore, by Art. 170,

$$\begin{aligned} \int u dx &= \iint D_y u dx dy; & \int v dx &= \iint D_y v dx dy; \\ \int v dy &= -\iint D_x v dx dy; & \int u dy &= -\iint D_x u dx dy; \end{aligned}$$


the integral in the first member of each equation being taken around the contour, and that in the second member being a surface-integral taken over the surface bounded by the contour.

We have, then, from [2],

$$\int_{z_0}^{z_0} fz \cdot dz = \iint (D_y u + D_x v) dx dy + i \iint (D_y v - D_x u) dx dy, \tag{3}$$

but  $D_x u = D_y v$ , and  $D_y u = -D_x v$  from [3], Art. 208. Therefore, [3] reduces to  $\int_{z_0}^{z_0} fz \cdot dz = 0$ .


From this result we get easily the very important fact that if  $fz$  is holomorphic in a given portion of the plane,  $\int_{z_0}^z fz \cdot dz$  will have the same value for all paths leading from  $z_0$  to  $Z$ , provided they lie wholly in the given part of the plane. For let  $z_0 a Z$  and  $z_0 b Z$  be any two paths not intersecting between  $z_0$  and  $Z$ . Then  $z_0 a Z b z_0$  is a closed contour, and



$$\int_{z_0}^{z_0} fz . dz \text{ (along } z_0 a Z b z_0) \\ = \int_{z_0}^z fz . dz \text{ (along } z_0 a Z) + \int_z^{z_0} fz . dz \text{ (along } Z b z_0) = 0 ;$$

but  $\int_{z_0}^{z_0} fz . dz \text{ (along } Z b z_0) = - \int_{z_0}^z fz . dz \text{ (along } z_0 b Z)$   
by Art. 215.

Therefore,  $\int_{z_0}^z fz . dz \text{ (along } z_0 a Z) = \int_{z_0}^z fz . dz \text{ (along } z_0 b Z)$ .



If the paths  $z_0 a Z$  and  $z_0 b Z$  intersect, a third path  $z_0 c Z$  may be drawn not intersecting either of them, and by the proof just given


$$\int_{z_0}^z fz . dz \text{ (along } z_0 a Z) = \int_{z_0}^z fz . dz \text{ (along } z_0 c Z),$$

$$\int_{z_0}^z fz . dz \text{ (along } z_0 b Z) = \int_{z_0}^z fz . dz \text{ (along } z_0 c Z) ;$$

therefore,

$$\int_{z_0}^z fz . dz \text{ (along } z_0 a Z) = \int_{z_0}^z fz . dz \text{ (along } z_0 b Z).$$

217. If  $fz$ , while in other respects holomorphic in a given portion of the plane, becomes infinite for a value  $T$  of  $z$ , then  $\int fz . dz$  taken around a closed contour embracing  $T$ , while not zero, is, however, equal to the integral taken around any other closed path surrounding  $T$ .



For let  $ABCD$  be any closed contour about  $T$ . With  $T$  as a centre, and a radius  $\epsilon$ , describe a circumference, taking  $\epsilon$  so small that the circumference lies wholly within  $ABCD$ . Join the two contours by a line  $AA'$ . Then  $ABCDAA'D'C'B'A'A$  is a closed path within which  $fz$  is holomorphic.

Therefore,

$$\int fz \cdot dz \text{ (along } ABCDAA'D'C'B'A'A) = 0,$$

or  $\int fz \cdot dz \text{ (along } ABCDA) + \int fz \cdot dz \text{ (along } AA')$   
 $+ \int fz \cdot dz \text{ (along } A'D'C'B'A') + \int fz \cdot dz \text{ (along } A'A) = 0;$

but

$$\int fz \cdot dz \text{ (along } AA') = - \int fz \cdot dz \text{ along } (A'A),$$

and

$$\int fz \cdot dz \text{ (along } A'D'C'B'A') = - \int fz \cdot dz \text{ (along } A'B'C'D'A').$$

Hence

$$\int fz \cdot dz \text{ (along } ABCDA) = \int fz \cdot dz \text{ (along } A'B'C'D'A').$$

218. That the integral of a function of  $z$  around a closed contour embracing a point at which the function is infinite is not necessarily zero is easily shown by an example.

$fz = \frac{1}{z-t}$ ,  $t$  being a given constant, is single-valued, continuous, and finite throughout the whole of the plane except at the point  $t$ , at which  $\frac{1}{z-t}$  becomes infinite, without, however, ceasing to be single-valued.


Let us take  $\int \frac{dz}{z-t}$  around a circle whose centre is  $t$ , and whose radius is any arbitrarily chosen value  $\epsilon$ . If  $z$  is on the circumference of this circle

$$z - t = \epsilon (\cos \phi + i \sin \phi)$$

$$= \epsilon e^{i\phi} \quad \text{by [5], Art. 31.}$$

$$z = t + \epsilon e^{i\phi}$$

and  $dz = i\epsilon e^{i\phi} d\phi.$



Hence 
$$\int \frac{dz}{z-t} \text{ (around } abc) = \int_0^{2\pi} \frac{i\epsilon e^{\phi i} d\phi}{\epsilon e^{\phi i}} = 2\pi i.$$

From what has been proved in Art. 217, it follows that  $\int \frac{dz}{z-t}$  around any closed contour embracing  $t$  must also be equal to  $2\pi i$ .

As another example let us consider  $\int \frac{Fz}{z-t} dz$ , when  $Fz$  is supposed to be holomorphic in the portion of the plane considered, and where the integral is to be taken around any closed contour embracing the point  $z = t$ .

$\frac{Fz}{z-t}$  is holomorphic except at the point  $z = t$ , where it becomes infinite. The required integral is, then, equal to the integral around a circumference described from the point  $t$  as a centre, with any given radius  $\epsilon$ , that is, by the reasoning just used in the case of  $\int \frac{dz}{z-t}$ , to

$$\int_0^{2\pi} \frac{F(t + \epsilon e^{\phi i}) i\epsilon e^{\phi i} d\phi}{\epsilon e^{\phi i}}, \text{ or } i \int_0^{2\pi} F(t + \epsilon e^{\phi i}) d\phi;$$

and in this expression  $\epsilon$  may be taken at pleasure. If now  $\epsilon$  is made infinitesimal  $\epsilon e^{\phi i}$  is infinitesimal, and since  $Fz$  is continuous  $F(t + \epsilon e^{\phi i})$  is equal to  $Ft + \eta$  where  $\eta$  is some infinitesimal, and  $F(t + \epsilon e^{\phi i}) d\phi$  is equal to  $Ft \cdot d\phi + \eta \cdot d\phi$ .

Now, by I. Art. 161,

$$\int_0^{2\pi} (Ft \cdot d\phi + \eta d\phi) = \int_0^{2\pi} Ft \cdot d\phi.$$

Hence 
$$i \int_0^{2\pi} F(t + \epsilon e^{\phi i}) d\phi = i \int_0^{2\pi} Ft \cdot d\phi = 2\pi i Ft;$$

and we get the important result that  $\int \frac{Fz}{z-t} dz$ , taken around any contour including the point  $z = t$ , is equal to  $2\pi i \cdot Ft$ .

From this we have 
$$Ft = \frac{1}{2\pi i} \int \frac{Fz}{z-t} dz;$$

and we see that a holomorphic function is determined everywhere inside a closed contour if its value is given at every point of the contour.

If in the formula 
$$Ft = \frac{1}{2\pi i} \int \frac{Fz}{z-t} dz \tag{1}$$

we change  $t$  to  $t + \Delta t$ , we get

$$\Delta Ft = \frac{1}{2\pi i} \int Fz \cdot dz \left( \frac{1}{z-t-\Delta t} - \frac{1}{z-t} \right) = \frac{1}{2\pi i} \int \frac{Fz \cdot dz \cdot \Delta t}{(z-t)(z-t-\Delta t)};$$

whence

$$\lim_{\Delta t \rightarrow 0} \left[ \frac{\Delta Ft}{\Delta t} \right] = \frac{1}{2\pi i} \int Fz \cdot dz \cdot \lim_{\Delta t \rightarrow 0} \left[ \frac{1}{(z-t)(z-t-\Delta t)} \right],$$

or 
$$F't = \frac{1}{2\pi i} \int \frac{Fz \cdot dz}{(z-t)^2}; \tag{2}$$

and in like manner we get

$$F''t = \frac{2!}{2\pi i} \int \frac{Fz \cdot dz}{(z-t)^3}; \tag{3}$$

and in general 
$$F^{(n)}t = \frac{n!}{2\pi i} \int \frac{Fz \cdot dz}{(z-t)^{n+1}}, \tag{4}$$

each of the integrals in these formulas being taken around a closed contour lying wholly in that portion of the plane in which  $Fz$  is holomorphic, and enclosing the point  $z = t$ .

219. *The integral of a holomorphic function along any given path is finite and determinate*, for, by [3], Art. 215, it is equal to the sum of four line integrals, each of which is finite and determinate (Art. 166).

*If a series  $w_0 + w_1 + w_2 + \dots$ , where  $w_0, w_1, w_2 \dots$  are holomorphic functions of  $z$ , is convergent for all values of  $z$  in a certain portion of the plane, the integral of the series along any given path lying in that portion of the plane is the series formed of the integrals of the terms of the given series along the path in question, and the new series is convergent.*



$$\begin{aligned} \text{For let } S &= w_0 + w_1 + w_2 + \dots + w_n + w_{n+1} + \dots \\ &= w_0 + w_1 + w_2 + \dots + w_n + R, \end{aligned}$$

$$\text{where } R = w_{n+1} + w_{n+2} + \dots,$$

$$\text{and where by hypothesis } \lim_{n=\infty} [R] = 0. \quad (1)$$

$$\int Sdz = \int w_0 dz + \int w_1 dz + \dots + \int w_n dz + \int R. dz$$

for any given value of  $n$ .

By the proposition at the beginning of this article,  $\int Sdz$  along the given path is finite and determinate, as are also

$$\int w_0 dz, \int w_1 dz, \text{ etc.}$$

If, now,  $n$  is indefinitely increased,

$$\int Sdz = \int w_0 dz + \int w_1 dz + \int w_2 dz + \dots + \lim_{n=\infty} \int Rdz.$$

The modulus of  $\int Rdz$  is not greater than the line-integral along the given path of the modulus of  $R$  (v. Art. 215). As  $n$  increases each value of the modulus of  $R$  approaches zero as its limit (1); consequently each element of the cylindrical surface representing the line-integral of the modulus of  $R$  (v. Art. 166) approaches zero, and  $\lim_{n=\infty} \int Rdz = 0$ .

$$\text{Therefore, } \int Sdz = \int w_0 dz + \int w_1 dz + \int w_2 dz + \dots; \quad [2]$$

and, since the first member is finite and determinate, the second member is a *convergent series*.

*Taylor's and Maclaurin's Theorems.*

$$220. \quad \frac{1-q^n}{1-q} = 1 + q + q^2 + q^3 + \dots + q^{n-1}$$

identically, if  $n$  is a positive integer, even when  $q$  is imaginary.


If the modulus of  $q$  is less than 1,

$$\lim_{n \rightarrow \infty} [q^n] = 0.$$

Hence  $1 + q + q^2 + q^3 + \dots = \lim_{n \rightarrow \infty} \left[ \frac{1 - q^n}{1 - q} \right] = \frac{1}{1 - q}$ , [1]

even when  $q$  is imaginary, provided that the modulus of  $q$  is less than 1.

Suppose, now, that everywhere within and on a certain circumference described with the point  $z = a$  as a centre  $Fz$  is holomorphic. Let  $z = t$  be any point within this circumference, and  $z = Z$  be a point on the circumference. Then the modulus of  $Z - a$  is the distance from  $a$  to  $Z$ , and the modulus of  $t - a$  is the distance from  $a$  to  $t$ ;



hence  $\text{mod}(t - a) < \text{mod}(Z - a)$ ,

and  $\text{mod}\left(\frac{t - a}{Z - a}\right) < 1$ .

$$\begin{aligned} \frac{1}{Z - t} &= \frac{1}{Z - a - (t - a)} = \frac{1}{Z - a} \cdot \frac{1}{1 - \frac{t - a}{Z - a}}, \\ &= \frac{1}{Z - a} \left[ 1 + \frac{t - a}{Z - a} + \frac{(t - a)^2}{(Z - a)^2} + \frac{(t - a)^3}{(Z - a)^3} + \dots \right], \end{aligned}$$

by [1]

Hence

$$\frac{1}{Z - t} = \frac{1}{Z - a} + \frac{t - a}{(Z - a)^2} + \frac{(t - a)^2}{(Z - a)^3} + \frac{(t - a)^3}{(Z - a)^4} + \dots, [2]$$

and the second member of [2] is a convergent series.

Multiply [2] by  $\frac{FZ}{2\pi i}$ , and the series will still be convergent for each value of  $z$  which we have to consider; we get

$$\begin{aligned} &\frac{1}{2\pi i} \frac{FZ}{Z - t} \\ &= \frac{1}{2\pi i} \left[ \frac{FZ}{Z - a} + (t - a) \frac{FZ}{(Z - a)^2} + (t - a)^2 \frac{FZ}{(Z - a)^3} + \dots \right]. [3] \end{aligned}$$

Integrate now both members of [3] around the circumference, and we have

$$\frac{1}{2\pi i} \int \frac{FZ}{Z-t} dZ = \frac{1}{2\pi i} \left[ \int \frac{FZ}{Z-a} dZ + (t-a) \int \frac{FZ}{(Z-a)^2} dZ + (t-a)^2 \int \frac{FZ}{(Z-a)^3} dZ + \dots \right]; \quad [4]$$

and, since each of the functions to be integrated is holomorphic on the contour around which the integral is taken, and the second member of [3] is convergent, each integral will be finite and determinate, and the second member of [4] will be convergent.

Substituting in [4] the values obtained in Art. 218, [1], [2], [3], and [4], we have

$$Ft = Fa + (t-a)F'a + \frac{(t-a)^2}{2!}F''a + \frac{(t-a)^3}{3!}F'''a + \dots + \frac{(t-a)^n}{n!}F^{(n)}a + \dots \quad [5]$$

If the point  $z = a$  is at the origin,  $a = 0$  and [5] becomes

$$Ft = Fo + tF'o + \frac{t^2}{2!}F''o + \frac{t^3}{3!}F'''o + \dots, \quad [6]$$

which is Maclaurin's Theorem.

That [5] is merely a new form of Taylor's Theorem is easily seen if we let  $t - a = h$ , whence  $t = a + h$ , and [5] becomes

$$F(a+h) = Fa + hF'a + \frac{h^2}{2!}F''a + \frac{h^3}{3!}F'''a + \dots \quad [7]$$

[6] can, of course, be written

$$Fz = Fo + zF'o + \frac{z^2}{2!}F''o + \frac{z^3}{3!}F'''o + \dots, \quad [8]$$

and [5] as

$$Fz = Fa + (z-a)F'a + \frac{(z-a)^2}{2!}F''a + \frac{(z-a)^3}{3!}F'''a + \dots; \quad [9]$$

and we get the very important result that if a function of  $z$  is holomorphic within a circle whose centre is at the origin it may be developed by Maclaurin's Theorem, and the development will hold, that is, the series will be convergent, for all values of  $z$  lying within the circle.

If a function of  $z$  is holomorphic within a circle described from  $z=a$  as a centre it can be developed by Taylor's Theorem into a series arranged according to powers of  $z-a$ , and the development will hold for all values of  $z$  lying within the circle.

The question of the convergency of either Taylor's or Maclaurin's Series for the case when  $z$  lies on the circumference of the circle needs special investigation, and will not be considered here.

If the function which we wish to develop is single-valued, in drawing our circle of convergence we need avoid only those points at which the function becomes infinite; but if it is multiple-valued we must avoid also those at which its derivative is zero or infinite (v. Art. 214).

221. We are now able to investigate from a new point of view the question of the convergence of the series obtained by Taylor's and Maclaurin's Theorems in I. Chap. IX.

Let us begin with the *Binomial Theorem*,

$$(a+h)^n = a^n + na^{n-1}h + n \frac{(n-1)}{2!} a^{n-2}h^2 + \dots, \quad [1]$$

or, following the notation of [9], Art. 220,

$$z^n = a^n + na^{n-1}(z-a) + \frac{n(n-1)}{2!} a^{n-2}(z-a)^2 + \dots. \quad [2]$$

If  $n$  is a positive integer,  $z^n$  is holomorphic throughout the whole plane, and [2] holds for all values of  $z$  and  $a$ , and [1] for all values of  $a$  and  $h$ .

If  $n$  is a negative integer,  $z^n$  is single-valued, and it is finite and continuous except for  $z=0$ , where  $z^n$  becomes infinite. [2] is, then, convergent for all values of  $z$  lying within a circle described with  $a$  as a centre and passing through the origin;

that is, for all values of  $z$ , such that  $\text{mod}(z - a) < \text{mod } a$ ; and consequently [1] holds if  $\text{mod } h < \text{mod } a$ .

If  $n$  is a fraction,  $z^n$  is multiple-valued, and our circle of convergence must avoid the points at which  $\frac{dz^n}{dz}$  becomes zero or infinite; but as the origin is the only point of this character, the circle of convergence is the same as in the case last considered, and [1] holds for all cases where  $\text{mod } h < \text{mod } a$ .

When  $a$  and  $h$  are real our results agree with those obtained in I. Art. 131.

$$(b) \quad e^x = e^{x+yi} = e^x (\cos y + i \sin y) \quad ([4], \text{ Art. } 31)$$

is single-valued and continuous, and becomes infinite only when  $x = \infty$ . Maclaurin's development for  $e^x$  holds, then, for all finite values of  $z$ .

$$(c) \quad \log z = \log(r \text{ cis } \phi) = \log r + \phi i \quad (\text{Art. } 33)$$

is finite and continuous throughout the whole plane. It is, however, multiple-valued, but its derivative  $\frac{1}{z}$  becomes infinite only when  $z = 0$ , and does not become zero for any finite value of  $z$ .  $\log z$ , then, can be developed into a convergent series, arranged according to powers of  $z - a$ , for all values of  $z$  within a circle having the centre  $a$  and passing through the origin; that is, for all cases where  $\text{mod}(z - a) < \text{mod } a$ .

If  $z - a = h$ , we get

$$\log(a + h) = \log a + \frac{h}{a} - \frac{h^2}{2a^2} + \frac{h^3}{3a^3} - \frac{h^4}{4a^4} + \dots, \quad [3]$$

[3] holding for all cases where  $\text{mod } h < \text{mod } a$ .

If  $a = 1$  and  $h = z$ , we get

$$\log(1 + z) = \frac{z}{1} - \frac{z^2}{2} + \frac{z^3}{3} - \frac{z^4}{4} + \dots, \quad [4]$$

which holds for all values of  $z$  where  $\text{mod } z < 1$ .

$$(d) \quad \sin z = \sin(x + yi) = \sin x \cdot \frac{e^y + e^{-y}}{2} + i \cos x \cdot \frac{e^y - e^{-y}}{2},$$

and  $\cos z = \cos(x + yi) = \cos x \cdot \frac{e^y + e^{-y}}{2} - i \sin x \cdot \frac{e^y - e^{-y}}{2}$ ,  
(v. [3] and [4], Art. 35)

are single-valued, and are finite and continuous throughout the plane. Therefore, Maclaurin's developments for  $\sin z$  and  $\cos z$  hold for all values of  $z$ .

(e)  $\tan z = \frac{\sin z}{\cos z}$ , and  $\sec z = \frac{1}{\cos z}$ , are single-valued and continuous, and become infinite only when  $\cos z = 0$ ; that is, when  $z = \frac{\pi}{2}$ . Therefore, Maclaurin's developments for  $\tan z$  and  $\sec z$  (I. Art. 138), hold for every value of  $z$  whose modulus is less than  $\frac{\pi}{2}$ .

(f)  $\csc z = \frac{1}{\sin z}$ , and  $\cot z = \frac{\cos z}{\sin z}$  become infinite when  $z = 0$ , and cannot be developed by Maclaurin's Theorem.

(g)  $\sin^{-1} z$  is finite and continuous throughout the plane; it is, however, multiple-valued, and its derivative  $\frac{1}{\sqrt{1-z^2}}$  becomes infinite when  $z = 1$ , and when  $z = -1$ . Therefore, the development for  $\sin^{-1} z$  (I. Art. 135 [2]), holds for any value of  $z$  whose modulus is less than 1.

(h)  $\tan^{-1} z$  is finite and continuous throughout the plane; it is multiple-valued, and its derivative  $\frac{1}{1+z^2}$  becomes infinite when  $z = i$ , and when  $z = -i$ . Therefore, the development for  $\tan^{-1} z$  (I. Art. 135 [1]), holds if  $\text{mod } z < \text{mod } i$ ; that is, if  $\text{mod } z < 1$ .

#### EXAMPLES.

(1) Show that the development of  $\frac{z}{1+z} + \log(1+z)$ , given in I. Art. 136, Ex. 1, holds if  $\text{mod } z < 1$ .

(2) Show that the development of  $\log(1+e^z)$ , given in I. Art. 136, Ex. 2, holds if  $\text{mod } z < \pi$ .

(3) Obtain the following developments, and find for what real values of  $x$  they hold good:

- (a)  $\log(x + \sqrt{x^2 + a^2}) = \log a + \frac{x}{a} - \frac{1}{2} \cdot \frac{x^3}{3a^3} + \frac{1.3}{2.4} \cdot \frac{x^5}{5a^5} - \dots$
- (b)  $(e^x + e^{-x})^n = 2^n \left( 1 + \frac{nx^2}{2!} + n(3n-2) \frac{x^4}{4!} + \dots \right)$
- (c)  $e^x \cdot \cos x = 1 + x - \frac{2x^3}{3!} - \frac{4x^4}{4!} \dots$
- (d)  $e^{\cos x} = e \left( 1 - \frac{x^2}{2!} + \frac{4x^4}{4!} - \frac{31x^6}{6!} \dots \right)$
- (e)  $e^x \log(1+x) = x + \frac{x^2}{2!} + \frac{2x^3}{3!} + \frac{9x^4}{5!} \dots$
- (f)  $\tan^4 x = x^4 + \frac{4x^5}{3} + \frac{6x^6}{5} \dots$
- (g)  $(1 + 2x + 3x^2)^{-\frac{1}{2}} = 1 - x + 2x^2 - \frac{7x^4}{4} \dots$
- (h)  $e^{\tan^{-1} x} = 1 + x + \frac{x^2}{2} - \frac{x^3}{6} \dots$
- (i)  $\log\left(\frac{1+x}{1-x}\right) = 2 \left( x + \frac{x^3}{3} + \frac{x^5}{5} \dots \right)$
- (j)  $\tan x = x + \frac{2}{3!}x^3 + \frac{2^4}{5!}x^5 + \frac{2^4 \cdot 17}{7!}x^7 + \frac{2^8 \cdot 31}{9!}x^9 + \frac{2^9 \cdot 691}{11!}x^{11} \dots$
- (k)  $x \cdot \text{ctn } x = 1 - \frac{x^2}{3} - \frac{x^4}{45} - \frac{2x^6}{945} - \frac{x^8}{4725} - \frac{2x^{10}}{93555} \dots$
- (l)  $\log \tan\left(\frac{\pi}{4} + x\right) = \log \tan \frac{\pi}{4} + 2x + \frac{2^3}{3!}x^3 + \frac{5 \cdot 2^5}{5!}x^5 + \frac{61 \cdot 2^7}{7!}x^7 \dots$
- (m)  $e^{\sin x} = 1 + \frac{x}{1!} + \frac{x^2}{2!} - \frac{3x^4}{4!} - \frac{8x^5}{5!} - \frac{3x^6}{6!} + \frac{56x^7}{7!} + \dots$
- (n)  $e^{\tan x} = 1 + x + \frac{x^2}{2!} + \frac{3x^3}{3!} + \frac{9x^4}{4!} + \frac{37x^5}{5!} + \frac{177x^6}{6!} \dots$
- (o)  $(\text{ver } \sin^{-1} x)^2 = 2 \left( x + \frac{x^2}{3.2} + \frac{1.2}{3.5} \frac{x^3}{3} + \frac{1.2.3}{3.5.7} \frac{x^4}{4} + \dots \right)$
- (p)  $\frac{1}{2-2x+x^2} = \frac{2}{4} + \frac{2x}{4} + \frac{x^2}{4} - \frac{2x^4}{4^2} - \frac{2x^5}{4^2} - \frac{x^6}{4^2} \dots$
- (q)  $\frac{1+x}{6-5x+x^2} = \frac{3}{2-x} - \frac{4}{3-x} = \left(\frac{3}{2} - \frac{4}{3}\right) + \left(\frac{3}{2^2} - \frac{4}{3^2}\right)x + \dots$

*Answers.*

- (a)  $-a < x < a$ ;  
(b)  $-\infty < x < \infty$  if  $n > 0$ ,  $-\frac{\pi}{2} < x < \frac{\pi}{2}$  if  $n < 0$ ;  
(c)  $-\infty < x < \infty$ ; (d)  $-\infty < x < \infty$ ;  
(e)  $-1 < x < 1$ ; (f)  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ ;  
(g)  $-\frac{1}{3} < x < \frac{1}{3}$ ; (h)  $-1 < x < 1$ ;  
(i)  $-1 < x < 1$ ; (j)  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ ;  
(k)  $-\pi < x < \pi$ ; (l)  $-\frac{\pi}{4} < x < \frac{\pi}{4}$ ;  
(m)  $-\infty < x < \infty$ ; (n)  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ ;  
(o)  $-2 < x < 2$ ; (p)  $-\sqrt{2} < x < \sqrt{2}$ ;  
(q)  $-2 < x < 2$ .



## CHAPTER XVIII.

## KEY TO THE SOLUTION OF DIFFERENTIAL EQUATIONS.

222. In this chapter an analytical key leads to a set of concise, practical rules, embodying most of the ordinary methods employed in solving differential equations; and the attempt has been made to render these rules so explicit that they may be understood and applied by any one who has mastered the Integral Calculus proper.

The key is based upon "Boole's Differential Equations" (London: Macmillan & Co.), to which or to "Forsyth's Differential Equations" (London: Macmillan & Co.), we refer the student who wishes to become familiar with the theoretical considerations upon which the working rules are based.

223. A *differential equation* is an expressed relation involving derivatives with or without the primitive variables from which they are derived.

For example:

$$(1+x)y + (1-y)x \frac{dy}{dx} = 0, \quad (1)$$

$$x \frac{dy}{dx} - ay = x + 1, \quad (2)$$

$$x \frac{dy}{dx} - y + x\sqrt{x^2 - y^2} = 0, \quad (3)$$

$$\left(\frac{dy}{dx}\right)^3 = y^4 \left(y + x \frac{dy}{dx}\right), \quad (4)$$

$$\frac{d^4 y}{dx^4} - 2 \frac{d^3 y}{dx^3} + 2 \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = 1, \quad (5)$$

$$\sin^2 x \frac{d^2 y}{dx^2} + \sin x \cos x \frac{dy}{dx} - y = x - \sin x, \quad (6)$$

$$x(1-x)^2 \frac{d^2 y}{dx^2} - 2y = 0, \quad (7)$$

$$\frac{d^2 y}{dx^2} + \left(1 - \frac{2}{x^2}\right)y = 0, \quad (8)$$

$$D_x^2 z - a^2 D_y^2 z = 0, \quad (9)$$

are differential equations.

The *order* of a differential equation is the same as that of the derivative of highest order which appears in the equation.

Equations (1), (2), (3), and (4) are of the first order; (6), (7), (8), and (9) of the second order; and (5) of the fourth order.

The *degree* of a differential equation is the same as the power to which the derivative of highest order in the equation is raised, that derivative being supposed to enter into the equation in a rational form.

Equations (1), (2), (3), (5), (6), (7), (8), and (9) are all of the first degree; (4) is of the third degree.

A differential equation is *linear* when it would be of the first degree if the dependent variable and all its derivatives were regarded as unknown quantities.

Equations (2), (5), (6), (7), (8), and (9) are *linear*.

The equation not containing differentials or derivatives, and expressing the most general relation between the primitive variables consistent with the given differential equation, is called its *general solution* or *complete primitive*. A general solution will always contain arbitrary constants or arbitrary functions.

The differential equation is formed from the complete primitive by direct differentiation, or by differentiation and the subsequent elimination of constants or functions between the primitive and the derived equations.

If it has been formed by differentiation *only* without subsequent elimination or reduction, the differential equation is said to be *exact*.

A *singular solution* of a differential equation is a relation between the primitive variables which satisfies the differential equation by means of the values which it gives to the derivatives, but which cannot be obtained from the complete primitive by giving particular values to the arbitrary constants.

224. We shall illustrate the use of the key by solving equations (1), (2), (3), (4), (5), (6), (7), (8), and (9) of Art. 223 by its aid.

$$(1) \quad (1+x)y + (1-y)x \frac{dy}{dx} = 0, \text{ or } (1+x)ydx + (1-y)xdy = 0.$$

Beginning at the beginning of the key, we see that we have a single equation, and hence look under I., p. 310; it involves ordinary derivatives: we are then directed to II., p. 310; it contains two variables: we go to III., p. 310; it is of the first order, IV., p. 310. and of the first degree, V., p. 310.

It is reducible to the form

$$\frac{1+x}{x} dx + \frac{1-y}{y} dy = 0,$$

which comes under  $Xdx + Ydy = 0$ .

Hence we turn to (1), p. 314, and there find the specific directions for its solution. Integrating each term separately, we get

$$\log x + x + \log y - y = c, \text{ or } \log(xy) + x - y = c,$$

the required primitive equation.

$$(2) \quad x \frac{dy}{dx} - ay = x + 1.$$

Beginning again at the beginning of the key, we are directed through I., II., III., IV., to V., p. 310. Looking under V., we see that it will come under either the third or the fourth head. Let us try the fourth; we are referred to (4), p. 314, for specific directions.

Obeying instructions, the work is as follows:

$$x \frac{dy}{dx} - ay = 0,$$

$$\begin{aligned}
 xdy - aydx &= 0, \\
 \frac{dy}{y} - \frac{adx}{x} &= 0, \\
 \log y - a \log x &= c, \\
 \log \frac{y}{x^a} &= c; \\
 \frac{y}{x^a} &= C, \\
 y &= Cx^a, \\
 \frac{dy}{dx} &= aCx^{a-1} + x^a \frac{dC}{dx}.
 \end{aligned} \tag{1}$$

Substitute in the given equation,

$$\begin{aligned}
 aCx^a + x^{a+1} \frac{dC}{dx} - aCx^a &= x + 1, \\
 x^{a+1} \frac{dC}{dx} - (x + 1) &= 0, \\
 dC - \frac{x + 1}{x^{a+1}} dx &= 0, \\
 C + \frac{1}{(a - 1)x^{a-1}} + \frac{1}{ax^a} &= C'.
 \end{aligned}$$

Substitute this value for  $C$  in (1), and we get

$$y = C'x^a - \left( \frac{1}{a} + \frac{x}{a-1} \right),$$

the required primitive.

$$(3) \quad x \frac{dy}{dx} - y + x \sqrt{x^2 - y^2} = 0.$$

Beginning at the beginning of the key, we are directed through I., II., III., IV., to V., page 310. Looking under V., we find that our equation does not come under any of the special forms there given. We are consequently driven to obtaining a solution in the form of a series, and for specific instructions we are referred to (13), page 316. Obeying these, our process is the following:

$$\frac{dy}{dx} = \frac{y}{x} - \sqrt{x^2 - y^2},$$

$$\frac{dy_0}{dx_0} = \frac{y_0}{x_0} - \sqrt{x_0^2 - y_0^2},$$

$$\frac{d^2y}{dx^2} = -y - \frac{2}{x}\sqrt{x^2 - y^2},$$

$$\frac{d^2y_0}{dx_0^2} = -y_0 - \frac{2}{x_0}\sqrt{x_0^2 - y_0^2},$$

$$\frac{d^3y}{dx^3} = -\frac{3y}{x} + \sqrt{x^2 - y^2},$$

$$\frac{d^3y_0}{dx_0^3} = -\frac{3y_0}{x_0} + \sqrt{x_0^2 - y_0^2},$$

$$\frac{d^4y}{dx^4} = y + \frac{4}{x}\sqrt{x^2 - y^2},$$

$$\frac{d^4y_0}{dx_0^4} = y_0 + \frac{4}{x_0}\sqrt{x_0^2 - y_0^2},$$

$$\frac{d^5y}{dx^5} = \frac{5y}{x} - \sqrt{x^2 - y^2},$$

$$\frac{d^5y_0}{dx_0^5} = \frac{5y_0}{x_0} - \sqrt{x_0^2 - y_0^2},$$

$$\frac{d^6y}{dx^6} = -y - \frac{6}{x}\sqrt{x^2 - y^2},$$

$$\frac{d^6y_0}{dx_0^6} = -y_0 - \frac{6}{x_0}\sqrt{x_0^2 - y_0^2},$$

$$\frac{d^7y}{dx^7} = -\frac{7y}{x} + \sqrt{x^2 - y^2},$$

$$\frac{d^7y_0}{dx_0^7} = -\frac{7y_0}{x_0} + \sqrt{x_0^2 - y_0^2};$$

and the general value of  $y$  is

$$\begin{aligned} y &= y_0 + (x - x_0) \left( \frac{y_0}{x_0} - \sqrt{x_0^2 - y_0^2} \right) - \frac{(x - x_0)^2}{2!} \left( y_0 + \frac{2}{x_0} \sqrt{x_0^2 - y_0^2} \right) \\ &\quad - \frac{(x - x_0)^3}{3!} \left( \frac{3y_0}{x_0} - \sqrt{x_0^2 - y_0^2} \right) \\ &\quad + \frac{(x - x_0)^4}{4!} \left( y_0 + \frac{4}{x_0} \sqrt{x_0^2 - y_0^2} \right) + \frac{(x - x_0)^5}{5!} \left( \frac{5y_0}{x_0} - \sqrt{x_0^2 - y_0^2} \right) \\ &\quad - \frac{(x - x_0)^6}{6!} \left( y_0 + \frac{6}{x_0} \sqrt{x_0^2 - y_0^2} \right) \dots \end{aligned}$$

This result can be very greatly simplified by breaking up the series; we have

$$\begin{aligned} y &= y_0 \left( 1 - \frac{(x - x_0)^2}{2!} + \frac{(x - x_0)^4}{4!} - \frac{(x - x_0)^6}{6!} \dots \right) \\ &\quad + y_0 \frac{(x - x_0)}{x_0} \left( 1 - \frac{(x - x_0)^2}{2!} + \frac{(x - x_0)^4}{4!} - \frac{(x - x_0)^6}{6!} \dots \right) \\ &\quad - \sqrt{x_0^2 - y_0^2} \left( (x - x_0) - \frac{(x - x_0)^3}{3!} + \frac{(x - x_0)^5}{5!} - \frac{(x - x_0)^7}{7!} \dots \right) \end{aligned}$$

$$-\sqrt{x_0^2 - y_0^2} \left( \frac{x - x_0}{x_0} \right) \left( (x - x_0) - \frac{(x - x_0)^3}{3!} + \frac{(x - x_0)^5}{5!} - \frac{(x - x_0)^7}{7!} \dots \right),$$

or

$$y = x \left[ \frac{y_0}{x_0} \cos(x - x_0) - \frac{\sqrt{x_0^2 - y_0^2}}{x_0} \sin(x - x_0) \right]$$

$$= x \left[ \frac{y_0}{x_0} \cos(x - x_0) - \sqrt{1 - \frac{y_0^2}{x_0^2}} \cdot \sin(x - x_0) \right].$$

$\frac{y_0}{x_0}$  is entirely arbitrary; call it  $\sin a$ , then

$$y = x [\sin a \cos(x - x_0) - \cos a \sin(x - x_0)] = x \sin[a - (x - x_0)],$$

$y = x \sin(c - x)$ , where  $c$  is any constant.

$$(4) \quad \left( \frac{dy}{dx} \right)^2 = y^4 \left( y + x \frac{dy}{dx} \right).$$

Beginning at the beginning of the key, we are directed through I., II., III., IV., to VI., page 311. Looking under VI. we see that the equation is of the first degree in  $x$ ; we are referred to (17), page 318, for our specific instructions.

Obedying these, we first replace  $\frac{dy}{dx}$  by  $p$ ; the equation becomes

$$p^2 = y^4 (y + xp).$$

Differentiate relatively to  $y$ , and we get

$$3p^2 \frac{dp}{dy} = 4y^3 (y + xp) + 2y^4 + xy^4 \frac{dp}{dy}.$$

Eliminate  $x$ ,

$$3p^2 \frac{dp}{dy} = 4 \frac{p^2}{y} + 2y^4 + \left( p^2 - \frac{y^5}{p} \right) \frac{dp}{dy},$$

or

$$\frac{2p^2 + y^5}{p} \frac{dp}{dy} - 2 \frac{(2p^2 + y^5)}{y} = 0.$$

Striking out the factor  $2p^2 + y^2$ , we have

$$\frac{1}{p} \frac{dp}{dy} - \frac{2}{y} = 0,$$

a differential equation of the first order and degree in which the variables are separated, and which therefore can be solved by (1), page 314.

Its solution is  $\log p - \log y^2 = C$ ,

or  $\frac{p}{y^2} = c$ .

Eliminating  $p$  between this and the given equation, and reducing, we have  $cy(x - c^2) = 1$ , as our required solution.

$$(5) \quad \frac{d^4 y}{dx^4} - 2 \frac{d^3 y}{dx^3} + 2 \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = 1. \quad (1)$$

Beginning at the beginning of the key, we are directed through I., II., III., VII., to (22) (a), page 319, for our specific directions.

We see at once that  $y = 1$  is a particular solution.

Obedient directions, we have now to solve

$$\frac{d^4 y}{dx^4} - 2 \frac{d^3 y}{dx^3} + 2 \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = 0 \quad \text{by (21)}. \quad (2)$$

Let  $y = e^{mx}$ , and we have

$$m^4 - 2m^3 + 2m^2 - 2m + 1 = 0,$$

as our auxiliary algebraic equation in  $m$ . Its roots are

$$1, 1, \sqrt{-1}, -\sqrt{-1}.$$

The solution of (2) is then

$$y = (A + Bx)e^x + C \cos x + D \sin x,$$

and of (1) is

$$y = (A + Bx)e^x + C \cos x + D \sin x + 1.$$

$$(6) \quad \sin^2 x \frac{d^2 y}{dx^2} + \sin x \cos x \frac{dy}{dx} - y = x - \sin x. \quad (1)$$

Beginning at the beginning of the key, we are directed through I., II., III., VII., to (24), page 321, for our specific instructions.

Dividing through by  $\sin^2 x$ , the equation becomes

$$\frac{d^2 y}{dx^2} + \operatorname{ctn} x \frac{dy}{dx} - \operatorname{csc}^2 x \cdot y = x \operatorname{csc}^2 x - \operatorname{csc} x. \quad (2)$$

$y = \operatorname{ctn} x$  is found by inspection to be a solution of

$$\frac{d^2 y}{dx^2} + \operatorname{ctn} x \frac{dy}{dx} - \operatorname{csc}^2 x \cdot y = 0;$$

(2) can then be solved by (24) (a).

Substitute  $y = z \operatorname{ctn} x$  in (2), and it becomes

$$\operatorname{ctn} x \frac{d^2 z}{dx^2} + (\operatorname{ctn}^2 x - 2 \operatorname{csc}^2 x) \frac{dz}{dx} = x \operatorname{csc}^2 x - \operatorname{csc} x,$$

$$\text{or} \quad \frac{d^2 z}{dx^2} - (\tan x + \sec x \operatorname{csc} x) \frac{dz}{dx} = x \sec x \operatorname{csc} x - \sec x. \quad (3)$$

Referring to (25), page 323, and obeying instructions, we let  $z' = \frac{dz}{dx}$ , and (3) becomes

$$\frac{dz'}{dx} - (\tan x + \sec x \operatorname{csc} x) z' = x \sec x \operatorname{csc} x - \sec x,$$

a linear differential equation of the first order in  $z'$ , whose solution by (4), page 314, is

$$z' = A \tan x \sec x - x \sec^2 x + \tan x \sec x (\log \tan \frac{x}{2} - \log \sin x);$$

but  $z' = \frac{dz}{dx}$ , whence integrating, we have

$$z = B + A \sec x - x \tan x - (1 + \sec x) \log (1 + \cos x),$$

and

$$y = A \operatorname{csc} x + B \operatorname{ctn} x - x - (\operatorname{csc} x + \operatorname{ctn} x) \log (1 + \cos x).$$



$$(7) \quad x(1-x)^2 \frac{d^2 y}{dx^2} - 2y = 0.$$

Beginning at the beginning of the key, we are directed through I., II., III., VII., to (24), page 321, for our specific instructions.

Let us try the method of (24) (e), page 323.

Assume  $y = \sum a_n x^n$ , and substitute in the given equation; we have

$$\begin{aligned} \Sigma [m(m-1)a_n x^{n-1} - 2m(m-1)a_n x^n \\ + m(m-1)a_n x^{n+1} - 2a_n x^n] = 0. \end{aligned}$$

Writing the coefficient of  $x^n$  in this sum equal to zero, we have

$$m(m+1)a_{n-1} - 2[m(m-1)+1]a_n + (m-1)(m-2)a_{n-1} = 0,$$

and we wish to choose the simplest set of values that will satisfy this relation.

Substituting  $m=0$ ,  $m=-1$ ,  $m=-2$ , etc., in this relation, we find

$$a_{-1} = a_0, \quad a_{-2} = a_{-1}, \quad a_{-3} = a_{-2}, \quad \dots$$

Hence if we take  $a_0 = 0$ , it follows that

$$a_{-1} = a_{-2} = a_{-3} \dots = 0,$$

and no negative powers of  $x$  will occur in our particular solution.

Substituting now  $m=1$ ,  $m=2$ ,  $m=3$ , etc., we have

$$a_1 = a_2 = a_3 = a_4 = \dots$$

Taking  $a_1 = 1$ , we get as our required particular solution of the given equation

$$y = x + x^2 + x^3 + x^4 + \dots$$

This can be written in finite form, since we know that

$$1 + x + x^2 + x^3 \dots = \frac{1}{1-x}.$$

Hence 
$$y = \frac{x}{1-x}$$

is a particular solution.

Turning now to (24) (a), page 321, we find

$$y = \frac{cx}{1-x} + c' \left( 1 + x + \frac{2x}{1-x} \log x \right).$$

$$(8) \quad \frac{d^2 y}{dx^2} + \left( 1 - \frac{2}{x^2} \right) y = 0.$$

Beginning at the beginning of the key, we are directed through I., II., III., VII., to (24), page 321, for our specific instructions. Let us try again the method (24) (e), page 323.

Assume  $y = \sum a_m x^m$ , and substitute in the given equation,

$$\sum [m(m-1)a_m x^{m-2} + a_m x^m - 2a_m x^{m-2}] = 0.$$

The terms containing  $x^m$  are

$$(m+2)(m+1)a_{m+2}x^m + a_m x^m - 2a_{m+2}x^m;$$

writing the sum of the coefficients equal to zero, we have

$$m(m+3)a_{m+2} + a_m = 0. \tag{1}$$

Letting  $m = 0$  and  $m = -3$ , we get  $a_0 = 0$  and  $a_{-3} = 0$ ; and all terms of  $y$  involving even negative powers of  $x$  disappear, as do all terms involving odd negative powers, except the  $-1$ st.

In general 
$$a_{m+2} = -\frac{a_m}{m(m+3)}. \tag{2}$$

From this we get

$$a_4 = -\frac{a_2}{2 \cdot 5} = -\frac{1}{3!5}, \text{ if we take } a_2 = \frac{1}{3},$$

$$a_6 = \frac{a_2}{2 \cdot 4 \cdot 5 \cdot 7} = \frac{1}{5!7},$$

$$a_8 = -\frac{a_2}{2 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 9} = -\frac{1}{7!9},$$

$$a_{10} = \frac{a_2}{2 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9 \cdot 11} = \frac{1}{9!11}.$$

Hence 
$$y = \frac{x^2}{3} - \frac{x^4}{3!5} + \frac{x^6}{5!7} - \frac{x^8}{7!9} + \frac{x^{10}}{9!11} \dots$$

is a particular solution of the given equation. This can be thrown into finite form without much labor.

We have 
$$xy = \frac{x^3}{3} - \frac{x^5}{3!5} + \frac{x^7}{5!7} - \frac{x^9}{7!9} + \frac{x^{11}}{9!11} \dots,$$

$$\begin{aligned} \frac{d(xy)}{dx} &= x^2 - \frac{x^4}{3!} + \frac{x^6}{5!} - \frac{x^8}{7!} + \frac{x^{10}}{9!} \dots, \\ &= x \left( x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} \dots \right), \\ &= x \sin x; \end{aligned}$$

whence  $xy = \sin x - x \cos x,$

and  $y = \frac{1}{x} (\sin x - x \cos x).$

By going back to (2), and using odd values of  $m$ , we get another solution of our given equation, namely,

$$y = \frac{1}{x} + \frac{x}{2} - \frac{x^3}{2!4} + \frac{x^5}{4!6} - \frac{x^7}{6!8},$$

which can be reduced to

$$y = \frac{1}{x} (\cos x + x \sin x).$$

Hence our complete solution is

$$y = \frac{1}{x} [A (\cos x + x \sin x) + B (\sin x - x \cos x)],$$

or 
$$y = A' \left[ \frac{\cos(x-c)}{x} + \sin(x-c) \right],$$

if we let  $\frac{B}{A} = \tan c.$

$$(9) \quad D_x^2 z - a^2 D_x^{-2} z = 0.$$

Beginning at the beginning of the key, we are directed through I. and IX. to (45), p. 331, for our specific instructions.

Obeying these, our work is as follows :

$$dy^2 - a^2 dx^2 = 0,$$

$$dy - adx = 0, \tag{1}$$

$$dy + adx = 0, \tag{2}$$

$$dpdy - a^2 dqdx = 0. \tag{3}$$

Combining (1) and (3), we get

$$dpdy - adqdy = 0,$$

or  $dp - adq = 0. \tag{4}$

(1) gives  $y - ax = a.$

(4) gives  $p - aq = \beta.$

(2) and (3) give us, in the same way,

$$y + ax = a_1,$$

$$p + aq = \beta_1;$$

and our two first integrals are

$$p - aq = f_1(y - ax), \tag{5}$$

$$p + aq = f_2(y + ax), \tag{6}$$

$f_1$  and  $f_2$  denoting arbitrary functions.

Determining  $p$  and  $q$ , from (5) and (6),

$$p = \frac{1}{2} [f_2(y + ax) + f_1(y - ax)],$$

$$q = \frac{1}{2a} [f_2(y + ax) - f_1(y - ax)];$$

$$dz = \frac{1}{2} [f_2(y + ax) + f_1(y - ax)] dx + \frac{1}{2a} [f_2(y + ax) - f_1(y - ax)] dy$$

$$= \frac{f_2(y + ax)(dy + adx) - f_1(y - ax)(dy - adx)}{2a}.$$

Hence,  $z = F(y + ax) + F_1(y - ax),$

where  $F$  and  $F_1$  denote arbitrary functions obtained by integrating  $f_1$  and  $f_2$ , which are arbitrary.

225. When a differential equation does not come under any of the forms given in the key, a change of dependent or independent variable, or of both, will often reduce it to one of the standard forms. No general rule can be laid down for such a substitution. It will, however, often suffice to introduce a new letter for the sum, or the difference, or the product, or the quotient of the variables, or for a power of one or of both. Sometimes an ingenious trigonometric substitution is effective, or a change from rectangular to polar coördinates; that is, the introduction of  $r \cos \phi$  for  $x$  and  $r \sin \phi$  for  $y$ .

The following examples of such substitutions are instructive.

(A.) *Change of dependent variable.*

$$(1) (x+y)^2 \frac{dy}{dx} = a^2, \text{ reduces to } \frac{z^2}{a^2+z^2} dz - dx = 0,$$

if we introduce  $z = x + y$ .

$$(2) \frac{d\theta}{d\phi} = \sin(\phi - \theta), \text{ reduces to } \frac{d\omega}{1 - \sin \omega} - d\phi = 0,$$

if  $\omega = \phi - \theta$ .

$$(3) (x - y^2) dx + 2xy dy = 0, \text{ reduces to } (x - z) dx + x dz = 0,$$

if  $z = y^2$ .

$$(4) x \frac{dy}{dx} - y + x \sqrt{x^2 - y^2} = 0, \text{ reduces to } \frac{dz}{\sqrt{1 - z^2}} + dx = 0,$$

if  $z = \frac{y}{x}$ .

$$(5) \frac{d^2 y}{dx^2} + \frac{2}{x} \frac{dy}{dx} - n^2 y = 0, \text{ reduces to } \frac{d^2 z}{dx^2} - n^2 z = 0,$$

if  $z = xy$ .

(B.) *Change of independent variable.*

$$(1) (1 - x^2)^2 \frac{d^2 y}{dx^2} + y = 0, \text{ reduces to}$$

$$\cos^2 \theta \frac{d^2 y}{d\theta^2} + \sin \theta \cos \theta \frac{dy}{d\theta} + y = 0, \text{ if } x = \sin \theta.$$

(2)  $\frac{d^2 y}{dx^2} + \tan x \frac{dy}{dx} + \cos^2 x \cdot y = 0$ , reduces to  $\frac{d^2 y}{dz^2} + y = 0$ ,

if  $z = \sin x$ .

(C.) *Change of both variables.*

(1)  $\left(1 - \frac{dy^2}{dx^2}\right)xy = \frac{dy}{dx}(x^2 - y^2 - a^2)$ , reduces to

$v - z \frac{dv^2}{dz^2} - \frac{dv}{dz}(z - v - a^2) = 0$ , if  $z = x^2$  and  $v = y^2$ .

(2)  $(y - x)(1 + x^2)^{\frac{1}{2}} \frac{dy}{dx} = (1 + y^2)^{\frac{3}{2}}$ , reduces to

$\sin(\phi - \theta) d\phi = d\theta$ , if  $x = \tan \theta$  and  $y = \tan \phi$ .

(3)  $\left(x \frac{dy}{dx} - y\right)^2 = a \left(1 + \frac{dy^2}{dx^2}\right)(x^2 + y^2)^{\frac{3}{2}}$ , reduces to

$\frac{dr}{\sqrt{r(1-ar)}} - \frac{d\phi}{\sqrt{a}} = 0$ , if  $x = r \cos \phi$  and  $y = r \sin \phi$ .

## KEY.

	Page
Single equation . . . . .	I. 310
System of simultaneous equations . . . . .	VIII. 313
<b>I. Involving ordinary derivatives . . . . .</b>	<b>II. 310</b>
Involving partial derivatives . . . . .	IX. 313
<b>II. Containing two variables . . . . .</b>	<b>III. 310</b>
Containing three variables and of first degree.	
General form, $Pdx + Qdy + Rdz = 0$ . . . . .	(36) 327
Containing more than three variables and of the first degree. General form, $Pdx_1 + Qdx_2$ $+ Rdx_3 + \dots = 0$ . . . . .	(37) 328
<b>III. Of first order . . . . .</b>	<b>IV. 310</b>
Not of first order . . . . .	VII. 312
<b>IV. Of first degree. General form, <math>Mdx + Ndy = 0</math> . . . . .</b>	<b>V. 310</b>
Not of first degree . . . . .	VI. 311
<b>V Of first degree. General form, <math>Mdx + Ndy = 0</math>.</b>	
Variables separated or separable; that is, of or reducible to the form $Xdx + Ydy = 0$ , where $X$ is a function of $x$ alone, and $Y$ is a function of $y$ alone* . . . . .	(1) 314
$M$ and $N$ homogeneous functions of $x$ and $y$ of the same degree . . . . .	(2) 314

\*Of course,  $X$  and  $Y$  may be constants.

	Page
Of the form $(ax + by + c) dx + (a'x + b'y + c') dy = 0$ . . . . .	(3) 314
Linear. General form, $\frac{dy}{dx} + X_1y = X_2$ , where $X_1$ and $X_2$ are functions of $x$ alone* . . . . .	(4) 314
Of the form $\frac{dy}{dx} + X_1y = X_2y^n$ , where $X_1$ and $X_2$ are functions of $x$ alone* . . . . .	(5) 315
$Mdx + Ndy$ an exact differential. Test, $D_yM = D_xN$ . . . . .	(6) 315
$Mx + Ny = 0$ . . . . .	(7) 315
$Mx - Ny = 0$ . . . . .	(8) 315
Of the form $F_1(xy) ydx + F_2(xy) xdy = 0$ . . . . .	(9) 315
$\frac{D_yM - D_xN}{N}$ , a function of $x$ alone . . . . .	(10) 315
$\frac{D_xN - D_yM}{M}$ , a function of $y$ alone . . . . .	(11) 316
$\frac{D_yM - D_xN}{Ny - Mx}$ , a function of $(xy)$ . . . . .	(12) 316
A solution in the form of a series can always be obtained . . . . .	(13) 316

VI. Not of first degree.

Can be solved as an algebraic equation in $p$ , where $p$ stands for $\frac{dy}{dx}$ . . . . .	(14) 317
Involves only one of the variables and $p$ , where $p$ stands for $\frac{dy}{dx}$ . . . . .	(15) 317
Of the first degree in $x$ and $y$ ; that is, of the form $xf_1p + yf_2p = f_3p$ , where $p$ stands for $\frac{dy}{dx}$ . . . . .	(16) 317
Of the first degree in $x$ or $y$ . . . . .	(17) 318
Homogeneous relatively to $x$ and $y$ . . . . .	(18) 318

\* Of course,  $X_1$  and  $X_2$  may be constants.



	Page
Of the form $F(\phi, \psi) = 0$ , where $\phi$ and $\psi$ are functions of $x, y$ , and $\frac{dy}{dx}$ , such that $\phi = a$ $\psi = b$ , will lead, on differentiation, to the same differential equation of the second order . . . . .	(19) 318
A singular solution will answer . . . . .	(20) 318

VII. Not of first order.

Linear, with constant coefficients; second member zero* . . . . .	(21) 319
Linear, with constant coefficients; second member not zero* . . . . .	(22) 319
Of the form $(u + bx)^n \frac{d^n y}{dx^n} + A(u + bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + Ly = X$ , where $X$ is a function of $x$ alone † . . . . .	(23) 321
Linear; of second order; coefficients not constant. General form, $\frac{d^2 y}{dx^2} + P \frac{dy}{dx} + Qy = R$ ; $P, Q$ , and $R$ being functions of $x$ . . . . .	(24) 321
Either of the primitive variables wanting . . . . .	(25) 323
Of the form $\frac{d^n y}{dx^n} = X$ , $X$ being a function of $x$ alone † . . . . .	(26) 323
Of the form $\frac{d^2 y}{dx^2} = Y$ , $Y$ being a function of $y$ alone † . . . . .	(27) 324
Of the form $\frac{d^n y}{dx^n} = f \frac{d^{n-1} y}{dx^{n-1}}$ . . . . .	(28) 324
Of the form $\frac{d^n y}{dx^n} = f \frac{d^{n-2} y}{dx^{n-2}}$ . . . . .	(29) 324
Homogeneous on the supposition that $x$ and	

\* The first member is supposed to contain only those terms involving the dependent variable or its derivatives.

† See note, p. 310.

	Page
$y$ are of the degree 1, $\frac{dy}{dx}$ of the degree 0, $\frac{d^2y}{dx^2}$ of the degree $-1$ , ... . . . . . (30)	325
Homogeneous on the supposition that $x$ is of the degree 1, $y$ of the degree $n$ , $\frac{dy}{dx}$ of the degree $n - 1$ , $\frac{d^2y}{dx^2}$ of the degree $n - 2$ , ... (31)	325
Homogeneous relatively to $y$ , $\frac{dy}{dx}$ , $\frac{d^2y}{dx^2}$ , ... . (32)	325
Containing the first power only of the deriva- tive of the highest order . . . . . (33)	325
Of the form $\frac{d^2y}{dx^2} + X\frac{dy}{dx} + Y\left[\frac{dy}{dx}\right]^2 = 0$ , where $X$ is a function of $x$ alone and $Y$ a func- tion of $y$ alone * . . . . . (34)	326
Singular integral will answer . . . . . (35)	326
VIII. Simultaneous equations of the first order . . (38)	328
Not of the first order . . . . . (39)	329
IX. All the partial derivatives taken with respect to one of the independent variables . . . (40)	329
Of the first order and Linear . . . . . X.	313
Of the first order and not Linear . . . . . XI.	313
Of the second order and containing the deriva- tives of the second order only in the first degree. General form $RD_x^2z + SD_xD_yz +$ $TD_y^2z = V$ , where $R, S, T$ , and $V$ may be functions of $x, y, z, D_xz$ , and $D_yz$ . . . (45)	331
X. Containing three variables . . . . . (41)	330
Containing more than three variables . . . (42)	330
XI. Containing three variables . . . . . (43)	330
Containing more than three variables . . . (44)	331

\* See note, p. 310.

- (1) Of or reducible to the form  $Xdx + Ydy = 0$ , where  $X$  is a function of  $x$  alone and  $Y$  is a function of  $y$  alone.

Integrate each term separately, and write the sum of their integrals equal to an arbitrary constant.

- (2)  $M$  and  $N$  homogeneous functions of  $x$  and  $y$  of the same degree.

Introduce in place of  $y$  the new variable  $v$  defined by the equation  $y = vx$ , and the equation thus obtained can be solved by (1).

Or, multiply the equation through by  $\frac{1}{Mx + Ny}$ , and its first member will become an exact differential, and the solution may be obtained by (6).

- (3) Of the form  $(ax + by + c)dx + (a'x + b'y + c')dy = 0$ .

If  $ab' - a'b = 0$ , the equation may be thrown into the form  $(ax + by + c)dx + \frac{a'}{a}(ax + by + c)dy = 0$ . If now  $z = ax + by$  be introduced in place of either  $x$  or  $y$ , the resulting equation can be solved by (1).

If  $ab' - a'b$  does not equal zero, the equation can be made homogeneous by assuming  $x = x' - \alpha$ ,  $y = y' - \beta$ , and determining  $\alpha$  and  $\beta$  so that the constant terms in the new values of  $M$  and  $N$  shall disappear, and it can then be solved by (2).

- (4) Linear. General form  $\frac{dy}{dx} + X_1y = X_2$ , where  $X_1$  and  $X_2$  are functions of  $x$  alone.

Solve on the supposition that  $X_2 = 0$  by (1); and from this solution obtain a value for  $y$ , involving of course an arbitrary constant  $C$ . Substitute this value of  $y$  in the given equation, regarding  $C$  as a variable, and there will result a differential equation, involving  $C$  and  $x$ , whose solution by (1) will express  $C$  as a function of  $x$ . Substitute this value for  $C$  in the expression already obtained for  $y$ , and the result will be the required solution.

- (5) Of the form  $\frac{dy}{dx} + X_1y = X_2y^n$ , where  $X_1$  and  $X_2$  are functions of  $x$  alone.

Divide through by  $y^n$ , and then introduce  $z = y^{1-n}$  in place of  $y$ , and the equation will become linear and may be solved by (4).

- (6)  $Mdx + Ndy$  an exact differential. Test  $D_yM = D_xN$ .

Find  $\int Mdx$ , regarding  $y$  as constant, and add an arbitrary function of  $y$ . Determine this function of  $y$  by the fact that the differential of the result just mentioned, taken on the supposition that  $x$  is constant, must equal  $Ndy$ . Write equal to an arbitrary constant the  $\int Mdx$  above mentioned plus the function of  $y$  just determined.

- (7)  $Mx + Ny = 0$ .

Divide the first term of  $Mdx + Ndy = 0$  by  $Mx$ , and the second by its equal  $-Ny$ , and integrate by (1).

- (8)  $Mx - Ny = 0$ .

Divide the first term of  $Mdx + Ndy = 0$  by  $Mx$ , and the second by its equal  $Ny$ , and integrate by (1).

- (9) Of the form  $f_1(xy)ydx + f_2(xy)xdy = 0$ .

Multiply through by  $\frac{1}{Mx - Ny}$ , and the first member will become an exact differential. The solution may then be found by (6).

- (10)  $\frac{D_yM - D_xN}{N}$ , a function of  $x$  alone.

Multiply the equation through by  $e^{\int \frac{D_yM - D_xN}{N} dx}$ , and the first member will become an exact differential. The solution may then be found by (6).

- (11)  $\frac{D_x N - D_y M}{M}$ , a function of  $y$  alone.

Multiply the equation through by  $e^{\int \frac{D_x N - D_y M}{M} \cdot dy}$ , and the first member will become an exact differential. The solution may then be found by (6).

- (12)  $\frac{D_y M - D_x N}{Ny - Mx}$ , a function of  $(xy)$ .

Multiply the equation through by  $e^{\int \frac{D_y M - D_x N}{Ny - Mx} \cdot dx}$  where  $v = xy$ , and the first member will become an exact differential. The solution may thus be found by (6).

- (13) A solution of  $Mdx + Ndy = 0$  in the form of a series can always be obtained.

Throw the given equation into the form  $\frac{dy}{dx} = -\frac{M}{N}$ , then differentiate, and in the result replace  $\frac{dy}{dx}$  by  $-\frac{M}{N}$ , thus obtaining a value of  $\frac{d^2 y}{dx^2}$  in terms of  $x$  and  $y$ ; by successive differentiations and substitutions get values of  $\frac{d^3 y}{dx^3}$ ,  $\frac{d^4 y}{dx^4}$ , etc., in terms of  $x$  and  $y$ .

If  $y_0$  is the value of  $y$  corresponding to any chosen value  $x_0$  of  $x$ ,  $y$  can now be developed by Taylor's Theorem.

$$\text{We have} \quad y = fx = f(x_0 + x - x_0)$$

$$= fx_0 + (x - x_0)f'x_0 + \frac{(x - x_0)^2}{2!}f''x_0 + \frac{(x - x_0)^3}{3!}f'''x_0 + \dots,$$

or

$$y = y_0 + (x - x_0)\frac{dy_0}{dx_0} + \frac{(x - x_0)^2}{2!}\frac{d^2 y_0}{dx_0^2} + \frac{(x - x_0)^3}{3!}\frac{d^3 y_0}{dx_0^3} + \dots,$$

where  $\frac{dy_0}{dx_0}$ ,  $\frac{d^2 y_0}{dx_0^2}$ ,  $\frac{d^3 y_0}{dx_0^3}$ , etc.,

are obtained by replacing  $x$  and  $y$  by  $x_0$  and  $y_0$  in the values of

$$\frac{dy}{dx}, \frac{d^2 y}{dx^2}, \frac{d^3 y}{dx^3}, \text{ etc.,}$$

described above.

In the general case  $y_0$  is entirely arbitrary, and if the given equation is at all complicated, the solution is apt to be too complicated to be of much service. If, however, in a special problem the value of  $y$  corresponding to some value of  $x$  is given, and these values are taken as  $y_0$  and  $x_0$ , the solution will generally be useful.

- (14) Can be solved as an algebraic equation in  $p$ , where  $p$  stands for  $\frac{dy}{dx}$ .

Solve as an algebraic equation in  $p$ , and, after transposing all the terms to the first member, express the first member as the product of factors of the first order and degree. Write each of these factors separately equal to zero, and find its solution in the form  $V - c = 0$  by (V.). Write the product of the first members of these solutions equal to zero, using the same arbitrary constant in each.

- (15) Involves only one of the variables and  $p$ , where  $p$  stands for  $\frac{dy}{dx}$ .

By algebraic solution express the variable as an explicit function of  $p$ , and then differentiate through relatively to the other variable, regarding  $p$  as a new variable and remembering that  $\frac{dx}{dy} = \frac{1}{p}$ . There will result a differential equation of the first order and degree between the second variable and  $p$  which can be solved by (1). Eliminate  $p$  between this solution and the given equation, and the resulting equation will be the required solution.

- (16) Of the form  $xf_1p + yf_2p = f_3p$ , where  $p$  stands for  $\frac{dy}{dx}$ .

Differentiate the equation relatively to one of the variables, regarding  $p$  as a new variable, and, with the aid of the given equation, eliminate the original variable. There will result a linear differential equation of the first

order between  $p$  and the remaining variable, which may be simplified by striking out any factor not containing  $\frac{dp}{dx}$  or  $\frac{dy}{dx}$ , and can be solved by (4). Eliminate  $p$  between this solution and the given equation, and the result will be the required solution.

- (17) Of the first degree in  $x$  or  $y$ .

The equation can sometimes be solved by the method of (16), differentiating relatively to the variable which does not enter to the first degree.

- (18) Homogeneous relatively to  $x$  and  $y$ .

Let  $y = vx$ , and solve algebraically relatively to  $p$  or  $v$ ,  $p$  standing for  $\frac{dy}{dx}$ . The result will be of the form  $p = fv$ , or  $v = Fp$ . If

$$p = fv, \quad \frac{dy}{dx} = fv, \quad \frac{d(vx)}{dx} = fv, \quad x \frac{dv}{dx} + v = fv,$$

an equation that can be solved by (1). If

$$v = Fp, \quad \frac{y}{x} = Fp, \quad y = xFp,$$

an equation that can be solved by (16).

- (19) Of the form  $F(\phi, \psi) = 0$ , where  $\phi$  and  $\psi$  are functions of  $x$ ,  $y$ , and  $\frac{dy}{dx}$ , such that  $\phi = a$  and  $\psi = b$  will lead, on differentiation, to the same differential equations of the second order.

Eliminate  $\frac{dy}{dx}$  between  $\phi = a$  and  $\psi = b$ , where  $a$  and  $b$  are arbitrary constants subject to the relation that  $F(a, b) = 0$ , and the result will be the required solution.

- (20) Singular solution will answer.

Let  $\frac{dy}{dx} = p$ , and express  $p$  as an explicit function of  $x$  and  $y$ . Take  $\frac{dp}{dy}$ , regarding  $x$  as constant, and see

whether it can be made infinite by writing equal to zero any expression involving  $y$ . If so, and if the equation thus formed will satisfy the given differential equation, it is a singular solution.

Or take  $\frac{d\left(\frac{1}{p}\right)}{dx}$ , regarding  $y$  as constant, and see whether

it can be made infinite by writing equal to zero any expression involving  $x$ . If so, and if the equation thus formed is consistent with the given equation, it is a singular solution.

- (21) Linear, with constant coefficients. Second member zero.

Assume  $y = e^{mx}$ ;  $m$  being constant, substitute in the given equation, and then divide through by  $e^{mx}$ . There will result an algebraic equation in  $m$ . Solve this equation, and the complete value of  $y$  will consist of a series of terms characterized as follows: For every distinct real value of  $m$  there will be a term  $Ce^{mx}$ ; for each pair of imaginary values,  $a + b\sqrt{-1}$ ,  $a - b\sqrt{-1}$ , a term  $Ae^{ax} \cos bx + Be^{ax} \sin bx$ ; each of the coefficients  $A$ ,  $B$ , and  $C$  being an arbitrary constant, if the root or pair of roots occurs but once; and an algebraic polynomial in  $x$  of the  $(r-1)$ st degree with arbitrary constant coefficients, if the root or pair of roots occurs  $r$  times.

- (22) Linear, with constant coefficients. Second member not zero.

(a) If a particular solution of the given equation can be obtained by inspection, this value plus the value of  $y$  obtained by (21) on the hypothesis that the second member is zero, will be the complete value of the dependent variable.



(b) If the second member of the given equation can be got rid of by differentiation, or by differentiation and elimination between the given and the derived equations, solve the new differential equation thus obtained, by (21), and determine the superfluous arbitrary constants so that the given equation shall be satisfied.

In determining these superfluous constants, it will generally save labor to solve the original equation on the hypothesis that its second member is zero, and then to strike out from the preceding solution the terms which are duplicates of the ones in the second solution before proceeding to differentiate, as from the nature of the case they would drop out in the course of the work.

(c) If the given equation is of the second order, solve on the hypothesis that the second member is zero, by (21), obtain from this solution a simple particular solution by letting one of the arbitrary constants equal zero and the other equal unity, and let  $y = vz$  be this last solution; then substitute  $vz$  for  $y$  in the given equation; there will result a differential equation of the second order between  $x$  and  $z$  in which the dependent variable  $z$  will be wanting, and which can be completely solved by (25). Substitute the value of  $z$  thus obtained in  $y = vz$  and there will result the required solution of the given equation.

(d) Solve, on the hypothesis that the second member is zero, and obtain the complete value of  $y$  by (21). Denoting the order of the given equation by  $n$ , form the  $n - 1$  successive derivatives  $\frac{dy}{dx}, \frac{d^2y}{dx^2} \dots \frac{d^{n-1}y}{dx^{n-1}}$ . Then differentiate  $y$  and each of the values just obtained, regarding the arbitrary constants as new variables, and substitute the resulting values in the given equation; and by its aid, and that of the  $n - 1$  equations of condition formed by writing each of the derivatives of the second set,

except the  $n$ th, equal to the derivative of the same order in the first set, determine the arbitrary coefficients and substitute their values in the original expression for  $y$ .

(23) Of the form

$$(a + bx)^n \frac{d^n y}{dx^n} + A(a + bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + Ly = X,$$

where  $X$  is a function of  $x$  alone.

Assume  $a + bx = e^t$ , and change the independent variable in the given equation so as to introduce  $t$  in place of  $x$ . The solution can then be obtained by (22).

(24) Linear; of second order; coefficients not constants.

$$\text{General form } \frac{d^2 y}{dx^2} + P \frac{dy}{dx} + Qy = R.$$

(a) If a particular solution  $y = v$  of the equation

$$\frac{d^2 y}{dx^2} + P \frac{dy}{dx} + Qy = 0$$

can be found by inspection or other means, substitute  $y = vz$  in the given equation, which will then reduce to the form

$$v \frac{d^2 z}{dx^2} + \left( 2 \frac{dv}{dx} + Pv \right) \frac{dz}{dx} = R,$$

and can be solved by (25). Substitute the value of  $z$  thus found in  $y = vz$ , and the result will be the general solution of the given equation.

(b) The substitution of  $y = vz$  in the given equation, where  $v$  is given by the auxiliary differential equation

$$2 \frac{dv}{dx} + Pv = 0,$$

and can be found by (1), and should be used in the simplest possible form, will lead to a differential equation in  $z$  of the form

$$\frac{d^2 z}{dx^2} + Iz = R,$$

which is often simpler than the original equation.

(c) The introduction of  $z$  in place of the independent variable  $x$ ,  $z$  being a solution of the auxiliary differential equation

$$\frac{d^2 z}{dx^2} + P \frac{dz}{dx} = 0,$$

the simpler the better, will reduce the given equation to the form

$$\frac{d^2 y}{dz^2} + Iy = S,$$

which is often simpler than the original equation.

(d) If the first member of the given equation regarded as an operation performed on  $y$  can be resolved into the product of two operations, the equation can always be solved. The conditions of such a resolution are the following: let the given equation be

$$u \frac{d^2 y}{dx^2} + v \frac{dy}{dx} + wy = R,$$

where  $u$ ,  $v$ ,  $w$ , and  $R$  are functions of  $x$ ; this can be resolved into

$$\left( p \frac{d}{dx} + q \right) \left( r \frac{d}{dx} + s \right) y = R,$$

where  $p$ ,  $q$ ,  $r$ , and  $s$  are functions of  $x$ , if

$$pr = u, \quad qr + p \left( \frac{dr}{dx} + s \right) = v, \quad \text{and} \quad qs + p \frac{ds}{dx} = w;$$

and the values of  $p$ ,  $q$ ,  $r$ , and  $s$  can usually be obtained

by inspection. We have first to solve  $p \frac{dz}{dx} + qz = R$  by (4), and then to solve  $r \frac{dy}{dx} + sy = z$  by (4).

(e) A particular solution of the equation

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Qy = 0,$$

can often be obtained by assuming that  $y$  is of the form  $\Sigma a_m x^m$ ,  $m$  being an integer, substituting this value for  $y$  in the given equation, writing the sum of the coefficients of  $x^m$  equal to zero, since the equation must be identically true, and thus obtaining a relation between successive coefficients of the assumed series. The simplest set of values consistent with this relation should be substituted in the assumed value of  $y$ , which will then be a particular solution of the equation. If this solution can be expressed in finite form, the complete solution of the given equation can be obtained from it by the method described in (24) (a). If, however, two different particular solutions can be found by the method just described, each of them should be multiplied by an arbitrary constant, and the sum of these products will be the complete solution of the given equation.

(25) Either of the primitive variables wanting.

Assume  $z$  equal to the derivative of lowest order in the equation, and express the equation in terms of  $z$  and its derivatives with respect to the primitive variable actually present, and the order of the resulting equation will be lower than that of the given one.

(26) Of the form  $\frac{d^n y}{dx^n} = X$ .  $X$  being a function of  $x$  alone.

Solve by integrating  $n$  times successively with regard to  $x$ .

Or solve by (22).

(27) Of the form  $\frac{d^2 y}{dx^2} = Y$ .  $Y$  being a function of  $y$  alone.

Multiply by  $2 \frac{dy}{dx}$  and integrate relatively to  $x$ . There will result the equation  $\left(\frac{dy}{dx}\right)^2 = 2 \int Y dy + C$ , whence  $\frac{dy}{dx} = \left(2 \int Y dy + C\right)^{\frac{1}{2}}$ , an equation that may be solved by (1).

(28) Of the form  $\frac{d^n y}{dx^n} = f \frac{d^{n-1} y}{dx^{n-1}}$

Assume

$$\frac{d^{n-1} y}{dx^{n-1}} = z, \text{ then } \frac{dz}{dx} = fz \text{ or } dx = \frac{dz}{fz}, x = \int \frac{dz}{fz} + C.$$

After effecting this integration, express  $z$  in terms of  $x$  and  $C$ . Then, since  $z = \frac{d^{n-1} y}{dx^{n-1}}$ ,  $\frac{d^{n-1} y}{dx^{n-1}} = F(x, C)$ , an equation that may be treated by (26).

Or, since

$$\frac{d^{n-1} y}{dx^{n-1}} = z, \frac{d^{n-2} y}{dx^{n-2}} = \int z dx + c = \int \frac{z dz}{fz} + c, \text{ since } dx = \frac{dz}{fz}.$$

$$\frac{d^{n-3} y}{dx^{n-3}} = \int dx \left( \int \frac{z dz}{fz} + c \right) + c_1 = \int \frac{dz}{fz} \left( \int \frac{z dz}{fz} + c \right) + c_1, \dots$$

Continue this process until  $y$  is expressed in terms of  $z$  and  $n-1$ , arbitrary constants, and then eliminate  $z$  by the aid of the equation  $x = \int \frac{dz}{fz} + C$ .

(29) Of the form  $\frac{d^n y}{dx^n} = f \frac{d^{n-2} y}{dx^{n-2}}$ .

Let  $\frac{d^{n-2} y}{dx^{n-2}} = z$ , and the equation becomes  $\frac{d^2 z}{dx^2} = fz$ , and may be solved by (27).

- (30) Homogeneous on the supposition that  $x$  and  $y$  are of the degree 1,  $\frac{dy}{dx}$  of the degree 0,  $\frac{d^2y}{dx^2}$  of the degree  $-1$ , ...

Assume  $x = e^\theta$ ,  $y = e^\theta z$ , and by changing the variables introduce  $\theta$  and  $z$  into the equation in the place of  $x$  and  $y$ . Divide through by  $e^\theta$  and there will result an equation involving only  $z$ ,  $\frac{dz}{d\theta}$ ,  $\frac{d^2z}{d\theta^2}$ , ..., whose order may be depressed by (25).

- (31) Homogeneous on the supposition that  $x$  is of the degree 1,  $y$  of the degree  $n$ ,  $\frac{dy}{dx}$  of the degree  $n-1$ ,  $\frac{d^2y}{dx^2}$  of the degree  $n-2$ , ...

Assume  $x = e^\theta$ ,  $y = e^{n\theta} z$ , and by changing the variables introduce  $\theta$  and  $z$  into the equation in the place of  $x$  and  $y$ . The resulting equation may be freed from  $\theta$  by division and treated by (25).

- (32) Homogeneous relatively to  $y$ ,  $\frac{dy}{dx}$ ,  $\frac{d^2y}{dx^2}$ , ...

Assume  $y = e^x$ , and substitute in the given equation. Divide through by  $e^x$  and treat by (25).

- (33) Containing the first power only of the derivative of the highest order.

The equation may be *exact*.

Call its first member  $\frac{dV}{dx}$ . If  $n$  is the order of the equation, represent  $\frac{d^{n-1}y}{dx^{n-1}}$  by  $p$  and  $\frac{d^n y}{dx^n}$  by  $\frac{dp}{dx}$ . Multiply the term containing  $\frac{dp}{dx}$  by  $dx$  and integrate it as if  $p$  were the only variable, calling the result  $U_1$ ; then replacing  $p$  by  $\frac{d^{n-1}y}{dx^{n-1}}$ ,

find the complete derivative  $\frac{dU_1}{dx}$ , and form the expression  $\frac{dV}{dx} - \frac{dU_1}{dx}$ , representing it by  $\frac{dV_1}{dx}$ . If  $\frac{dV_1}{dx}$  contains the first power only of the highest derivative of  $y$ , it may itself be an exact derivative, and is to be treated precisely as the first member of the given equation  $\frac{dV}{dx}$  has been. Continue this process until a remainder  $\frac{dV_{n-1}}{dx}$  of the first order occurs.

Write this equal to zero, and see if the equation thus formed is *exact*, see (6). If so, solve it by (6), throwing its solution into the form  $V_{n-1} = C$ . A complete first integral of the given equation will be  $U_1 + U_2 + \dots + V_{n-1} = C$ . The occurrence at any step of the process of a remainder  $\frac{dV_k}{dx}$ , containing a higher power than the first of its highest derivative of  $y$ , or the failure of the resulting equation of the first order above described to be *exact*, shows that the first member of the given equation was not an exact derivative, and that this method will not apply.

(34) Of the form  $\frac{d^2y}{dx^2} + X \frac{dy}{dx} + Y \left[ \frac{dy}{dx} \right]^2 = 0$ , where  $X$  is a function of  $x$  alone and  $Y$  a function of  $y$  alone. Multiply through by  $\left[ \frac{dy}{dx} \right]^{-1}$  and the equation will become exact, and may be solved by (33).

(35) Singular integral will answer.

Call  $\frac{d^{n-1}y}{dx^{n-1}} = p$ , and  $\frac{d^n y}{dx^n} = q$ , and find  $\frac{dq}{dp}$ , regarding  $p$  and  $q$  as the only variables, and see whether  $\frac{dq}{dp}$  can be made infinite by writing equal to zero any factor containing  $p$ .

If so, eliminate  $q$  between this equation and the given equation, and if the result is a solution it will be a singular integral.

(36) General form,  $Pdx + Qdy + Rdz = 0$ .

If the equation can be reduced to the form  $Xdx + Ydy + Zdz = 0$ , where  $X$  is a function of  $x$  alone,  $Y$  a function of  $y$  alone, and  $Z$  a function of  $z$  alone, integrate each term separately, and write the sum of the integrals equal to an arbitrary constant.

If not, integrate the equation by (V.) on the supposition that one of the variables is constant and its differential zero, writing an arbitrary function of that variable in place of the arbitrary constant in the result. Transpose all the terms to the first member, and then take its complete differential, regarding all the original variables as variable, and write it equal to the first member of the given equation, and from this equation of condition determine the arbitrary function. Substitute for the arbitrary function in the first integral its value thus determined, and the result will be the solution required.

If the equation of condition contains any other variables than the one involved in the arbitrary function, they must be eliminated by the aid of the primitive equation already obtained; and if this elimination cannot be performed, the given equation is not derivable from a single primitive equation, but must have come from two simultaneous primitive equations.

In that case, assume any arbitrary equation  $f(x,y,z) = 0$  as one primitive, differentiate it, and eliminate between it its derived equation and the given equation, one variable, and its differential. There will result a differential equation containing only two variables, which may be solved by (III.), and will lead to the second primitive of the given equation.



(37) General form,  $Pdx_1 + Qdx_2 + Rdx_3 + \dots = 0$ .

If the equation can be reduced to the form  $X_1dx_1 + X_2dx_2 + X_3dx_3 + \dots = 0$ , where  $X_1$  is a function of  $x_1$  alone,  $X_2$  a function of  $x_2$  alone,  $X_3$  a function of  $x_3$  alone, etc., integrate each term separately, and write the sum of their integrals equal to an arbitrary constant.

If not, integrate the equation by (V.), on the supposition that all the variables but two are constant and their differentials zero, writing an arbitrary function of these variables in place of the arbitrary constant in the result. Transpose all the terms to the first member, and then take its complete differential, regarding all of the original variables as variable, and write it equal to the first member of the given equation, and from this equation of condition determine the arbitrary function. Substitute for the arbitrary function in the first integral its value thus determined, and the result will be the solution required.

If the equation of condition cannot, even with the aid of the primitive equation first obtained, be thrown into a form where the complete differential of the arbitrary function is given equal to an exact differential, the function cannot be determined, and the given equation is not derivable from a single primitive equation.

(38) System of simultaneous equations of the first order.

If any of the equations of the set can be integrated separately by (II.) so as to lead to single primitives, the problem can be simplified; for by the aid of these primitives a number of variables equal to the number of solved equations can be eliminated from the remaining equations of the series, and there will be formed a simpler set of simultaneous equations whose primitives, together with the primitives already found, will form the primitive system of the given equations.

There must be  $n$  equations connecting  $n + 1$  variables, in order that the system may be determinate.

Let  $x, x_1, x_2, \dots, x_n$  be the original variables. Choose

any two,  $x$  and  $x_1$ , as the independent and the principal dependent variable, and by successive eliminations form the  $n$  equations  $\frac{dx_1}{dx} = f_1(x, x_1, x_2, \dots, x_n)$ ,  $\frac{dx_2}{dx} = f_2(x, x_1, x_2, \dots, x_n)$ , ..., up to  $\frac{dx_n}{dx} = f_n(x, x_1, x_2, \dots, x_n)$ . Differentiate the first of these with respect to  $x$   $n - 1$  times, substituting for  $\frac{dx_2}{dx}$ ,  $\frac{dx_3}{dx}$ , ...,  $\frac{dx_n}{dx}$ , after each step their values in terms of the original variables. There will result  $n$  equations, which will express each of the  $n$  successive derivatives  $\frac{dx_1}{dx}$ ,  $\frac{d^2x_1}{dx^2}$ ,  $\frac{d^3x_1}{dx^3}$ , ...,  $\frac{d^nx_1}{dx^n}$ , in terms of  $x$ ,  $x_1$ ,  $x_2$ , ...,  $x_n$ . Eliminate from these all the variables except  $x$  and  $x_1$ , obtaining a single equation of the  $n$ th order between  $x$  and  $x_1$ . Solve this by (VII.), and so get a value of  $x_1$  in terms of  $x$  and  $n$  arbitrary constants. Find by differentiating this result values for  $\frac{dx_1}{dx}$ ,  $\frac{d^2x_1}{dx^2}$ , ...,  $\frac{d^{n-1}x_1}{dx^{n-1}}$ , and write them equal to the ones already obtained for them in terms of the original variables. The  $n - 1$  equations thus formed, together with the equation expressing  $x_1$  in terms of  $x$  and arbitrary constants, are the complete primitive system required.

- (39) System of simultaneous equations not of the first order. Regard each derivative of each dependent variable, from the first to the next to the highest as a new variable, and the given equations, together with the equations defining these new variables, will form a system of simultaneous equations of the first order which may be solved by (38). Eliminate the new variables representing the various derivatives from the equations of the solution, and the equations obtained will be the complete primitive system required.
- (40) All the partial derivatives taken with respect to one of the independent variables.

Integral by  $\Sigma$  is if that one were the only independent variable, replacing each arbitrary constant by an arbitrary function of the other independent variables.

10. The given equation is linear, containing three variables, and of the form  $P_1 D_1 z + Q_1 D_2 z = R$ .

Form the auxiliary system of ordinary differential equations  $\frac{dx}{P_1} = \frac{dy}{Q_1} = \frac{dz}{R}$ , and integrate by (38). Express their solutions in the form  $x = a$ ,  $y = b$ ,  $z = c$  and  $b = f(a)$ , where  $f$  is an arbitrary function.  $a = c$  is the required solution.

11. The given equation is linear, containing more than three variables, and of the form  $P_1 D_1 z + P_2 D_2 z + \dots = R$ , where  $x_1, x_2, \dots, x_n$  are the independent and  $z$  the dependent variables.

Form the auxiliary system of ordinary differential equations  $\frac{dx_1}{P_1} = \frac{dx_2}{P_2} = \dots = \frac{dz}{R}$ , and integrate them by (38). Express the solutions in the form  $x_1 = a$ ,  $x_2 = b$ ,  $x_3 = c$ , and  $b = f(a)$ ,  $c = g(a)$ , where  $f$  is an arbitrary function.  $a = z$  is the required solution.

12. The given equation is not linear, containing three variables, and of the form  $P_1 D_1 z + P_2 D_2 z = R$ .

Express  $x_2$  in terms of  $x_1$  and  $z$  from the given equation, and substitute this value thus obtained in the auxiliary system of ordinary differential equations  $\frac{dx_1}{P_1} = \frac{dx_2}{P_2} = \frac{dz}{R}$ . Deduce by integration from these equations, by (38), a value of  $x_1$  involving an arbitrary constant, and substitute it with the corresponding value of  $x_2$  in the equation  $dx = p dx + q dy$ . Integrate this result by (36), if possible; and if a single primitive equation be obtained, it will be a complete primitive of the given equation.

A singular solution may be obtained by finding the partial derivatives  $D_p z$  and  $D_q z$  from the given equation, writing them separately equal to zero, and eliminating  $p$  and  $q$  between them and the given equation.

- (44) Of the first order and not linear, containing more than three variables.  $F(x_1, x_2, \dots, x_n, z, p_1, p_2, \dots, p_n) = 0$ , where  $p_1 = D_{x_1} z$ ,  $p_2 = D_{x_2} z$ , ....

Form the linear partial differential equation  $\Sigma_i [(D_{x_i} F + p_i D_z F) D_{p_i} \Phi - D_{p_i} F (D_{x_i} \Phi + p_i D_z \Phi)] = 0$ , where  $\Phi$  is an unknown function of  $(x_1, \dots, x_n, p_1, \dots, p_n)$ , and where  $\Sigma_i$  means the sum of all the terms of the given form that can be obtained by giving  $i$  successively the values 1, 2, 3, ...,  $n$ .

Form, by (42), its auxiliary system of ordinary differential equations, and from them get, by (38),  $n - 1$  integrals,  $\Phi_1 = \alpha_1$ ,  $\Phi_2 = \alpha_2$ , ...,  $\Phi_{n-1} = \alpha_{n-1}$ . By these equations and the given equation express  $p_1, p_2, \dots, p_n$  in terms of the original variables, and substitute their values in the equation  $dz = p_1 dx_1 + p_2 dx_2 + \dots + p_n dx_n$ . Integrate this by (37), and the result will be the required complete primitive.

- (45) Of the second order and containing the derivatives of the second order only in the first degree. General form,  $RD_x^2 z + SD_x D_y z + TD_y^2 z = V$ , where  $R, S, T$ , and  $V$  may be functions of  $x, y, z, D_x z$ , and  $D_y z$ .

Call  $D_x z$   $p$  and  $D_y z$   $q$ .

Form first the equation

$$Rdy^2 - Sdx dy + Tdx^2 = 0, \quad [1]$$

and resolve it, supposing the first member not a complete square, into two equations of the form

$$dy - m_1 dx = 0, \quad dy - m_2 dx = 0. \quad [2]$$

From the first of these, and from the equation

$$Rdp dy + Tdq dx - V dx dy = 0, \quad [3]$$

combined if needful with the equation

$$dz = p dx + q dy,$$

seek to obtain two integrals  $u_1 = \alpha$ ,  $v_1 = \beta$ . Proceeding in the same way with the second equation of [2], seek two other integrals  $u_2 = \alpha_1$ ,  $v_2 = \beta_1$ ; then the first integrals of the proposed equation will be

$$u_1 = f_1 v_1, \quad u_2 = f_2 v_2, \quad [4]$$

where  $f_1$  and  $f_2$  denote arbitrary functions.

To deduce the final integral, we must either integrate one of these, or, determining from the two  $p$  and  $q$  in terms of  $x$ ,  $y$ , and  $z$ , substitute those values in the equation

$$dz = p dx + q dy,$$

which will then become integrable. Its solution will give the final integral sought.

If the values of  $m_1$  and  $m_2$  are equal, only one first integral will be obtained, and the final solution must be sought by its integration.

When it is not possible so to combine the auxiliary equations as to obtain two auxiliary integrals  $u = \alpha$ ,  $v = \beta$ , no first integral of the proposed equation exists, and this method of solution fails.

## EXAMPLES.

- (1)  $\sin x \cos y \cdot dx - \cos x \sin y \cdot dy = 0$ .    *Ans.*  $\cos y = c \cos x$ .
- (2)  $(x + y)^2 \frac{dy}{dx} = a^2$ .    *Ans.*  $y - a \tan^{-1} \frac{x + y}{a} = c$ .
- (3)  $\frac{d\theta}{d\phi} = \sin(\phi - \theta)$ .    *Ans.*  $\text{ctn} \left[ \frac{\pi}{4} - \frac{\phi - \theta}{2} \right] = \phi + c$ .
- (4)  $x \frac{dy}{dx} - y + x \sqrt{x^2 - y^2} = 0$ .    *Ans.*  $\sin^{-1} \frac{y}{x} = c - x$ .
- (5)  $(y - x)(1 + x^2)^{\frac{1}{2}} \frac{dy}{dx} = (1 + y^2)^{\frac{1}{2}}$ .  
*Ans.*  $\text{ctn} \left[ \frac{\pi}{4} - \frac{1}{2} (\tan^{-1} y - \tan^{-1} x) \right] = \tan^{-1} y + c$ .
- (6)  $\left( x \frac{dy}{dx} - y \right)^2 = a \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right] (x^2 + y^2)^{\frac{1}{2}}$ .  
*Ans.*  $2a(x^2 + y^2) = (x^2 + y^2)^{\frac{1}{2}} - x \cos c + y \sin c$ .
- (7)  $[2\sqrt{(xy)} - x] dy + y dx = 0$ .    *Ans.*  $y = ce^{-\sqrt{\frac{x}{y}}}$ .
- (8)  $(x - y^2) + 2xy \frac{dy}{dx} = 0$ .    *Ans.*  $xe^{\frac{x}{y}} = c$ .
- (9)  $(2x - y + 1)dx + (2y - x - 1) dy = 0$ .  
*Ans.*  $x^2 - xy + y^2 + x - y = c$ .
- (10)  $\frac{dy}{dx} + y \cos x = \frac{\sin 2x}{2}$ .    *Ans.*  $y = \sin x - 1 + ce^{-\sin x}$ .
- (11)  $(1 - x^2) \frac{dy}{dx} - xy = axy^2$ .    *Ans.*  $y = [c\sqrt{(1 - x^2)} - a]^{-1}$ .
- (12)  $xy(1 + xy^2) \frac{dy}{dx} = 1$ .    *Ans.*  $\frac{1}{x} = 2 - y^2 + ce^{-\frac{y^2}{x}}$ .
- (13)  $y(x^2 + y^2 + a^2) \frac{dy}{dx} + x(x^2 + y^2 - a^2) = 0$ .  
*Ans.*  $(x^2 + y^2)^2 - 2a^2(x^2 - y^2) = c$ .

$$(14) \quad xdx + ydy + \frac{xdy - ydx}{x^2 + y^2} = 0. \quad \text{Ans. } \frac{x^2 + y^2}{2} + \tan^{-1} \frac{y}{x} = c.$$

$$(15) \quad \left(\frac{dy}{dx}\right)^2 - \frac{a^2}{x^2} = 0. \quad \text{Ans. } (y - a \log x - c)(y + a \log x - c) = 0.$$

$$(16) \quad \left(\frac{dy}{dx}\right)^2 + 2y \operatorname{ctn} x \frac{dy}{dx} = y^2. \quad \text{Ans. } \left(y \sin^2 \frac{x}{2} - c\right) \left(y \cos^2 \frac{x}{2} - c\right) = 0.$$

$$(17) \quad \frac{dy}{dx} \left(\frac{dy}{dx} + y\right) = x(x + y). \quad \text{Ans. } (2y - x^2 - c)[\log(x + y - 1) + x - c] = 0.$$

$$(18) \quad \left(\frac{dy}{dx}\right)^3 - (x^2 + xy + y^2) \left(\frac{dy}{dx}\right)^2 + (x^3y + x^2y^2 + xy^3) \frac{dy}{dx} - x^3y^3 = 0. \quad \text{Ans. } \left(y - \frac{x^3}{3} - c\right) \left(x + \frac{1}{y} - c\right) \left(\log y - \frac{x^2}{2} - c\right) = 0.$$

$$(19) \quad \left(1 - y^2 - \frac{y^4}{x^2}\right) \left(\frac{dy}{dx}\right)^2 - \frac{2y}{x} \frac{dy}{dx} + \frac{y^2}{x^2} = 0. \quad \text{Ans. } \left(y + \log \frac{x + \sqrt{x^2 + y^2}}{y} - c\right) \left(y - \log \frac{x + \sqrt{x^2 + y^2}}{y} - c\right) = 0.$$

$$(20) \quad y = x \frac{dy}{dx} + \frac{dy}{dx} - \left(\frac{dy}{dx}\right)^2. \quad \text{Ans. } y = cx + c - c^2. \quad \text{Singular solution, } y = \frac{(x+1)^2}{4}.$$

$$(21) \quad y = y \left(\frac{dy}{dx}\right)^2 + 2x \frac{dy}{dx}. \quad \text{Ans. } y^2 = 2cx + c^2.$$

$$(22) \quad \left[1 - \left(\frac{dy}{dx}\right)^2\right] xy = (x^2 - y^2 - a^2) \frac{dy}{dx}. \quad \text{Ans. } y^2 - cx^2 + \frac{a^2c}{1+c} = 0.$$

$$(23) \quad y = 2x \frac{dy}{dx} + y^2 \left(\frac{dy}{dx}\right)^3. \quad \text{Ans. } y^2 = 2cx + c^3.$$

$$(24) \quad x^3 \left(\frac{dy}{dx}\right)^2 + x^2 y \frac{dy}{dx} + a^3 = 0. \quad \text{Ans. } c^2 + cxy + a^3x = 0.$$

$$(25) f\left[x\left(\frac{dy}{dx}\right)^2\right] + 2x\frac{dy}{dx} - y = 0. \quad \text{Ans. } (b+y)^2 = 4ax, f(a) + b = 0.$$

$$(26) x^2 - \frac{xy}{\frac{dy}{dx}} = f\left[y^2 - xy\frac{dy}{dx}\right]. \quad \text{Ans. } \frac{ax^2}{y^2 - a} = b, f(a) + b = 0.$$

$$(27) \frac{d^4y}{dx^4} - 4\frac{d^3y}{dx^3} + 6\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + y = 0. \quad \text{Ans. } y = (c_0 + c_1x + c_2x^2 + c_3x^3)e^x.$$

$$(28) \frac{d^2y}{dx^2} - 2k\frac{dy}{dx} + k^2y = e^x. \quad \text{Ans. } y = (c_1 + c_2x)e^{kx} + \frac{e^x}{(k-1)^2}.$$

$$(29) \frac{d^4y}{dx^4} + 2\frac{d^2y}{dx^2} + y = e^x. \quad \text{Ans. } y = \frac{e^x}{4} + (A + Bx)\cos x + (C + Dx)\sin x.$$

$$(30) \frac{d^3y}{dx^3} - 2\frac{dy}{dx} + 4y = e^x \cos x. \quad \text{Ans. } y = Ae^{-2x} + e^x\left[\left(B - \frac{x}{20}\right)\cos x + \left(C + \frac{3x}{20}\right)\sin x\right].$$

$$(31) \frac{d^4y}{dx^4} - 2\frac{d^3y}{dx^3} + \frac{d^2y}{dx^2} = x^3. \quad \text{Ans. } y = (A + Bx)e^x + (C + Dx) + 12x^2 + 3x^3 + \frac{x^4}{2} + \frac{x^5}{20}.$$

$$(32) \frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = x^2. \quad \text{Ans. } y = (A + Bx)e^{2x} + \frac{1}{8}(2x^2 + 4x + 3).$$

$$(33) \frac{d^2y}{dx^2} + y = \cos x. \quad \text{Ans. } y = A\cos x + B\sin x + \frac{x}{2}\sin x.$$

$$(34) \frac{d^2y}{dx^2} + 4y = x\sin^2 x. \quad \text{Ans. } y = \left(A - \frac{x}{32}\right)\cos 2x + \left(B - \frac{x^2}{16}\right)\sin 2x + \frac{x}{8}.$$

$$(35) x^2\frac{d^2y}{dx^2} - x\frac{dy}{dx} + 2y = x\log x. \quad \text{Ans. } y = Ax\cos(\log x) + Bx\sin(\log x) + x\log x.$$



- (36)  $x^2 \frac{d^2 y}{dx^2} - x^2 \frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} - 2y = x^2 + 3x$ .  
*Ans.*  $y = x(A + B \log x) + Cx^2 + \frac{x^3}{4} - 3x \left(1 + \frac{(\log x)^2}{2}\right)$
- (37)  $\frac{d^2 y}{dx^2} + \frac{2}{x} \frac{dy}{dx} - n^2 y = 0$ . *Ans.*  $y = \frac{1}{x} (Ae^{nx} + Be^{-nx})$ .
- (38)  $\frac{d^2 y}{dx^2} + \tan x \frac{dy}{dx} + \cos^2 x \cdot y = 0$ .  
*Ans.*  $y = A \cos(\sin x) + B \sin(\sin x)$ .
- (39)  $(1 - x^2)^2 \frac{d^2 y}{dx^2} + y = 0$ .  
*Ans.*  $y = \sqrt{1 - x^2} \left( A + B \log \frac{1+x}{1-x} \right)$ .
- (40)  $(1 + x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 2y = 0$ .  
*Ans.*  $y = Bx + A(1 - x^2)$ .
- (41)  $\frac{d^2 y}{dx^2} - \frac{x}{x-1} \frac{dy}{dx} + \frac{1}{x-1} y = x - 1$ .  
*Ans.*  $y = Ae^x + Bx - (1 + x^2)$ .
- (42)  $x^2 \frac{d^2 y}{dx^2} - 2x(1+x) \frac{dy}{dx} + 2(1+x)y = x^2$ .  
*Ans.*  $y = Axe^x + Bx - \frac{x^2}{2}$ .
- (43)  $\sin^2 x \frac{d^2 y}{dx^2} - 2y = 0$ . *Ans.*  $y = A \operatorname{ctn} x + B(1 - x \operatorname{ctn} x)$ .
- (44)  $\frac{d^2 y}{dx^2} + \frac{1}{x^2 \log x} y = e^x \left( \frac{2}{x} + \log x \right)$ .  
*Ans.*  $y = A \log x + e^x \log x + B \left( \log x \int \frac{dx}{\log x} - x \right)$ .
- (45)  $\frac{d^2 y}{dx^2} - 2 \left( n - \frac{a}{x} \right) \frac{dy}{dx} + \left( n^2 - 2 \frac{na}{x} \right) y = e^{ax}$ .  
*Ans.*  $y = e^{ax} \left( A + \frac{B}{x^{2a-1}} + \frac{x^2}{2(2a-1)} \right)$ .
- (46)  $\frac{d^2 y}{dx^2} - \frac{2}{x} \frac{dy}{dx} + \left( a^2 + \frac{2}{x^2} \right) y = 0$ .  
*Ans.*  $y = x(A \cos ax + B \sin ax)$ .

$$(47) \frac{d^2 y}{dx^2} - 2bx \frac{dy}{dx} + b^2 x^2 y = 0. \quad \text{Ans. } y = e^{\frac{bx^2}{2}} (A \cos x \sqrt{b} + B \sin x \sqrt{b}).$$

$$(48) \frac{d^2 y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 3)y = e^{x^2}. \quad \text{Ans. } y = e^{x^2} (Ae^x + Be^{-x} - 1).$$

$$(49) (1 - x^2) \frac{d^2 y}{dx^2} - 4x \frac{dy}{dx} - (1 + x^2)y = x. \quad \text{Ans. } y = \frac{1}{1 - x^2} (x + A \cos x + B \sin x).$$

$$(50) \frac{d^2 y}{dx^2} - \frac{1}{\sqrt{x}} \frac{dy}{dx} + \frac{x + \sqrt{x} - 8}{4x^2} y = 0. \quad \text{Ans. } y = e^{\sqrt{x}} \left( Ax^2 + \frac{B}{x} \right).$$

$$(51) \frac{d^2 y}{dx^2} + 2n \cot nx \frac{dy}{dx} + (m^2 - n^2)y = 0. \quad \text{Ans. } y = (A \cos mx + B \sin mx) \csc nx.$$

$$(52) (x^2 - 1) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - c^2 y = 0. \quad \text{Ans. } y = A(x + \sqrt{x^2 - 1})^c + B(x - \sqrt{x^2 - 1})^c.$$

$$(53) \frac{d^2 y}{dx^2} + \frac{2}{x} \frac{dy}{dx} + \frac{a^2}{x^4} y = 0. \quad \text{Ans. } y = A \sin \frac{a}{x} + B \cos \frac{a}{x}.$$

$$(54) \frac{d^2 y}{dx^2} - \frac{3x+1}{x^2-1} \frac{dy}{dx} + y \left[ \frac{6(x+1)}{(x-1)(3x+5)} \right]^2 = 0. \quad \text{Ans. } y = [A + B \log((x-1)^3(3x+5))] \sqrt{(x-1)^3(3x+5)}.$$

$$(55) (1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} - c^2 y = 0. \quad \text{Ans. } y = Ae^{c \sin^{-1} x} + Be^{-c \sin^{-1} x}.$$

$$(56) (1 + ax^2) \frac{d^2 y}{dx^2} + ax \frac{dy}{dx} - n^2 y = 0. \quad \text{Ans. } y = A(\sqrt{1 + ax^2} + x\sqrt{a})^{\frac{n}{\sqrt{a}}} + B(\sqrt{1 + ax^2} - x\sqrt{a})^{-\frac{n}{\sqrt{a}}}.$$

$$(57) (x-1)(x-2) \frac{d^2 y}{dx^2} - (2x-3) \frac{dy}{dx} + 2y = 0. \quad \text{Ans. } y = c(x-2)^2 + c'(x-2) [(x-2)^2].$$

$$(58) (3-x) \frac{d^2 y}{dx^2} - (9-4x) \frac{dy}{dx} + (6-3x)y = 0.$$

$$\text{Ans. } y = ce^{2x} + c'e^{3x} \left( \frac{183}{8} - \frac{75}{4}x + \frac{21}{4}x^2 - \frac{1}{2}x^3 \right).$$

$$(59) (a^2 - x^2) \frac{d^2 y}{dx^2} - 8x \frac{dy}{dx} - 12y = 0.$$

$$\text{Ans. } y = \frac{c}{(a-x)^3} + c' \frac{(a^2 + 3x^2)}{(a^2 - x^2)^3}.$$

$$(60) \frac{d^2 y}{dx^2} + \frac{2}{x} \frac{dy}{dx} + \left( n^2 - \frac{2}{x^2} \right) y = 0.$$

$$\text{Ans. } y = \frac{1}{x^2} [A (\sin nx - nx \cos nx) + B (\cos nx + nx \sin nx)].$$

$$(61) \frac{d^2 y}{dx^2} + \frac{1}{x} \frac{dy}{dx} = 0.$$

$$\text{Ans. } y = c \log x + c'.$$

$$(62) \left( 2x^2 \frac{dy}{dx} + x^2 y \right) \frac{d^2 y}{dx^2} + 4x^2 \left( \frac{dy}{dx} \right)^2 + 2xy \frac{dy}{dx} = 0.$$

$$\text{Ans. } c^2 + cxy = c'x.$$

$$(63) \left( x^2 + 2y^2 \frac{dy}{dx} \right) \frac{d^2 y}{dx^2} + 2y \left( \frac{dy}{dx} \right)^2 + 3x \frac{dy}{dx} + y = 0.$$

Find a first integral.

$$\text{Ans. } x^2 \frac{dy}{dx} + y^2 \left( \frac{dy}{dx} \right)^2 + xy = c.$$

$$(64) x^2 \frac{d^3 y}{dx^3} + x \frac{d^2 y}{dx^2} + (2xy - 1) \frac{dy}{dx} + y^2 = 0.$$

Find a first integral.

$$\text{Ans. } x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + xy^2 = c.$$

$$(65) \frac{dx}{x-a} + \frac{dy}{y-b} + \frac{dz}{z-c} = 0. \quad \text{Ans. } (x-a)(y-b)(z-c) = c.$$

$$(66) (y+z) dx + dy + dz = 0.$$

$$\text{Ans. } e^x (y+z) = c.$$

$$(67) \frac{dx}{dt} + 4x + \frac{y}{4} = 0, \quad \frac{dy}{dt} + 3y - x = 0.$$

$$\text{Ans. } x = ce^{\frac{7t}{4}} - \frac{y}{2}, \quad y = (ct + c_1) e^{-\frac{3t}{4}}.$$

$$(68) \frac{d^2x}{dt^2} + m^2x = 0, \quad \frac{d^2y}{dt^2} - m^2y = 0.$$

$$\text{Ans. } x = A \sin mt + B \cos mt, \quad x + y = Ct + D.$$

$$(69) D_x z = \frac{y}{x+z}.$$

$$\text{Ans. } e^{-\frac{z}{y}}(x+y+z) = \phi y.$$

$$(70) xx D_x z + yz D_y z = xy.$$

$$\text{Ans. } z^2 = xy + \phi\left(\frac{x}{y}\right).$$

$$(71) D_x z \cdot D_y z = 1.$$

$$\text{Ans. } z = ax + \frac{y}{a} + b.$$

$$(72) x^2 D_x^2 z + 2xy D_x D_y z + y^2 D_y^2 z = 0. \quad \text{Ans. } z = x\phi\left(\frac{y}{x}\right) + \psi\left(\frac{y}{x}\right).$$

$$(73) (D_y z)^2 D_x^2 z - 2D_x z D_y z D_x D_y z + (D_x z)^2 D_y^2 z = 0.$$

$$\text{Ans. } y = x\phi z + \psi z.$$

$$(74) D_x z \cdot D_x D_y z - D_y z \cdot D_x^2 z = 0.$$

$$\text{Ans. } x = \phi y + \psi z.$$





**APPENDIX.**





## CHAPTER V.

## INTEGRATION.

74. We are now able to extend materially our list of *formulas for direct integration* (Art. 55), one of which may be obtained from each of the derivative formulas in our last chapter. The following set contains the most important of these:—

$D_x \log x = \frac{1}{x}$	gives $\int_x \frac{1}{x} = \log x$ .
$D_x a^x = a^x \log a$	“ $\int_x a^x \log a = a^x$ .
$D_x e^x = e^x$	“ $\int_x e^x = e^x$ .
$D_x \sin x = \cos x$	“ $\int_x \cos x = \sin x$ .
$D_x \cos x = -\sin x$	“ $\int_x (-\sin x) = \cos x$ .
$D_x \log \sin x = \operatorname{ctn} x$	“ $\int_x \operatorname{ctn} x = \log \sin x$ .
$D_x \log \cos x = -\tan x$	“ $\int_x (-\tan x) = \log \cos x$ .
$D_x \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$	“ $\int_x \frac{1}{\sqrt{1-x^2}} = \sin^{-1} x$ .
$D_x \tan^{-1} x = \frac{1}{1+x^2}$	“ $\int_x \frac{1}{1+x^2} = \tan^{-1} x$ .
$D_x \operatorname{vers}^{-1} x = \frac{1}{\sqrt{2x-x^2}}$	“ $\int_x \frac{1}{\sqrt{2x-x^2}} = \operatorname{vers}^{-1} x$ .

The second, fifth, and seventh in the above list can be written in the more convenient forms,



$$\int_x a^x = \frac{a^x}{\log a};$$

$$\int_x \sin x = -\cos x;$$

$$\int_x \tan x = -\log \cos x.$$

75. When the expression to be integrated does not come under any of the forms in the preceding list, *it can often be prepared for integration by a suitable change of variable*, the new variable, of course, being a function of the old. This method is called *integration by substitution*, and is based upon a formula easily

deduced from  $D_x(Fy) = D_y Fy \cdot D_x y;$

which gives immediately

$$Fy = \int_x (D_y Fy \cdot D_x y).$$

Let  $u = D_y Fy,$

then  $Fy = \int_y u,$

and we have  $\int_y u = \int_x (uD_x y);$

or, interchanging  $x$  and  $y,$

$$\int_x u = \int_y (uD_y x). \quad [1]$$

For example, required  $\int_x (a + bx)^n.$

Let  $z = a + bx,$

and then  $\int_x (a + bx)^n = \int_x z^n = \int_x (z^n \cdot D_x x),$  by [1];

but  $x = \frac{z}{b} - \frac{a}{b},$

$$D_x x = \frac{1}{b};$$

hence  $\int_x (a + bx)^n = \frac{1}{b} \int_x z^n = \frac{1}{b} \frac{z^{n+1}}{n+1}.$

Substituting for  $z$  its value, we have

$$\int (a + bx)^n = \frac{1}{b} \frac{(a + bx)^{n+1}}{n + 1}.$$

EXAMPLE.

Find  $\int \frac{1}{a + bx}$ .

Ans.  $\frac{1}{b} \log(a + bx)$ .

76. If  $fz$  represents a function that can be integrated,  $f(a + bx)$  can always be integrated; for, if

$$z = a + bx,$$

then

$$D_x x = \frac{1}{b}$$

and

$$\int f(a + bx) = \int fz = \int fz D_x x = \frac{1}{b} \int fz.$$

EXAMPLES.

Find

(1)  $\int \sin ax$ .

Ans.  $-\frac{1}{a} \cos ax$ .

(2)  $\int \cos ax$ .

Ans.  $\frac{1}{a} \sin ax$ .

(3)  $\int \tan ax$ .

(4)  $\int \text{ctn } ax$ .

77. Required  $\int \frac{1}{\sqrt{a^2 - x^2}}$ .

$$\int \frac{1}{\sqrt{a^2 - x^2}} = \frac{1}{a} \int \frac{1}{\sqrt{1 - \left(\frac{x}{a}\right)^2}}$$

Let

$$z = \frac{x}{a},$$

then

$$x = az,$$

$$D_x x = a,$$

$$\begin{aligned} \frac{1}{a} \int_x \frac{1}{\sqrt{\left[1 - \left(\frac{x}{a}\right)^2\right]}} &= \frac{1}{a} \int_x \frac{1}{\sqrt{(1-z^2)}} = \frac{1}{a} \int_x \frac{1}{\sqrt{(1-z^2)}} D_x x \\ &= \int_x \frac{1}{\sqrt{(1-z^2)}} = \sin^{-1} z = \sin^{-1} \frac{x}{a}. \end{aligned}$$

## EXAMPLES.

Find

$$(1) \int_x \frac{1}{a^2 + x^2}. \quad \text{Ans. } \frac{1}{a} \tan^{-1} \frac{x}{a}.$$

$$(2) \int_x \frac{1}{\sqrt{(2ax - x^2)}}. \quad \text{Ans. } \text{vers}^{-1} \frac{x}{a}.$$

$$78. \text{ Required } \int_x \frac{1}{\sqrt{(x^2 + a^2)}}.$$

Let

$$z = x + \sqrt{(x^2 + a^2)};$$

then

$$z - x = \sqrt{(x^2 + a^2)},$$

$$z^2 - 2zx + x^2 = x^2 + a^2,$$

$$2zx = z^2 - a^2,$$

$$x = \frac{z^2 - a^2}{2z},$$

$$\sqrt{(x^2 + a^2)} = z - x = z - \frac{z^2 - a^2}{2z} = \frac{z^2 + a^2}{2z},$$

$$D_x x = \frac{z^2 + a^2}{2z^2}.$$

$$\begin{aligned} \int_x \frac{1}{\sqrt{(x^2 + a^2)}} &= \int_x \frac{2z}{z^2 + a^2} = \int_x \frac{2z}{z^2 + a^2} D_x x \\ &= \int_x \frac{2z}{z^2 + a^2} \frac{z^2 + a^2}{2z^2} = \int_x \frac{1}{z} = \log z = \log(x + \sqrt{x^2 + a^2}). \end{aligned}$$

## EXAMPLE.

$$\text{Find } \int_x \frac{1}{\sqrt{(x^2 - a^2)}}.$$

$$\text{Ans. } \log(x + \sqrt{x^2 - a^2}).$$

79. When the expression to be integrated can be factored, the required integral can often be obtained by the use of a formula

deduced from  $D_x(uv) = uD_xv + vD_xu$ ,

which gives  $uv = \int_x uD_xv + \int_x vD_xu$

or  $\int_x uD_xv = uv - \int_x vD_xu$ . [1]

This method is called *integrating by parts*.

(a) For example, required  $\int_x \log x$ .

$\log x$  can be regarded as the product of  $\log x$  by 1.

Call  $\log x = u$  and  $1 = D_xv$ ,

then  $D_xu = \frac{1}{x}$ ,

$$v = x;$$

and we have

$$\begin{aligned} \int_x \log x &= \int_x 1 \log x = \int_x uD_xv = uv - \int_x vD_xu \\ &= x \log x - \int_x \frac{x}{x} = x \log x - x. \end{aligned}$$

#### EXAMPLE.

Find  $\int_x x \log x$ .

*Suggestion:* Let  $\log x = u$  and  $x = D_xv$ .

$$\text{Ans. } \frac{1}{2} x^2 \left( \log x - \frac{1}{2} \right)$$

80. Required  $\int_x \sin^2 x$ .

Let  $u = \sin x$  and  $D_xv = \sin x$ ,

then  $D_xu = \cos x$ ,

$$v = -\cos x,$$

but

$$\cos^2 x = 1 - \sin^2 x,$$

so

$$\int_x \cos^2 x = \int_x 1 - \int_x \sin^2 x = x - \int_x \sin^2 x$$

and

$$\int_x \sin^2 x = x - \sin x \cos x - \int_x \sin^2 x.$$

$$2 \int_x \sin^2 x = x - \sin x \cos x.$$

$$\int_x \sin^2 x = \frac{1}{2}(x - \sin x \cos x).$$

## EXAMPLES.

(1) Find  $\int_x \cos^2 x$ .

$$\text{Ans. } \frac{1}{2}(x + \sin x \cos x).$$

(2)  $\int_x \sin x \cos x$ .

$$\text{Ans. } \frac{\sin^2 x}{2}.$$

81. *Very often both methods described above are required in the same integration.*

(a) Required  $\int_x \sin^{-1} x$ .Let  $\sin^{-1} x = y,$ 

then

$$x = \sin y;$$

$$D_y x = \cos y,$$

$$\int_x \sin^{-1} x = \int_x y = \int_y y \cos y.$$

Let  $u = y$  and  $D_y v = \cos y;$ 

then

$$D_y u = 1,$$

$$v = \sin y,$$

and

$$\int_y y \cos y = y \sin y - \int_y \sin y = y \sin y + \cos y = x \sin^{-1} x + \sqrt{1-x^2}.$$

Any inverse or anti-function can be integrated by this method if the direct function is integrable.

(b) Thus,  $\int_x f^{-1} x = \int_x y = \int_y y D_y f y = y f y - \int_y f y$ 

where

$$y = f^{-1} x.$$

EXAMPLES.

(1) Find  $\int_x \cos^{-1}x$ . *Ans.*  $x \cos^{-1}x - \sqrt{(1-x^2)}$ .

(2)  $\int_x \tan^{-1}x$ . *Ans.*  $x \tan^{-1}x - \frac{1}{2} \log(1+x^2)$ .

(3)  $\int_x \text{vers}^{-1}x$ . *Ans.*  $(x-1) \text{vers}^{-1}x + \sqrt{(2x-x^2)}$ .

82. Sometimes an algebraic transformation, either alone or in combination with the preceding methods, is useful.

(a) Required  $\int_x \frac{1}{x^2-a^2}$ .

$$\frac{1}{x^2-a^2} = \frac{1}{2a} \left( \frac{1}{x-a} - \frac{1}{x+a} \right),$$

and, by Art. 75 (Ex.),

$$\int_x \frac{1}{x^2-a^2} = \frac{1}{2a} [\log(x-a) - \log(x+a)] = \frac{1}{2a} \log \frac{x-a}{x+a}.$$

(b) Required  $\int_x \sqrt{\left(\frac{1+x}{1-x}\right)}$ .

$$\sqrt{\left(\frac{1+x}{1-x}\right)} = \frac{1+x}{\sqrt{(1-x^2)}} = \frac{1}{\sqrt{(1-x^2)}} + \frac{x}{\sqrt{(1-x^2)}},$$

$$\int_x \frac{1}{\sqrt{(1-x^2)}} = \sin^{-1}x.$$

$\int_x \frac{x}{\sqrt{(1-x^2)}}$  can be readily obtained by substituting  $y = (1-x^2)$ , and is  $-\sqrt{(1-x^2)}$ ;

hence  $\int_x \sqrt{\left(\frac{1+x}{1-x}\right)} = \sin^{-1}x - \sqrt{(1-x^2)}$ .

(c) Required  $\int_x \sqrt{(a^2-x^2)}$ .

$$\sqrt{(a^2-x^2)} = \frac{a^2-x^2}{\sqrt{(a^2-x^2)}} = \frac{a^2}{\sqrt{(a^2-x^2)}} - \frac{x^2}{\sqrt{(a^2-x^2)}},$$

and  $\int_x \sqrt{a^2 - x^2} = a^2 \int_x \frac{1}{\sqrt{a^2 - x^2}} - \int_x \frac{x^2}{\sqrt{a^2 - x^2}},$

whence  $\int_x \sqrt{a^2 - x^2} = a^2 \sin^{-1} \frac{x}{a} - \int_x \frac{x^2}{\sqrt{a^2 - x^2}},$  by Art. 77;

but  $\int_x \sqrt{a^2 - x^2} = x\sqrt{a^2 - x^2} + \int_x \frac{x^2}{\sqrt{a^2 - x^2}},$

by integration by parts, if we let

$$u = \sqrt{a^2 - x^2} \text{ and } D_x v = 1.$$

Adding our two equations, we have

$$2 \int_x \sqrt{a^2 - x^2} = x\sqrt{a^2 - x^2} + a^2 \sin^{-1} \frac{x}{a};$$

and  $\therefore \int_x \sqrt{a^2 - x^2} = \frac{1}{2} \left( x\sqrt{a^2 - x^2} + a^2 \sin^{-1} \frac{x}{a} \right).$

#### EXAMPLES.

Find

(1)  $\int_x \sqrt{x^2 + a^2}.$

Ans.  $\frac{1}{2} [x\sqrt{x^2 + a^2} + a^2 \log(x + \sqrt{x^2 + a^2})].$

(2)  $\int_x \sqrt{x^2 - a^2}.$

Ans.  $\frac{1}{2} [x\sqrt{x^2 - a^2} - a^2 \log(x + \sqrt{x^2 - a^2})].$

#### Applications.


83. To find the area of a segment of a circle.

Let the equation of the circle be

$$x^2 + y^2 = a^2,$$

and let the required segment be cut off by the double ordinates through  $(x_0, y_0)$  and  $(x, y)$ . Then the required area

$$A = 2 \int_x y + C.$$



From the equation of the circle,

$$y = \sqrt{(a^2 - x^2)},$$

hence

$$A = 2 \int_x \sqrt{(a^2 - x^2)} + C;$$

and therefore, by Art. 82 (c),

$$A = x\sqrt{(a^2 - x^2)} + a^2 \sin^{-1} \frac{x}{a} + C.$$

As the area is measured from the ordinate  $y_0$  to the ordinate  $y$ ,

$$A = 0 \text{ when } x = x_0;$$

therefore

$$0 = x_0\sqrt{(a^2 - x_0^2)} + a^2 \sin^{-1} \frac{x_0}{a} + C,$$

$$C = -x_0\sqrt{(a^2 - x_0^2)} - a^2 \sin^{-1} \frac{x_0}{a},$$

and we have

$$A = x\sqrt{(a^2 - x^2)} + a^2 \sin^{-1} \frac{x}{a} - x_0\sqrt{(a^2 - x_0^2)} - a^2 \sin^{-1} \frac{x_0}{a}$$

If  $x_0 = 0$ , and the segment begins with the axis of Y,

$$A = x\sqrt{(a^2 - x^2)} + a^2 \sin^{-1} \frac{x}{a}.$$

If, at the same time,  $x = a$ , the segment becomes a semicircle, and

$$A = a\sqrt{(a^2 - a^2)} + a^2 \sin^{-1} \frac{a}{a} = \frac{\pi a^2}{2}.$$

The area of the whole circle is  $\pi a^2$ .





## EXAMPLES.

(1) Show that, in the case of an ellipse,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

the area of a segment beginning with any ordinate  $y_0$  is

$$A = \frac{b}{a} \left[ x \sqrt{(a^2 - x^2)} + a^2 \sin^{-1} \frac{x}{a} - x_0 \sqrt{(a^2 - x_0^2)} - a^2 \sin^{-1} \frac{x_0}{a} \right].$$

That if the segment begins with the minor axis,

$$A = \frac{b}{a} \left[ x \sqrt{(a^2 - x^2)} + a^2 \sin^{-1} \frac{x}{a} \right].$$

That the area of the whole ellipse is  $\pi ab$ .

(2) The area of a segment of the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

is 
$$A = \frac{b}{a} \left[ x \sqrt{(x^2 - a^2)} - a^2 \log(x + \sqrt{x^2 - a^2}) - x_0 \sqrt{(x_0^2 - a^2)} + a^2 \log(x_0 + \sqrt{x_0^2 - a^2}) \right].$$

If  $x_0 = a$ , and the segment begins at the vertex,

$$A = \frac{b}{a} \left[ x \sqrt{(x^2 - a^2)} - a^2 \log(x + \sqrt{x^2 - a^2}) + a^2 \log a \right].$$

84. To find the length of any arc of a circle, the coördinates of its extremities being  $(x_0, y_0)$  and  $(x, y)$ .

By Art. 52, 
$$s = \int_x \sqrt{[1 + (D_x y)^2]}.$$

From the equation of the circle,

$$x^2 + y^2 = a^2,$$

we have

$$2x + 2yD_x y = 0,$$

$$D_x y = -\frac{x}{y},$$

$$1 + (D_x y)^2 = \frac{x^2 + y^2}{y^2} = \frac{a^2}{y^2},$$

$$s = \int \frac{a}{y} = a \int \frac{1}{\sqrt{(a^2 - x^2)}} = a \sin^{-1} \frac{x}{a} + C. \quad (\text{Art. 77.})$$

When

$$x = x_0, \quad s = 0;$$

hence

$$0 = a \sin^{-1} \frac{x_0}{a} + C,$$

$$C = -a \sin^{-1} \frac{x_0}{a},$$

and

$$s = a \left( \sin^{-1} \frac{x}{a} - \sin^{-1} \frac{x_0}{a} \right).$$

If  $x_0 = 0$ , and the arc is measured from the highest point of the circle,

$$s = a \sin^{-1} \frac{x}{a}.$$

If the arc is a quadrant,  $x = a$ ,

$$s = a \sin^{-1}(1) = \frac{\pi a}{2},$$

and the whole circumference =  $2\pi a$ .

85. To find the length of an arc of the parabola  $y^2 = 2mx$ .

We have

$$2yD_x y = 2m;$$

$$D_x y = \frac{m}{y};$$

$$\sqrt{[1 + (D_x y)^2]} = \sqrt{\left(\frac{m^2 + y^2}{y^2}\right)} = \frac{1}{y} \sqrt{(m^2 + y^2)};$$

$$s = \int_x \left[ \frac{1}{y} \sqrt{(m^2 + y^2)} \right] = \int_y \left[ \frac{1}{y} \sqrt{(m^2 + y^2)} D_y x \right];$$

$$D_y x = \frac{1}{D_x y} = \frac{y}{m}, \quad \text{by Art. 73;}$$

$$s = \frac{1}{m} \int_y \sqrt{m^2 + y^2} = \frac{1}{2m} [y \sqrt{m^2 + y^2} + m^2 \log(y + \sqrt{m^2 + y^2})] + C,$$

by Art. 82, Ex. 1.

If the arc is measured from the vertex,

$$s = 0 \text{ when } y = 0;$$

$$0 = \frac{1}{2m} (m^2 \log m) + C,$$

$$C = -\frac{1}{2} m \log m,$$

and 
$$s = \frac{1}{2} \left[ \frac{y \sqrt{(m^2 + y^2)}}{m} + m \log \frac{y + \sqrt{(m^2 + y^2)}}{m} \right].$$

EXAMPLE.

Find the length of the arc of the curve  $x^2 = 27y^2$  included between the origin and the point whose abscissa is 15.

*Ans.* 19.

A

SHORT TABLE OF INTEGRALS.

COMPILED BY

B. O. PEIRCE,

HOLLIS PROFESSOR OF MATHEMATICS AND NATURAL PHILOSOPHY  
IN HARVARD UNIVERSITY.



BOSTON, U.S.A.:

PUBLISHED BY GINN & COMPANY.

1895.

*The compiler will be grateful to any person who may send  
notice of errors in these formulas to*

**B. O. PEIRCE,**

*Harvard College, Cambridge.*

## I. FUNDAMENTAL FORMS.

---

1.  $\int a dx = ax.$
2.  $\int a f(x) dx = a \int f(x) dx.$
3.  $\int \frac{dx}{x} = \log x.$
4.  $\int x^m dx = \frac{x^{m+1}}{m+1},$  when  $m$  is different from  $-1.$
5.  $\int e^x dx = e^x.$
6.  $\int a^x \log a dx = a^x.$
7.  $\int \frac{dx}{1+x^2} = \tan^{-1} x.$
8.  $\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x.$
9.  $\int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x.$
10.  $\int \frac{dx}{\sqrt{2x-x^2}} = \text{versin}^{-1} x.$
11.  $\int \cos x dx = \sin x.$
12.  $\int \sin x dx = -\cos x.$

$$13. \int \operatorname{ctn} x dx = \log \sin x.$$

$$14. \int \tan x dx = -\log \cos x.$$

$$15. \int \tan x \sec x dx = \sec x.$$

$$16. \int \sec^2 x dx = \tan x.$$

$$17. \int \csc^2 x dx = -\operatorname{ctn} x.$$

In the following formulas,  $u$ ,  $v$ ,  $w$ , and  $y$  represent any functions of  $x$ :

$$18. \int (u + v + w + \text{etc.}) dx = \int u dx + \int v dx + \int w dx + \text{etc.}$$

$$19a. \int u dv = uv - \int v du.$$

$$19b. \int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx.$$

$$20. \int f(y) dx = \int \frac{f(y) dy}{\frac{dy}{dx}}.$$

## II. RATIONAL ALGEBRAIC FUNCTIONS.

A. — EXPRESSIONS INVOLVING  $(a + bx)$ .

The substitution of  $y$  or  $z$  for  $x$ , where  $y = bx = a + bx$ , gives

$$21. \int (a + bx)^m dx = \frac{1}{b} \int y^m dy.$$

$$22. \int x(a + bx)^m dx = \frac{1}{b^2} \int y^m (y - a) dy.$$

$$23. \int x^2(a + bx)^m dx = \frac{1}{b^{n+1}} \int y^m (y - a)^2 dy.$$

$$24. \int \frac{x^n dx}{(a + bx)^m} = \frac{1}{b^{n+1}} \int \frac{(y - a)^n dy}{y^m}.$$

$$25. \int \frac{dx}{x^n (a + bx)^m} = -\frac{1}{a^{m+n-1}} \int \frac{(z - b)^{m+n-2} dz}{z^n}.$$

Whence

$$26. \int \frac{dx}{a + bx} = \frac{1}{b} \log(a + bx).$$

$$27. \int \frac{dx}{(a + bx)^2} = -\frac{1}{b(a + bx)}.$$

$$28. \int \frac{dx}{(a + bx)^3} = -\frac{1}{2b(a + bx)^2}.$$

$$29. \int \frac{x dx}{a + bx} = \frac{1}{b^2} [a + bx - a \log(a + bx)].$$

$$30. \int \frac{x dx}{(a + bx)^2} = \frac{1}{b^2} \left[ \log(a + bx) + \frac{a}{a + bx} \right].$$



$$31. \int \frac{x dx}{(a+bx)^2} = \frac{1}{b^2} \left[ -\frac{1}{a+bx} + \frac{a}{2(a+bx)^2} \right].$$

$$32. \int \frac{x^2 dx}{a+bx} = \frac{1}{b^2} \left[ \frac{1}{2}(a+bx)^2 - 2a(a+bx) + a^2 \log(a+bx) \right].$$

$$33. \int \frac{x^2 dx}{(a+bx)^2} = \frac{1}{b^2} \left[ a+bx - 2a \log(a+bx) - \frac{a^2}{a+bx} \right].$$

$$34. \int \frac{dx}{x(a+bx)} = -\frac{1}{a} \log \frac{a+bx}{x}.$$

$$35. \int \frac{dx}{x(a+bx)^2} = \frac{1}{a(a+bx)} - \frac{1}{a^2} \log \frac{a+bx}{x}.$$

$$36. \int \frac{dx}{x^2(a+bx)} = -\frac{1}{ax} + \frac{b}{a^2} \log \frac{a+bx}{x}.$$

B. — EXPRESSIONS INVOLVING  $(a+bx^2)$ .

$$37. \int \frac{dx}{c^2+x^2} = \frac{1}{c} \tan^{-1} \frac{x}{c}.$$

$$38. \int \frac{dx}{c^2-x^2} = \frac{1}{2c} \log \frac{c+x}{c-x}.$$

$$39. \int \frac{dx}{a+bx^2} = \frac{1}{\sqrt{ab}} \tan^{-1} x \sqrt{\frac{b}{a}}, \text{ if } a > 0, b > 0.$$

$$40. \int \frac{dx}{a+bx^2} = \frac{1}{2\sqrt{-ab}} \log \frac{\sqrt{a}+x\sqrt{-b}}{\sqrt{a}-x\sqrt{-b}}, \text{ if } a > 0, b < 0.$$

$$41. \int \frac{dx}{(a+bx^2)^2} = \frac{x}{2a(a+bx^2)} + \frac{1}{2a} \int \frac{dx}{a+bx^2}.$$

$$42. \int \frac{dx}{(a+bx^2)^{m+1}} = \frac{1}{2ma} \frac{x}{(a+bx^2)^m} + \frac{2m-1}{2ma} \int \frac{dx}{(a+bx^2)^m}.$$

$$43. \int \frac{x dx}{a+bx^2} = \frac{1}{2b} \log \left( x^2 + \frac{a}{b} \right).$$

$$44. \int \frac{x dx}{(a + bx^2)^{m+1}} = \frac{1}{2} \int \frac{dz}{(a + bz)^{m+1}}, \text{ where } z = x^2.$$

$$45. \int \frac{dx}{x(a + bx^2)} = \frac{1}{2a} \log \frac{x^2}{a + bx^2}.$$

$$46. \int \frac{x^2 dx}{a + bx^2} = \frac{x}{b} - \frac{a}{b} \int \frac{dx}{a + bx^2}.$$

$$47. \int \frac{dx}{x^2(a + bx^2)} = -\frac{1}{ax} - \frac{b}{a} \int \frac{dx}{a + bx^2}.$$

$$48. \int \frac{x^2 dx}{(a + bx^2)^{m+1}} = \frac{-x}{2mb(a + bx^2)^m} + \frac{1}{2mb} \int \frac{dx}{(a + bx^2)^m}.$$

$$49. \int \frac{dx}{x^2(a + bx^2)^{m+1}} = \frac{1}{a} \int \frac{dx}{x^2(a + bx^2)^m} - \frac{b}{a} \int \frac{dx}{(a + bx^2)^{m+1}}.$$

$$50. \int \frac{dx}{a + bx^3} = \frac{k}{3a} \left[ \frac{1}{2} \log \left( \frac{(k+x)^2}{k^2 - kx + x^2} \right) + \sqrt{3} \tan^{-1} \frac{2x - k}{k\sqrt{3}} \right],$$

where  $bk^3 = a$ .

$$51. \int \frac{x dx}{a + bx^3} = \frac{1}{3bk} \left[ \frac{1}{2} \log \left( \frac{k^2 - kx + x^2}{(k+x)^2} \right) + \sqrt{3} \tan^{-1} \frac{2x - k}{k\sqrt{3}} \right],$$

where  $bk^3 = a$ .

$$52. \int \frac{dx}{x(a + bx^n)} = \frac{1}{an} \log \frac{x^n}{a + bx^n}.$$

$$53. \int \frac{dx}{(a + bx^n)^{m+1}} = \frac{1}{a} \int \frac{dx}{(a + bx^n)^m} - \frac{b}{a} \int \frac{x^n dx}{(a + bx^n)^{m+1}}.$$

$$54. \int \frac{x^n dx}{(a + bx^n)^{r+1}} = \frac{1}{b} \int \frac{x^{n-r}}{(a + bx^n)^r} - \frac{a}{b} \int \frac{x^{n-n}}{(a + bx^n)^{r+1}}.$$

$$55. \int \frac{dx}{x^n(a + bx^n)^{r+1}} = \frac{1}{a} \int \frac{dx}{x^n(a + bx^n)^r} - \frac{b}{a} \int \frac{dx}{x^{n-n}(a + bx^n)^{r+1}}.$$

$$56. \int x^{m-1}(a+bx)^p dx = \begin{cases} \frac{1}{b(m+np)} \left[ x^{m-n}(a+bx)^{p+1} - (m-n)a \int x^{m-n-1}(a+bx)^p dx \right] \\ \frac{1}{m+np} \left[ x^m(a+bx)^p + npa \int x^{m-1}(a+bx)^{p-1} dx \right] \\ \frac{1}{ma} \left[ x^m(a+bx)^{p+1} - (m+np+n)b \int x^{m+n-1}(a+bx)^p dx \right] \\ \frac{1}{an(p+1)} \left[ -x^m(a+bx)^{p+1} + (m+np+n) \int x^{m-1}(a+bx)^{p+1} dx \right] \end{cases}$$

C. — EXPRESSIONS INVOLVING  $(a+bx+cx^2)$ .

Let  $X = a + bx + cx^2$  and  $q = 4ac - b^2$ , then

$$57. \int \frac{dx}{X} = \frac{2}{\sqrt{q}} \tan^{-1} \frac{2cx+b}{\sqrt{q}}, \text{ when } q > 0.$$

$$60. \int \frac{dx}{X^2} = \frac{2cx+b}{q} \left( \frac{1}{2X^2} + \frac{3c}{qX} \right) + \frac{6c^2}{q^2} \int \frac{dx}{X}$$

$$58. \int \frac{dx}{X} = \frac{1}{\sqrt{-q}} \log \frac{2cx+b-\sqrt{-q}}{2cx+b+\sqrt{-q}}, \text{ when } q < 0.$$

$$61. \int \frac{dx}{X^{n+1}} = \frac{2cx+b}{nqX^n} + \frac{2(2n-1)c}{qn} \int \frac{dx}{X^n}$$

$$59. \int \frac{dx}{X^2} = \frac{2cx+b}{qX} + \frac{2c}{q} \int \frac{dx}{X}$$

$$62. \int \frac{x dx}{X} = \frac{1}{2c} \log X - \frac{b}{2c} \int \frac{dx}{X}$$

$$63. \int \frac{x dx}{X^2} = -\frac{bx + 2a}{qX} - \frac{b}{q} \int \frac{dx}{X}.$$

$$64. \int \frac{x dx}{X^{n+1}} = -\frac{2a + bx}{nqX^n} - \frac{b(2n-1)}{nq} \int \frac{dx}{X^n}.$$

$$65. \int \frac{x^2 dx}{X} = \frac{x}{c} - \frac{b}{2c^2} \log X + \frac{b^2 - 2ac}{2c^2} \int \frac{dx}{X}.$$

$$66. \int \frac{x^2 dx}{X^2} = \frac{(b^2 - 2ac)x + ab}{cqX} + \frac{2a}{q} \int \frac{dx}{X}.$$

$$67. \int \frac{x^m dx}{X^{n+1}} = -\frac{x^{m-1}}{(2n-m+1)cX^n} - \frac{n-m+1}{2n-m+1} \cdot \frac{b}{c} \int \frac{x^{m-1} dx}{X^{n+1}} \\ + \frac{m-1}{2n-m+1} \cdot \frac{a}{c} \int \frac{x^{m-2} dx}{X^{n+1}}.$$

$$68. \int \frac{dx}{xX} = \frac{1}{2a} \log \frac{x^2}{X} - \frac{b}{2a} \int \frac{dx}{X}.$$

$$69. \int \frac{dx}{x^2 X} = \frac{b}{2a^2} \log \frac{X}{x^2} - \frac{1}{ax} + \left( \frac{b^2}{2a^2} - \frac{c}{a} \right) \int \frac{dx}{X}.$$

$$70. \int \frac{dx}{x^m X^{n+1}} = -\frac{1}{(m-1)ax^{m-1}X^n} - \frac{n+m-1}{m-1} \cdot \frac{b}{a} \int \frac{dx}{x^{m-1}X^{n+1}} \\ - \frac{2n+m-1}{m-1} \cdot \frac{c}{a} \int \frac{dx}{x^{m-2}X^{n+1}}.$$

## D. — RATIONAL FRACTIONS.

Every proper fraction can be represented by the general form:

$$\frac{f(x)}{F(x)} = \frac{g_1 x^{n-1} + g_2 x^{n-2} + g_3 x^{n-3} + \dots + g_n}{x^n + k_1 x^{n-1} + k_2 x^{n-2} + \dots + k_n}.$$

$a, b, c$ , etc., are the roots of the equation  $F(x) = 0$ , so that

$$F(x) = (x-a)^r (x-b)^s (x-c)^t \dots,$$

$$\begin{aligned} \text{then } \frac{f(x)}{F(x)} &= \frac{A_1}{(x-a)^p} + \frac{A_2}{(x-a)^{p-1}} + \frac{A_3}{(x-a)^{p-2}} + \dots + \frac{A_p}{x-a} \\ &+ \frac{B_1}{(x-b)^q} + \frac{B_2}{(x-b)^{q-1}} + \frac{B_3}{(x-b)^{q-2}} + \dots + \frac{B_q}{x-b} \\ &+ \frac{C_1}{(x-c)^r} + \frac{C_2}{(x-c)^{r-1}} + \frac{C_3}{(x-c)^{r-2}} + \dots + \frac{C_r}{x-c} \\ &+ \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \end{aligned}$$

Where the numerators of the separate fractions may be determined by the equations

$$A_m = \frac{\phi_1^{[m-1]}(a)}{(m-1)!}, \quad B_m = \frac{\phi_2^{[m-1]}(b)}{(m-1)!}, \quad \text{etc., etc.}$$

$$\phi_1(x) = \frac{f(x)(x-a)^p}{F(x)}, \quad \phi_2(x) = \frac{f(x)(x-b)^q}{F(x)}, \quad \text{etc., etc.}$$

If  $a, b, c, \dots$ , are single roots, then  $p = q = r = \dots = 1$ , and

$$\frac{f(x)}{F(x)} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c} \dots$$

where  $A = \frac{f(a)}{F'(a)}, B = \frac{f(b)}{F'(b)}, \text{ etc.}$

The simpler fractions, into which the original fraction is thus divided, may be integrated by means of the following formulas:

$$71. \int \frac{h dx}{(mx+n)^l} = \int \frac{h d(mx+n)}{m(mx+n)^l} = \frac{h}{m(1-l)(mx+n)^{l-1}}.$$

$$72. \int \frac{h dx}{mx+n} = \frac{h}{m} \log(mx+n).$$

If any of the roots of the equation  $f(x) = 0$  are imaginary, the parts of the integral which arise from conjugate roots can be combined together and the integral brought into a real form. The following formula, in which  $i = \sqrt{-1}$ , is often useful in combining logarithms of conjugate complex quantities:

$$73. \log(x \pm yi) = \frac{1}{2} \log(x^2 + y^2) \pm i \tan^{-1} \frac{y}{x}$$

## III. IRRATIONAL ALGEBRAIC FUNCTIONS.

A.—EXPRESSIONS INVOLVING  $\sqrt{a+bx}$ .

The substitution of a new variable of integration,  
 $y = \sqrt{a+bx}$ , gives

$$74. \int \sqrt{a+bx} \, dx = \frac{2}{3b} \sqrt{(a+bx)^3}.$$

$$75. \int x \sqrt{a+bx} \, dx = -\frac{2(2a-3bx)\sqrt{(a+bx)^3}}{15b^2}.$$

$$76. \int x^2 \sqrt{a+bx} \, dx = \frac{2(8a^2-12abx+15b^2x^2)\sqrt{(a+bx)^3}}{105b^3}.$$

$$77. \int \frac{\sqrt{a+bx}}{x} \, dx = 2\sqrt{a+bx} + a \int \frac{dx}{x\sqrt{a+bx}}.$$

$$78. \int \frac{dx}{\sqrt{a+bx}} = \frac{2\sqrt{a+bx}}{b}.$$

$$79. \int \frac{x \, dx}{\sqrt{a+bx}} = -\frac{2(2a-bx)\sqrt{a+bx}}{3b^2}.$$

$$80. \int \frac{x^2 \, dx}{\sqrt{a+bx}} = \frac{2(8a^2-4abx+3b^2x^2)\sqrt{a+bx}}{15b^3}.$$

$$81. \int \frac{dx}{x\sqrt{a+bx}} = \frac{1}{\sqrt{a}} \log \left( \frac{\sqrt{a+bx} - \sqrt{a}}{\sqrt{a+bx} + \sqrt{a}} \right), \text{ for } a > 0.$$

$$82. \int \frac{dx}{x\sqrt{a+bx}} = \frac{2}{\sqrt{-a}} \tan^{-1} \sqrt{\frac{a+bx}{-a}}, \text{ for } a < 0.$$

$$83. \int \frac{dx}{x^2\sqrt{a+bx}} = -\frac{\sqrt{a+bx}}{ax} - \frac{b}{2a} \int \frac{dx}{x\sqrt{a+bx}}.$$

84.  $\int (a+bx)^{\pm \frac{1}{2}} dx = \frac{2}{b} \int y^{\pm 1} dy = \frac{2(a+bx)^{\frac{3 \pm 1}{2}}}{b(2 \pm 1)}$
85.  $\int x(a+bx)^{\pm \frac{1}{2}} dx = \frac{2}{b^2} \left[ \frac{(a+bx)^{\frac{3 \pm 1}{2}}}{4 \pm 1} - \frac{a(a+bx)^{\frac{3 \pm 1}{2}}}{2 \pm 1} \right]$
86.  $\int \frac{x^m dx}{\sqrt{a+bx}} = \frac{2x^m \sqrt{a+bx}}{(2m+1)b} - \frac{2ma}{(2m+1)b} \int \frac{x^{m-1} dx}{\sqrt{a+bx}}$
87.  $\int \frac{dx}{x^m \sqrt{a+bx}} = -\frac{\sqrt{a+bx}}{(n-1)ax^{n-1}} - \frac{(2n-3)b}{(2n-2)a} \int \frac{dx}{x^{n-1} \sqrt{a+bx}}$
88.  $\int \frac{(a+bx)^{\frac{n}{2}} dx}{x} = b \int (a+bx)^{\frac{n-2}{2}} dx + a \int \frac{(a+bx)^{\frac{n-2}{2}} dx}{x}$
89.  $\int \frac{dx}{x(a+bx)^{\frac{m}{2}}} = \frac{1}{a} \int \frac{dx}{x(a+bx)^{\frac{m-2}{2}}} - \frac{b}{a} \int \frac{dx}{(a+bx)^{\frac{m}{2}}}$

B. — EXPRESSIONS \* INVOLVING  $\sqrt{x^2 \pm a^2}$  AND  $\sqrt{a^2 - x^2}$ .

90.  $\int \sqrt{x^2 \pm a^2} dx = \frac{1}{2} [x \sqrt{x^2 \pm a^2} \pm a^2 \log(x + \sqrt{x^2 \pm a^2})]$
91.  $\int \sqrt{a^2 - x^2} dx = \frac{1}{2} (x \sqrt{a^2 - x^2} + a^2 \sin^{-1} \frac{x}{a})$
92.  $\int \frac{dx}{\sqrt{x^2 \pm a^2}} = \log(x + \sqrt{x^2 \pm a^2})$
93.  $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a}$
94.  $\int \frac{dx}{x \sqrt{x^2 - a^2}} = \frac{1}{a} \cos^{-1} \frac{a}{x}$
95.  $\int \frac{dx}{x \sqrt{a^2 \pm x^2}} = -\frac{1}{a} \log \left( \frac{a + \sqrt{a^2 \pm x^2}}{x} \right)$
96.  $\int \frac{\sqrt{a^2 \pm x^2}}{x} dx = \sqrt{a^2 \pm x^2} - a \log \frac{a + \sqrt{a^2 \pm x^2}}{x}$

\* These equations are all special cases of more general equations given in the next section.

$$97. \int \frac{\sqrt{x^2 - a^2}}{x} dx = \sqrt{x^2 - a^2} - a \cos^{-1} \frac{a}{x}.$$

$$98. \int \frac{x dx}{\sqrt{a^2 \pm x^2}} = \pm \sqrt{a^2 \pm x^2}.$$

$$99. \int \frac{x dx}{\sqrt{x^2 - a^2}} = \sqrt{x^2 - a^2}.$$

$$100. \int x \sqrt{x^2 \pm a^2} dx = \frac{1}{3} \sqrt{(x^2 \pm a^2)^3}.$$

$$101. \int x \sqrt{a^2 - x^2} dx = -\frac{1}{3} \sqrt{(a^2 - x^2)^3}.$$

$$102. \int \sqrt{(x^2 \pm a^2)^3} dx \\ = \frac{1}{4} \left[ x \sqrt{(x^2 \pm a^2)^3} \pm \frac{3a^2 x}{2} \sqrt{x^2 \pm a^2} + \frac{3a^4}{2} \log(x + \sqrt{x^2 \pm a^2}) \right].$$

$$103. \int \sqrt{(a^2 - x^2)^3} dx \\ = \frac{1}{4} \left[ x \sqrt{(a^2 - x^2)^3} + \frac{3a^2 x}{2} \sqrt{a^2 - x^2} + \frac{3a^4}{2} \sin^{-1} \frac{x}{a} \right].$$

$$104. \int \frac{dx}{\sqrt{(x^2 \pm a^2)^3}} = \frac{\pm x}{a^2 \sqrt{x^2 \pm a^2}}.$$

$$105. \int \frac{dx}{\sqrt{(a^2 - x^2)^3}} = \frac{x}{a^2 \sqrt{a^2 - x^2}}.$$

$$106. \int \frac{x dx}{\sqrt{(x^2 \pm a^2)^3}} = \frac{-1}{\sqrt{x^2 \pm a^2}}.$$

$$107. \int \frac{x dx}{\sqrt{(a^2 - x^2)^3}} = \frac{1}{\sqrt{a^2 - x^2}}.$$

$$108. \int x \sqrt{(x^2 \pm a^2)^3} dx = \frac{1}{5} \sqrt{(x^2 \pm a^2)^5}$$

$$109. \int x \sqrt{(a^2 - x^2)^3} dx = -\frac{1}{5} \sqrt{(a^2 - x^2)^5}$$



$$110. \int x^2 \sqrt{x^2 \pm a^2} dx \\ = \frac{x}{4} \sqrt{(x^2 \pm a^2)^3} \mp \frac{a^2}{8} (x \sqrt{x^2 \pm a^2} \pm a^2 \log (x + \sqrt{x^2 \pm a^2})).$$

$$111. \int x^2 \sqrt{a^2 - x^2} dx \\ = -\frac{x}{4} \sqrt{(a^2 - x^2)^3} + \frac{a^2}{8} \left( x \sqrt{a^2 - x^2} + a^2 \sin^{-1} \frac{x}{a} \right).$$

$$112. \int \frac{x^2 dx}{\sqrt{x^2 \pm a^2}} = \frac{x}{2} \sqrt{x^2 \pm a^2} \mp \frac{a^2}{2} \log (x + \sqrt{x^2 \pm a^2}).$$

$$113. \int \frac{x^2 dx}{\sqrt{a^2 - x^2}} = -\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}.$$

$$114. \int \frac{dx}{x^2 \sqrt{x^2 \pm a^2}} = \pm \frac{\sqrt{x^2 \pm a^2}}{a^2 x}.$$

$$115. \int \frac{dx}{x^2 \sqrt{a^2 - x^2}} = -\frac{\sqrt{a^2 - x^2}}{a^2 x}.$$

$$116. \int \frac{\sqrt{x^2 \pm a^2} dx}{x^2} = -\frac{\sqrt{x^2 \pm a^2}}{x} + \log x + \sqrt{x^2 \pm a^2}.$$

$$117. \int \frac{\sqrt{a^2 - x^2} dx}{x^2} = -\frac{\sqrt{a^2 - x^2}}{x} - \sin^{-1} \frac{x}{a}.$$

$$118. \int \frac{x^2 dx}{\sqrt{(x^2 \pm a^2)^3}} = \frac{-x}{\sqrt{x^2 \pm a^2}} + \log (x + \sqrt{x^2 \pm a^2}).$$

$$119. \int \frac{x^2 dx}{\sqrt{(a^2 - x^2)^3}} = \frac{x}{\sqrt{a^2 - x^2}} - \sin^{-1} \frac{x}{a}.$$

### C. — EXPRESSIONS INVOLVING $\sqrt{a + bx + cx^2}$ .

Let  $X = a + bx + cx^2$ ,  $q = 4ac - b^2$ , and  $k = \frac{4c}{q}$ . In order to rationalize the function  $f(x, \sqrt{a + bx + cx^2})$  we may put  $\sqrt{a + bx + cx^2} = \sqrt{\pm c} \sqrt{A + Bx \pm x^2}$ , according as  $c$  is positive or negative, and then substitute for  $x$  a new variable  $z$ , such that

$$z = \sqrt{A+Bx+x^2} - x, \text{ if } c > 0.$$

$$z = \frac{\sqrt{A+Bx-x^2} - \sqrt{A}}{x}, \text{ if } c < 0 \text{ and } \frac{a}{-c} > 0.$$

$$z = \sqrt{\frac{x-\beta}{\alpha-x}}, \text{ where } \alpha \text{ and } \beta \text{ are the roots of the equation } A+Bx-x^2=0, \text{ if } c < 0 \text{ and } \frac{a}{-c} < 0.$$

By rationalization, or by the aid of reduction formulas, may be obtained the values of the following integrals :

$$120. \int \frac{dx}{\sqrt{X}} = \frac{1}{\sqrt{c}} \log \left( \sqrt{X} + x\sqrt{c} + \frac{b}{2\sqrt{c}} \right), \text{ if } c > 0.$$

$$121. \int \frac{dx}{\sqrt{X}} = \frac{1}{\sqrt{-c}} \sin^{-1} \left( \frac{-2cx-b}{\sqrt{b^2-4ac}} \right), \text{ if } c < 0.$$

$$122. \int \frac{dx}{X\sqrt{X}} = \frac{2(2cx+b)}{q\sqrt{X}}.$$

$$123. \int \frac{dx}{X^2\sqrt{X}} = \frac{2(2cx+b)}{3q\sqrt{X}} \left( \frac{1}{X} + 2k \right).$$

$$124. \int \frac{dx}{X^n\sqrt{X}} = \frac{2(2cx+b\sqrt{X})}{(2n-1)qX^n} + \frac{2k(n-1)}{2n-1} \int \frac{dx}{X^{n-1}\sqrt{X}}.$$

$$125. \int \sqrt{X} dx = \frac{(2cx+b)\sqrt{X}}{4c} + \frac{1}{2k} \int \frac{dx}{\sqrt{X}}.$$

$$126. \int X\sqrt{X} dx = \frac{(2cx+b)\sqrt{X}}{8c} \left( X + \frac{3}{2k} \right) + \frac{3}{8k^2} \int \frac{dx}{\sqrt{X}}.$$

$$127. \int X^2\sqrt{X} dx = \frac{(2cx+b)\sqrt{X}}{12c} \left( X^2 + \frac{5X}{4k} + \frac{15}{8k^2} \right) + \frac{5}{16k^2} \int \frac{dx}{\sqrt{X}}.$$

$$128. \int X^n\sqrt{X} dx = \frac{(2cx+b)X^n\sqrt{X}}{4(n+1)c} + \frac{2n+1}{2(n+1)k} \int \frac{X^n dx}{\sqrt{X}}.$$

$$129. \int \frac{x dx}{\sqrt{X}} = \frac{\sqrt{X}}{c} - \frac{b}{2c} \int \frac{dx}{\sqrt{X}}.$$

$$130. \int \frac{x dx}{X\sqrt{X}} = -\frac{2(bx+2a)}{q\sqrt{X}}$$

$$131. \int \frac{x dx}{X^n\sqrt{X}} = -\frac{\sqrt{X}}{(2n-1)cX^n} - \frac{b}{2c} \int \frac{dx}{X^n\sqrt{X}}$$

$$132. \int \frac{x^2 dx}{\sqrt{X}} = \left(\frac{x}{2c} - \frac{3b}{4c^2}\right)\sqrt{X} + \frac{3b^2-4ac}{8c^2} \int \frac{dx}{\sqrt{X}}$$

$$133. \int \frac{x^2 dx}{X\sqrt{X}} = \frac{(2b^2-4ac)x+2ab}{cq\sqrt{X}} + \frac{1}{c} \int \frac{dx}{\sqrt{X}}$$

$$134. \int \frac{x^2 dx}{X^n\sqrt{X}} = \frac{(2b^2-4ac)x+2ab}{(2n-1)cqX^{n-1}\sqrt{X}} + \frac{4ac+(2n-3)b^2}{(2n-1)cq} \int \frac{dx}{X^{n-1}\sqrt{X}}$$

$$135. \int \frac{x^2 dx}{\sqrt{X}} = \left(\frac{x^2}{3c} - \frac{5bx}{12c^2} + \frac{5b^2}{8c^2} - \frac{2a}{3c^2}\right)\sqrt{X} + \left(\frac{3ab}{4c^2} - \frac{5b^2}{16c^2}\right) \int \frac{dx}{\sqrt{X}}$$

$$136. \int x\sqrt{X} dx = \frac{X\sqrt{X}}{3c} - \frac{b}{2c} \int \sqrt{X} dx.$$

$$137. \int xX\sqrt{X} dx = \frac{X^2\sqrt{X}}{5c} - \frac{b}{2c} \int X\sqrt{X} dx.$$

$$138. \int \frac{xX^n dx}{\sqrt{X}} = \frac{X^n\sqrt{X}}{(2n+1)c} - \frac{b}{2c} \int \frac{X^n dx}{\sqrt{X}}$$

$$139. \int x^2\sqrt{X} dx = \left(x - \frac{5b}{6c}\right)\frac{X\sqrt{X}}{4c} + \frac{5b^2-4ac}{16c^2} \int \sqrt{X} dx$$

$$140. \int \frac{x^2X^n dx}{\sqrt{X}} = \frac{xX^n\sqrt{X}}{2(n+1)c} - \frac{(2n+3)b}{4(n+1)c} \int \frac{xX^n dx}{\sqrt{X}} \\ - \frac{a}{2(n+1)c} \int \frac{X^n dx}{\sqrt{X}}$$

$$141. \int x^3\sqrt{X} dx = \left(x^2 - \frac{7bx}{8c} + \frac{35b^2}{48c^2} - \frac{2a}{3c}\right)\frac{X\sqrt{X}}{5c} \\ + \left(\frac{3ab}{8c^2} - \frac{7b^2}{32c^2}\right) \int \sqrt{X} dx.$$

$$142. \int \frac{dx}{x\sqrt{X}} = -\frac{1}{\sqrt{a}} \log \left( \frac{\sqrt{X} + \sqrt{a}}{x} + \frac{b}{2\sqrt{a}} \right), \text{ if } a > 0.$$

$$143. \int \frac{dx}{x\sqrt{X}} = \frac{1}{\sqrt{-a}} \sin^{-1} \left( \frac{bx + 2a}{x\sqrt{b^2 - 4ac}} \right), \text{ if } a < 0.$$

$$144. \int \frac{dx}{x\sqrt{X}} = -\frac{2\sqrt{X}}{bx}, \text{ if } a = 0.$$

$$145. \int \frac{dx}{xX^n\sqrt{X}} = \frac{\sqrt{X}}{(2n-1)aX^n} + \frac{1}{a} \int \frac{dx}{xX^{n-1}\sqrt{X}} - \frac{b}{2a} \int \frac{dx}{X^n\sqrt{X}}.$$

$$146. \int \frac{dx}{x^2\sqrt{X}} = -\frac{\sqrt{X}}{ax} - \frac{b}{2a} \int \frac{dx}{x\sqrt{X}}.$$

$$147. \int \frac{\sqrt{X} dx}{x} = \sqrt{X} + \frac{b}{2} \int \frac{dx}{\sqrt{X}} + a \int \frac{dx}{x\sqrt{X}}.$$

$$148. \int \frac{X^n dx}{x\sqrt{X}} = \frac{X^n}{(2n-1)\sqrt{X}} + a \int \frac{X^{n-1} dx}{x\sqrt{X}} + \frac{b}{2} \int \frac{X^{n-1} dx}{\sqrt{X}}.$$

$$149. \int \frac{\sqrt{X} dx}{x^2} = -\frac{\sqrt{X}}{x} + \frac{b}{2} \int \frac{dx}{x\sqrt{X}} + c \int \frac{dx}{\sqrt{X}}.$$

$$150. \int \frac{x^m dx}{X^n\sqrt{X}} = \frac{1}{c} \int \frac{x^{m-2} dx}{X^{n-1}\sqrt{X}} - \frac{b}{c} \int \frac{x^{m-1} dx}{X^n\sqrt{X}} - \frac{a}{c} \int \frac{x^{m-2} dx}{X^n\sqrt{X}}.$$

$$151. \int \frac{x^m X^n dx}{\sqrt{X}} = \frac{x^{m-1} X^n \sqrt{X}}{(2n+m)c} - \frac{(2n+2m-1)b}{2c(2n+m)} \int \frac{x^{m-1} X^n dx}{\sqrt{X}} \\ - \frac{(m-1)a}{(2n+m)c} \int \frac{x^{m-2} X^n dx}{\sqrt{X}}.$$

$$152. \int \frac{dx}{x^m X^n \sqrt{X}} = -\frac{\sqrt{X}}{(m-1)ax^{m-1}X^n} \\ - \frac{(2n+2m-3)b}{2a(m-1)} \int \frac{dx}{x^{m-1}X^n\sqrt{X}} - \frac{(2n+m-2)c}{(m-1)a} \int \frac{dx}{x^{m-2}X^n\sqrt{X}}$$

$$153. \int \frac{X^n dx}{x^m \sqrt{X}} = -\frac{X^{n-1} \sqrt{X}}{(m-1)x^{m-1}} + \frac{(2n-1)b}{2(m-1)} \int \frac{X^{n-1} dx}{x^{m-1} \sqrt{X}} \\ + \frac{(2n-1)c}{m-1} \int \frac{X^{n-1} dx}{x^{m-2} \sqrt{X}}.$$

## D. — MISCELLANEOUS EXPRESSIONS.

$$154. \int \sqrt{2ax - x^2} dx = \frac{x-a}{2} \sqrt{2ax - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x-a}{a}.$$

$$155. \int \frac{dx}{\sqrt{2ax - x^2}} = \operatorname{versin}^{-1} \frac{x}{a}.$$

$$156. \int \frac{dx}{(x+1)\sqrt{x^2-1}} = +\sqrt{\frac{x-1}{x+1}}.$$

$$157. \int \frac{dx}{(x-1)\sqrt{x^2-1}} = -\sqrt{\frac{x+1}{x-1}}.$$

$$158. \int \sqrt{\frac{1+x}{1-x}} dx = \sin^{-1} x - \sqrt{1-x^2}.$$

$$159. \int \sqrt{\frac{x+a}{x+b}} dx = \sqrt{(x+a)(x+b)} \\ + (a-b) \log(\sqrt{x+a} + \sqrt{x+b}).$$

$$160. \int \frac{dx}{\sqrt{(x-a)(\beta-x)}} = 2 \sin^{-1} \sqrt{\frac{x-a}{\beta-a}}.$$

$$161. \int \frac{dx}{\sqrt{(a+bx)(a-\beta x)}} = \frac{2}{\sqrt{b\beta}} \sin^{-1} \sqrt{\frac{\beta(a+bx)}{a\beta+ba}}.$$

$$162. \int \sqrt[3]{a+bx} dx = \frac{3}{4b} \sqrt[3]{(a+bx)^4}.$$

$$163. \int \frac{dx}{\sqrt[3]{a+bx}} = \frac{3}{2b} \sqrt[3]{(a+bx)^2}.$$

$$164. \int \frac{x dx}{\sqrt[3]{a+bx}} = -\frac{3(3a-2bx)}{10b^2} \sqrt[3]{(a+bx)^2}.$$

$$165. \int \frac{dx}{x\sqrt{x^n-a^2}} = \frac{2}{an} \sec^{-1}\left(\frac{x^{\frac{n}{2}}}{a}\right).$$

$$166. \int \frac{dx}{x\sqrt{x^n+a^2}} = \frac{1}{an} \log \frac{\sqrt{a^2+x^n}-a}{\sqrt{a^2+x^n}+a}.$$

## IV. TRANSCENDENTAL FUNCTIONS.

$$167. \int \sin x dx = -\cos x.$$

$$168. \int \sin^2 x dx = -\frac{1}{2} \cos x \sin x + \frac{1}{2} x.$$

$$169. \int \sin^3 x dx = -\frac{1}{2} \cos x (\sin^2 x + 2).$$

$$170. \int \sin^n x dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x dx.$$

$$171. \int \cos x dx = \sin x.$$

$$172. \int \cos^2 x dx = \frac{1}{2} \sin x \cos x + \frac{1}{2} x.$$

$$173. \int \cos^3 x dx = \frac{1}{2} \sin x (\cos^2 x + 2).$$

$$174. \int \cos^n x dx = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} \int \cos^{n-2} x dx.$$

$$175. \int \sin x \cos x dx = \frac{1}{2} \sin^2 x.$$

$$176. \int \sin^2 x \cos^2 x dx = -\frac{1}{8} (\frac{1}{2} \sin 4x - x).$$

$$177. \int \sin x \cos^m x dx = -\frac{\cos^{m+1} x}{m+1}.$$

$$178. \int \sin^m x \cos x dx = \frac{\sin^{m+1} x}{m+1}.$$

$$179. \int \cos^m x \sin^n x dx = \frac{\cos^{m-1} x \sin^{n+1} x}{m+n} \\ + \frac{m-1}{m+n} \int \cos^{m-2} x \sin^n x dx.$$

$$180. \int \cos^m x \sin^n x dx = -\frac{\sin^{n-1} x \cos^{m+1} x}{m+n} \\ + \frac{n-1}{m+n} \int \cos^m x \sin^{n-2} x dx.$$

$$181. \int \frac{\cos^m x dx}{\sin^n x} = -\frac{\cos^{m+1} x}{(n-1)\sin^{n-1} x} - \frac{m-n+2}{n-1} \int \frac{\cos^m x dx}{\sin^{n-2} x}.$$

$$182. \int \frac{\cos^m x dx}{\sin^n x} = \frac{\cos^{m-1} x}{(m-n)\sin^{n-1} x} + \frac{m-1}{m-n} \int \frac{\cos^{m-2} x dx}{\sin^n x}.$$

$$183. \int \frac{\sin^m x dx}{\cos^n x} = -\int \frac{\cos^m\left(\frac{\pi}{2}-x\right) d\left(\frac{\pi}{2}-x\right)}{\sin^n\left(\frac{\pi}{2}-x\right)}.$$

$$184. \int \frac{dx}{\sin^m x \cos^n x}$$

$$= \frac{1}{n-1} \cdot \frac{1}{\sin^{n-1} x \cdot \cos^{n-1} x} + \frac{m+n-2}{n-1} \int \frac{dx}{\sin^m x \cdot \cos^{n-2} x}$$

$$= -\frac{1}{m-1} \cdot \frac{1}{\sin^{m-1} x \cdot \cos^{m-1} x} + \frac{m+n-2}{m-1} \int \frac{dx}{\sin^{m-2} x \cdot \cos^n x}.$$

$$185. \int \frac{dx}{\sin^m x} = -\frac{1}{m-1} \cdot \frac{\cos x}{\sin^{m-1} x} + \frac{m-2}{m-1} \int \frac{dx}{\sin^{m-2} x}.$$

$$186. \int \frac{dx}{\cos^n x} = \frac{1}{n-1} \cdot \frac{\sin x}{\cos^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\cos^{n-2} x}.$$

$$187. \int \tan x dx = -\log \cos x.$$

$$188. \int \tan^2 x dx = \tan x - x.$$

$$189. \int \tan^n x dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x dx.$$

$$190. \int \operatorname{ctn} x dx = \log \sin x.$$

$$191. \int \operatorname{ctn}^2 x dx = -\operatorname{ctn} x - x.$$

$$192. \int \operatorname{ctn}^n x dx = -\frac{\operatorname{ctn}^{n-1} x}{n-1} - \int \operatorname{ctn}^{n-2} x dx.$$

$$193. \int \sec x dx = \log \tan\left(\frac{\pi}{4} + \frac{x}{2}\right).$$

$$194. \int \sec^2 x dx = \tan x.$$



$$195. \int \sec^n x dx = \int \frac{dx}{\cos^n x}.$$

$$196. \int \csc x dx = \log \tan \frac{1}{2} x.$$

$$197. \int \csc^2 x dx = -\cot x.$$

$$198. \int \csc^n x dx = \int \frac{dx}{\sin^n x}.$$

$$199. \int \frac{dx}{a + b \cos x} = \frac{-1}{\sqrt{a^2 - b^2}} \cdot \sin^{-1} \left[ \frac{b + a \cos x}{a + b \cos x} \right],$$

$$\text{or } \frac{1}{\sqrt{b^2 - a^2}} \log \left[ \frac{b + a \cos x + \sqrt{b^2 - a^2} \cdot \sin x}{a + b \cos x} \right].$$

$$200. \int \frac{dx}{a + b \cos x + c \sin x}$$

$$= \frac{-1}{\sqrt{a^2 - b^2 - c^2}} \cdot \sin^{-1} \left[ \frac{b^2 + c^2 + a(b \cos x + c \sin x)}{\sqrt{b^2 + c^2}(a + b \cos x + c \sin x)} \right]$$

$$= \frac{1}{\sqrt{b^2 + c^2 - a^2}} \cdot \log$$

$$\left[ \frac{b^2 + c^2 + a(b \cos x + c \sin x) + \sqrt{b^2 + c^2 - a^2}(b \sin x - c \cos x)}{\sqrt{b^2 + c^2}(a + b \cos x + c \sin x)} \right].$$

$$201. \int x \sin x dx = \sin x - x \cos x.$$

$$202. \int x^2 \sin x dx = 2x \sin x - (x^2 - 2) \cos x.$$

$$203. \int x^3 \sin x dx = (3x^2 - 6) \sin x - (x^3 - 6x) \cos x.$$

$$204. \int x^m \sin x dx = -x^m \cos x + m \int x^{m-1} \cos x dx.$$

$$205. \int x \cos x dx = \cos x + x \sin x.$$

$$206. \int x^2 \cos x dx = 2x \cos x + (x^2 - 2) \sin x.$$

$$207. \int x^3 \cos x dx = (3x^2 - 6) \cos x + (x^3 - 6x) \sin x.$$

208.  $\int x^m \cos x dx = x^m \sin x - m \int x^{m-1} \sin x dx.$
209.  $\int \frac{\sin x}{x^m} dx = -\frac{1}{m-1} \cdot \frac{\sin x}{x^{m-1}} + \frac{1}{m-1} \int \frac{\cos x}{x^{m-1}} dx.$
210.  $\int \frac{\cos x}{x^m} dx = -\frac{1}{m-1} \cdot \frac{\cos x}{x^{m-1}} - \frac{1}{m-1} \int \frac{\sin x}{x^{m-1}} dx.$
211.  $\int \frac{\sin x}{x} dx = x - \frac{x^3}{3 \cdot 3!} + \frac{x^5}{5 \cdot 5!} - \frac{x^7}{7 \cdot 7!} + \frac{x^9}{9 \cdot 9!} \dots$
212.  $\int \frac{\cos x}{x} dx = \log x - \frac{x^2}{2 \cdot 2!} + \frac{x^4}{4 \cdot 4!} - \frac{x^6}{6 \cdot 6!} + \frac{x^8}{8 \cdot 8!} \dots$
213.  $\int \sin mx \sin nx dx = \frac{\sin(m-n)x}{2(m-n)} - \frac{\sin(m+n)x}{2(m+n)}$
214.  $\int \cos mx \cos nx dx = \frac{\sin(m-n)x}{2(m-n)} + \frac{\sin(m+n)x}{2(m+n)}$
215.  $\int \sin^{-1} x dx = x \sin^{-1} x + \sqrt{1-x^2}.$
216.  $\int \cos^{-1} x dx = x \cos^{-1} x - \sqrt{1-x^2}.$
217.  $\int \tan^{-1} x dx = x \tan^{-1} x - \frac{1}{2} \log(1+x^2).$
218.  $\int \text{ctn}^{-1} x dx = x \text{ctn}^{-1} x + \frac{1}{2} \log(1+x^2).$
219.  $\int \text{versin}^{-1} x dx = (x-1) \text{versin}^{-1} x + \sqrt{2x-x^3}.$
220.  $\int (\sin^{-1} x)^2 dx = x (\sin^{-1} x)^2 - 2x + 2\sqrt{1-x^2} \sin^{-1} x.$
221.  $\int x \cdot \sin^{-1} x dx = \frac{1}{4} [(2x^2-1) \sin^{-1} x + x\sqrt{1-x^2}].$
222.  $\int x^n \sin^{-1} x dx = \frac{x^{n+1} \sin^{-1} x}{n+1} - \frac{1}{n+1} \int \frac{x^{n+1} dx}{\sqrt{1-x^2}}.$
223.  $\int x^n \cos^{-1} x dx = \frac{x^{n+1} \cos^{-1} x}{n+1} + \frac{1}{n+1} \int \frac{x^{n+1} dx}{\sqrt{1-x^2}}.$
224.  $\int x^n \tan^{-1} x dx = \frac{x^{n+1} \tan^{-1} x}{n+1} - \frac{1}{n+1} \int \frac{x^{n+1} dx}{1+x^2}.$

$$225. \int \log x dx = x \log x - x.$$

$$226. \int \frac{(\log x)^n}{x} dx = \frac{1}{n+1} (\log x)^{n+1}.$$

$$227. \int \frac{dx}{x \log x} = \log \cdot \log x.$$

$$228. \int \frac{dx}{x (\log x)^n} = -\frac{1}{(n-1) (\log x)^{n-1}}.$$

$$229. \int x^m \log x dx = x^{m+1} \left[ \frac{\log x}{m+1} - \frac{1}{(m+1)^2} \right]$$

$$230. \int e^{ax} dx = \frac{e^{ax}}{a}.$$

$$231. \int x e^{ax} dx = \frac{e^{ax}}{a^2} (ax - 1).$$

$$232. \int x^m e^{ax} dx = \frac{x^m e^{ax}}{a} - \frac{m}{a} \int x^{m-1} e^{ax} dx.$$

$$233. \int \frac{e^{ax}}{x^m} dx = -\frac{1}{m-1} \frac{e^{ax}}{x^{m-1}} + \frac{a}{m-1} \int \frac{e^{ax}}{x^{m-1}} dx.$$

$$234. \int e^{ax} \log x dx = \frac{e^{ax} \log x}{a} - \frac{1}{a} \int \frac{e^{ax}}{x} dx.$$

$$235. \int e^{ax} \sin x dx = \frac{e^{ax} (a \sin x - \cos x)}{a^2 + 1}.$$

$$236. \int e^{ax} \cos x dx = \frac{e^{ax} (a \cos x + \sin x)}{a^2 + 1}.$$

## DEFINITE INTEGRALS.

$$237. \int_0^{\infty} \frac{a dx}{a^2 + x^2} = \frac{\pi}{2}, \text{ if } a > 0; 0, \text{ if } a = 0; -\frac{\pi}{2}, \text{ if } a < 0.$$

$$238. \int_0^{\infty} x^{n-1} e^{-x} dx = \int_0^1 \left[ \log \frac{1}{x} \right]^{n-1} dx = \Gamma(n).$$

$$\Gamma(n+1) = n \cdot \Gamma(n).$$

$$\Gamma(2) = \Gamma(1) = 1.$$

$$\Gamma(n+1) = n!, \text{ if } n \text{ is an integer} \quad \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$

$$239. \int_0^1 x^{m-1} (1-x)^{n-1} dx = \int_0^{\infty} \frac{x^{m+1} dx}{(1+x)^{m+n}} = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}.$$

$$240. \int_0^{\frac{\pi}{2}} \sin^n x dx = \int_0^{\frac{\pi}{2}} \cos^n x dx$$

$$= \frac{1 \cdot 3 \cdot 5 \dots (n-1)}{2 \cdot 4 \cdot 6 \dots (n)} \cdot \frac{\pi}{2}, \text{ if } n \text{ is an even integer.}$$

$$= \frac{2 \cdot 4 \cdot 6 \dots (n-1)}{1 \cdot 3 \cdot 5 \cdot 7 \dots n}, \text{ if } n \text{ is an odd integer.}$$

$$= \frac{1}{2} \sqrt{\pi} \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2}+1\right)}, \text{ for any value of } n.$$

$$241. \int_0^{\infty} \frac{\sin mx dx}{x} = \frac{\pi}{2}, \text{ if } m > 0; 0, \text{ if } m = 0; -\frac{\pi}{2}, \text{ if } m < 0.$$

$$242. \int_0^{\infty} \frac{\sin x \cdot \cos mx dx}{x} = 0, \text{ if } m < -1 \text{ or } m > 1;$$

$$\frac{\pi}{4}, \text{ if } m = -1 \text{ or } m = 1; \frac{\pi}{2}, \text{ if } -1 < m < 1.$$

$$243. \int_0^{\infty} \frac{\sin^2 x dx}{x^2} = \frac{\pi}{2}.$$

$$244. \int_0^{\infty} \cos(x^2) dx = \int_0^{\infty} \sin(x^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2}}.$$

$$245. \int_0^{\infty} \frac{\cos mx dx}{1+x^2} = \frac{\pi}{2} \cdot e^{-m}.$$

$$246. \int_0^{\infty} \frac{\cos x dx}{\sqrt{x}} = \int_0^{\infty} \frac{\sin x dx}{\sqrt{x}} = \sqrt{\frac{\pi}{2}}.$$

$$247. \int_0^{\frac{\pi}{2}} \frac{dx}{\sqrt{1-k^2 \sin^2 x}}$$

$$= \frac{\pi}{2} \left[ 1 + \left(\frac{1}{2}\right)^2 k^2 + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 k^4 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 k^6 + \dots \right], \text{ if } k^2 < 1$$

$$= K.$$

$$248. \int_0^{\frac{\pi}{2}} \sqrt{1-k^2 \sin^2 x} \cdot dx$$

$$= \frac{\pi}{2} \left[ 1 - \left(\frac{1}{2}\right)^2 k^2 - \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \frac{k^4}{3} - \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 \frac{k^6}{5} - \dots \right], \text{ if } k^2 < 1.$$

$$= E.$$

$$249. \int_0^{\infty} e^{-ax^2} dx = \frac{1}{2a} \sqrt{\pi} = \frac{1}{2a} \Gamma\left(\frac{1}{2}\right).$$

$$250. \int_0^{\infty} x^n e^{-ax} dx = \frac{\Gamma(n+1)}{a^{n+1}} = \frac{n!}{a^{n+1}}.$$

$$251. \int_0^{\infty} x^{2n} e^{-ax^2} dx = \frac{1 \cdot 3 \cdot 5 \dots (2n-1)}{2^{n+1} a^n} \sqrt{\frac{\pi}{a}}.$$

$$252. \int_0^{\infty} e^{-x^2 - \frac{a^2}{x^2}} dx = \frac{e^{-2a} \sqrt{\pi}}{2}.$$

$$253. \int_0^{\infty} e^{-ax} \cos mx dx = \frac{a}{a^2 + m^2}, \text{ if } a > 0.$$

$$254. \int_0^{\infty} e^{-ax} \sin mx dx = \frac{m}{a^2 + m^2}, \text{ if } a > 0.$$

$$255. \int_0^{\infty} e^{-ax^2} \cos bx dx = \frac{\sqrt{\pi} \cdot e^{-\frac{b^2}{4a^2}}}{2a}.$$

$$256. \int_0^1 \frac{\log x}{1-x} dx = -\frac{\pi^2}{6}.$$

$$257. \int_0^1 \frac{\log x}{1+x} dx = -\frac{\pi^2}{12}.$$

$$258. \int_0^1 \frac{\log x}{1-x^2} dx = -\frac{\pi^2}{8}.$$

$$259. \int_0^1 \log\left(\frac{1+x}{1-x}\right) \cdot \frac{dx}{x} = \frac{\pi^2}{4}.$$

$$260. \int_0^\infty \log\left(\frac{e^x+1}{e^x-1}\right) dx = \frac{\pi^2}{4}.$$

$$261. \int_0^1 \frac{dx}{\sqrt{\log\left(\frac{1}{x}\right)}} = \sqrt{\pi}.$$

$$262. \int_0^1 x^m \log\left(\frac{1}{x}\right)^n dx = \frac{\Gamma(n+1)}{(m+1)^{n+1}}.$$

$$263. \int_0^{\frac{\pi}{2}} \log \sin x dx = \int_0^{\frac{\pi}{2}} \log \cos x dx = -\frac{\pi}{2} \cdot \log 2.$$

$$264. \int_0^\pi x \cdot \log \sin x dx = -\frac{\pi^2}{2} \log 2.$$

## AUXILIARY FORMULAS.

The following formulas are sometimes useful in the reduction of integrals:

$$265. \quad \log u = \log cu + \text{a constant.}$$

$$266. \quad \log(-u) = \log u + \text{a constant.}$$

$$267. \quad \sin^{-1}u = \begin{cases} -\sin^{-1}\sqrt{1-u^2} & + \text{a constant.} \\ -\frac{1}{2}\sin^{-1}(2u^2-1) & + \text{a constant.} \\ \frac{1}{2}\sin^{-1}2u\sqrt{1-u^2} & + \text{a constant.} \end{cases}$$

$$268. \quad \tan^{-1}u = \begin{cases} -\tan^{-1}\frac{1}{u} & + \text{a constant.} \\ \tan^{-1}\frac{u+c}{1-cu} & + \text{a constant.} \end{cases}$$

$$269. \quad \log(x \pm yi) = \frac{1}{2}\log(x^2 + y^2) \pm i\tan^{-1}\frac{y}{x}.$$

$$270. \quad \sin^{-1}u = \cos^{-1}\sqrt{1-u^2} = \tan^{-1}\frac{u}{\sqrt{1-u^2}} = \csc^{-1}\frac{1}{u}.$$

$$271. \quad \cos^{-1}u = \sin^{-1}\sqrt{1-u^2} = \tan^{-1}\sqrt{\frac{1}{u^2}-1} = \sec^{-1}\frac{1}{u}.$$

$$272. \quad \tan^{-1}x \pm \tan^{-1}y = \tan^{-1}\left(\frac{x \pm y}{1 \mp xy}\right).$$

$$273. \quad \sin^{-1}x \pm \sin^{-1}y = \sin^{-1}(x\sqrt{1-y^2} \pm y\sqrt{1-x^2}).$$

$$274. \quad \cos^{-1}x \pm \cos^{-1}y = \cos^{-1}(xy \mp \sqrt{(1-x^2)(1-y^2)}).$$

$$275. \quad \sin x = \frac{e^{xi} - e^{-xi}}{2i}.$$

$$276. \quad \cos x = \frac{e^{xi} + e^{-xi}}{2}.$$

$$277. \quad \sin xi = \frac{1}{2}i(e^x - e^{-x}) = i\sinh x.$$

$$278. \quad \cos xi = \frac{1}{2}(e^x + e^{-x}) = \cosh x.$$

$$279. \quad \log_e x = (2.3025851) \log_{10} x.$$

The Natural Logarithms of Numbers between 1.0 and 9.9.

N.	0	1	2	3	4	5	6	7	8	9
1.	0.000	0.095	0.182	0.262	0.336	0.405	0.470	0.531	0.588	0.642
2.	0.693	0.742	0.788	0.833	0.875	0.916	0.956	0.993	1.030	1.065
3.	1.099	1.131	1.163	1.194	1.224	1.253	1.281	1.308	1.335	1.361
4.	1.386	1.411	1.435	1.459	1.482	1.504	1.526	1.548	1.569	1.589
5.	1.609	1.629	1.649	1.668	1.686	1.705	1.723	1.740	1.758	1.775
6.	1.792	1.808	1.825	1.841	1.856	1.872	1.887	1.902	1.917	1.932
7.	1.946	1.960	1.974	1.988	2.001	2.015	2.028	2.041	2.054	2.067
8.	2.079	2.092	2.104	2.116	2.128	2.140	2.152	2.163	2.175	2.186
9.	2.197	2.208	2.219	2.230	2.241	2.251	2.262	2.272	2.282	2.293

The Natural Logarithms of Whole Numbers from 10 to 100.

N.	0	1	2	3	4	5	6	7	8	9
1	2.303	2.398	2.485	2.565	2.639	2.708	2.773	2.833	2.890	2.944
2	2.996	3.045	3.091	3.135	3.178	3.219	3.258	3.296	3.332	3.367
3	3.401	3.434	3.466	3.497	3.526	3.555	3.584	3.611	3.638	3.664
4	3.689	3.714	3.738	3.761	3.784	3.807	3.829	3.850	3.871	3.892
5	3.912	3.932	3.951	3.970	3.989	4.007	4.025	4.043	4.060	4.078
6	4.094	4.111	4.127	4.143	4.159	4.174	4.190	4.205	4.220	4.234
7	4.248	4.263	4.277	4.290	4.304	4.317	4.331	4.344	4.357	4.369
8	4.382	4.394	4.407	4.419	4.431	4.443	4.454	4.466	4.477	4.489
9	4.500	4.511	4.522	4.533	4.543	4.554	4.564	4.575	4.585	4.595
10	4.605	4.615	4.625	4.635	4.644	4.654	4.663	4.673	4.682	4.691

The Values in Circular Measure of Angles which are given in Degrees and Minutes.

1'	0.0003	9'	0.0026	3°	0.0524	20°	0.3491	100°	1.7453
2'	0.0006	10'	0.0029	4°	0.0698	30°	0.5236	110°	1.9199
3'	0.0009	20'	0.0058	5°	0.0873	40°	0.6981	120°	2.0944
4'	0.0012	30'	0.0087	6°	0.1047	50°	0.8727	130°	2.2689
5'	0.0015	40'	0.0116	7°	0.1222	60°	1.0472	140°	2.4435
6'	0.0017	50'	0.0145	8°	0.1396	70°	1.2217	150°	2.6180
7'	0.0020	1°	0.0175	9°	0.1571	80°	1.3963	160°	2.7925
8'	0.0023	2°	0.0349	10°	0.1745	90°	1.5708	170°	2.9671



## NATURAL TRIGONOMETRIC FUNCTIONS.

Angle.	Sin.	Csc.	Tan.	Ctn.	Sec.	Cos.	
0°	0.000	∞	0.000	∞	1.000	1.000	90°
1	0.017	57.30	0.017	57.29	1.000	1.000	89
2	0.035	28.65	0.035	28.64	1.001	0.999	88
3	0.052	19.11	0.052	19.08	1.001	0.999	87
4	0.070	14.34	0.070	14.30	1.002	0.998	86
5°	0.087	11.47	0.087	11.43	1.004	0.996	85°
6	0.105	9.567	0.105	9.514	1.006	0.995	84
7	0.122	8.206	0.123	8.144	1.008	0.993	83
8	0.139	7.185	0.141	7.115	1.010	0.990	82
9	0.156	6.392	0.158	6.314	1.012	0.988	81
10°	0.174	5.759	0.176	5.671	1.015	0.985	80°
11	0.191	5.241	0.194	5.145	1.019	0.982	79
12	0.208	4.810	0.213	4.705	1.022	0.978	78
13	0.225	4.445	0.231	4.331	1.026	0.974	77
14	0.242	4.134	0.249	4.011	1.031	0.970	76
15°	0.259	3.864	0.268	3.732	1.035	0.966	75°
16	0.276	3.628	0.287	3.487	1.040	0.961	74
17	0.292	3.420	0.306	3.271	1.046	0.956	73
18	0.309	3.236	0.325	3.078	1.051	0.951	72
19	0.326	3.072	0.344	2.904	1.058	0.946	71
20°	0.342	2.924	0.364	2.747	1.064	0.940	70°
21	0.358	2.790	0.384	2.605	1.071	0.934	69
22	0.375	2.669	0.404	2.475	1.079	0.927	68
23	0.391	2.559	0.424	2.356	1.086	0.921	67
24	0.407	2.459	0.445	2.246	1.095	0.914	66
25°	0.423	2.366	0.466	2.145	1.103	0.906	65°
26	0.438	2.281	0.488	2.050	1.113	0.899	64
27	0.454	2.203	0.510	1.963	1.122	0.891	63
28	0.469	2.130	0.532	1.881	1.133	0.883	62
29	0.485	2.063	0.554	1.804	1.143	0.875	61
30°	0.500	2.000	0.577	1.732	1.155	0.866	60°
31	0.515	1.942	0.601	1.664	1.167	0.857	59
32	0.530	1.887	0.625	1.600	1.179	0.848	58
33	0.545	1.836	0.649	1.540	1.192	0.839	57
34	0.559	1.788	0.675	1.483	1.206	0.829	56
35°	0.574	1.743	0.700	1.428	1.221	0.819	55°
36	0.588	1.701	0.727	1.376	1.236	0.809	54
37	0.602	1.662	0.754	1.327	1.252	0.799	53
38	0.616	1.624	0.781	1.280	1.269	0.788	52
39	0.629	1.589	0.810	1.235	1.287	0.777	51
40°	0.643	1.556	0.839	1.192	1.305	0.766	50°
41	0.656	1.524	0.869	1.150	1.325	0.755	49
42	0.669	1.494	0.900	1.111	1.346	0.743	48
43	0.682	1.466	0.933	1.072	1.367	0.731	47
44	0.695	1.440	0.966	1.036	1.390	0.719	46
45°	0.707	1.414	1.000	1.000	1.414	0.707	45°
	Cos.	Sec.	Ctn.	Tan.	Csc.	Sin.	Angle.

Values of the Complete Elliptic Integrals,  $K$  and  $E$ , for Different Values of the Modulus,  $k$ .

$\sin^{-1}k$	$K$	$E$	$\sin^{-1}k$	$K$	$E$	$\sin^{-1}k$	$K$	$E$
0°	1.5708	1.5708	30°	1.6858	1.4675	60°	2.1565	1.2111
1°	1.5709	1.5707	31°	1.6941	1.4608	61°	2.1842	1.2015
2°	1.5713	1.5703	32°	1.7028	1.4539	62°	2.2132	1.1920
3°	1.5719	1.5697	33°	1.7119	1.4469	63°	2.2435	1.1826
4°	1.5727	1.5689	34°	1.7214	1.4397	64°	2.2754	1.1732
5°	1.5738	1.5678	35°	1.7312	1.4223	65°	2.3088	1.1638
6°	1.5711	1.5665	36°	1.7415	1.4248	66°	2.3439	1.1545
7°	1.5767	1.5649	37°	1.7522	1.4171	67°	2.3809	1.1453
8°	1.5785	1.5632	38°	1.7633	1.4092	68°	2.4198	1.1362
9°	1.5805	1.5611	39°	1.7748	1.4013	69°	2.4610	1.1272
10°	1.5828	1.5589	40°	1.7868	1.3931	70°	2.5046	1.1184
11°	1.5854	1.5564	41°	1.7992	1.3849	71°	2.5507	1.1096
12°	1.5882	1.5537	42°	1.8122	1.3765	72°	2.5998	1.1011
13°	1.5913	1.5507	43°	1.8256	1.3680	73°	2.6521	1.0927
14°	1.5946	1.5476	44°	1.8396	1.3594	74°	2.7081	1.0844
15°	1.5981	1.5442	45°	1.8541	1.3506	75°	2.7681	1.0764
16°	1.6020	1.5405	46°	1.8691	1.3418	76°	2.8327	1.0686
17°	1.6061	1.5367	47°	1.8848	1.3329	77°	2.9026	1.0611
18°	1.6105	1.5326	48°	1.9011	1.3238	78°	2.9786	1.0538
19°	1.6151	1.5283	49°	1.9180	1.3147	79°	3.0617	1.0468
20°	1.6200	1.5238	50°	1.9356	1.3055	80°	3.1534	1.0401
21°	1.6252	1.5191	51°	1.9539	1.2963	81°	3.2553	1.0338
22°	1.6307	1.5141	52°	1.9729	1.2870	82°	3.3699	1.0278
23°	1.6365	1.5090	53°	1.9927	1.2776	83°	3.5004	1.0223
24°	1.6426	1.5037	54°	2.0133	1.2681	84°	3.6519	1.0172
25°	1.6490	1.4981	55°	2.0347	1.2587	85°	3.8317	1.0127
26°	1.6557	1.4924	56°	2.0571	1.2492	86°	4.0528	1.0087
27°	1.6627	1.4864	57°	2.0804	1.2397	87°	4.3387	1.0052
28°	1.6701	1.4803	58°	2.1047	1.2301	88°	4.7427	1.0021
29°	1.6777	1.4740	59°	2.1300	1.2206	89°	5.4349	1.0000

The Common Logarithms of  $\Gamma(n)$  for Values of  $n$  between 1 and 2.

$n$	$\log_{10} \Gamma(n)$	$n$	$\log_{10} \Gamma(n)$	$n$	$\log_{10} \Gamma(n)$	$n$	$\log_{10} \Gamma(n)$	$n$	$\log_{10} \Gamma(n)$
1.01	1.9975	1.21	1.9617	1.41	1.9478	1.61	1.9517	1.81	1.9704
1.02	1.9951	1.22	1.9605	1.42	1.9476	1.62	1.9523	1.82	1.9717
1.03	1.9928	1.23	1.9594	1.43	1.9475	1.63	1.9529	1.83	1.9730
1.04	1.9905	1.24	1.9583	1.44	1.9473	1.64	1.9536	1.84	1.9743
1.05	1.9883	1.25	1.9573	1.45	1.9473	1.65	1.9543	1.85	1.9757
1.06	1.9862	1.26	1.9564	1.46	1.9472	1.66	1.9550	1.86	1.9771
1.07	1.9841	1.27	1.9554	1.47	1.9473	1.67	1.9558	1.87	1.9786
1.08	1.9821	1.28	1.9546	1.48	1.9473	1.68	1.9566	1.88	1.9800
1.09	1.9802	1.29	1.9538	1.49	1.9474	1.69	1.9575	1.89	1.9815
1.10	1.9783	1.30	1.9530	1.50	1.9475	1.70	1.9584	1.90	1.9831
1.11	1.9765	1.31	1.9523	1.51	1.9477	1.71	1.9593	1.91	1.9846
1.12	1.9748	1.32	1.9516	1.52	1.9479	1.72	1.9603	1.92	1.9862
1.13	1.9731	1.33	1.9510	1.53	1.9482	1.73	1.9613	1.93	1.9878
1.14	1.9715	1.34	1.9505	1.54	1.9485	1.74	1.9623	1.94	1.9895
1.15	1.9699	1.35	1.9500	1.55	1.9488	1.75	1.9633	1.95	1.9912
1.16	1.9684	1.36	1.9495	1.56	1.9492	1.76	1.9644	1.96	1.9929
1.17	1.9669	1.37	1.9491	1.57	1.9496	1.77	1.9656	1.97	1.9946
1.18	1.9655	1.38	1.9487	1.58	1.9501	1.78	1.9667	1.98	1.9964
1.19	1.9642	1.39	1.9483	1.59	1.9506	1.79	1.9679	1.99	1.9982
1.20	1.9629	1.40	1.9481	1.60	1.9511	1.80	1.9691	2.00	0.0000



To avoid fine, this book should be returned on  
or before the date last stamped below

SON-12-50-26433

--	--	--





**MATHEMATICAL  
SCIENCES  
LIBRARY**



~~571,3  
13983  
202~~

Stanford University Libraries



3 6105 002 057 391

**STANFORD UNIVERSITY LIBRARY**  
Stanford, California

<p><del>_____</del> <del>_____</del> <del>_____</del></p> <p>JUN 05 2002</p>			
--	--	--	--

